


REGIOSPECIFIC PREPARATION OF 2-OXAZOLINES PROMOTED BY SILICON TETRAFLUORIDE

Makoto Shimizu* and Hirosuke Yoshioka*

The Institute of Physical and Chemical Research (RIKEN) Wako, Saitama 351-01, Japan

Abstract- Silicon tetrafluoride promotes reaction of oxiranes with nitriles to give 2-oxazolines in good yield.

2-Oxazolines have received considerable attentions as excellent synthetic intermediates, *e. g.*, masked carboxylic acids, and possess diverse utilities for industrial applications.¹ A variety of synthetic procedures have been reported until now, and among them the reaction of epoxides with nitriles constitutes the most straightforward and attractive approach. However, the only procedure reported for that employs sulfuric acid at 0 °C --room temp as a promoter, and the yields of 2-oxazolines were generally less than 20 %.² We report herein that SiF₄³ promotes reaction of oxiranes with nitriles effectively to form 2-oxazolines in good yields with high regioselectivities.



Stirring a solution of styrene oxide (0.5 mmol) in acetonitrile (3 ml) under atmosphere of SiF₄ at room temp. for 1 h. gave 4-phenyl-2-oxazoline in 80 % yield.⁴ A variety of 2-oxazolines was prepared under similar conditions, and Table 1 summarizes the results. As shown, 2-oxazolines were formed regioselectively, *i. e.*, the nitrogen atom was introduced into the more substituted carbons. The reaction proceeds generally in the nitrile as solvent under an excess of SiF₄. When a stoichiometric amount of nitrile was used, formation of a considerable amount of fluorohydrin³ was observed. Although an electron-withdrawing group on the oxirane ring decreased the reactivity (run 10), the 1-cyano-2-oxazoline was formed in good yield under the more forcing conditions (at 80 °C). Furthermore, the present reaction has an important chemoselectivity: mono-substituted aliphatic oxiranes did not undergo 2-oxazoline formation but the starting materials were


recovered almost unaffected.

Table 1. Preparation of 2-Oxazolines^a

Run	R ¹	R ²	R ³	R ⁴	Yield of 2 (%) ^b	
1	Ph	H	H	CH ₃	80	
2	C ₅ H ₁₁	C ₅ H ₁₁	H	CH ₃	65	
3	C ₅ H ₁₁	C ₅ H ₁₁	H	C ₂ H ₅	51	
4	C ₆ H ₁₃	CH ₃	H	CH ₃	52	
5	--(CH ₂) ₁₁ --		H	CH ₃	45	
6	CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ OBn	CH ₃	61
7	CH ₃	CH ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ OBn	C ₂ H ₅	63
8	CH ₃	--(CH ₂) ₄ --		CH ₃	33	
9	H	--(CH ₂) ₁₀ --		CH ₃	38	
10 ^c	C ₅ H ₁₁	C ₅ H ₁₁	CN	CH ₃	74	

^aThe reaction was carried out at 0 °C--room temp. for 1--12 h under an atmosphere of SiF₄. ^bAll new compounds were fully characterized by nmr, ir, mass spectra, and/or combustion analyses, and in some cases by direct comparison with the authentic samples prepared by the reported method.² Yields refer to isolated materials. ^cThe reaction was carried out at 80 °C.

The stereochemistry of the reaction was examined using 4-*t*-butylcyclohexane derivatives (**3** and **4**).⁵ The oxirane (**3**) gave **5** exclusively in 67 % yield, whereas **4** gave a mixture of isomers together with fluorohydrin (**7**), indicating that an S_N1 process was involved in the case of **4** probably due to a substantial torsional strain on an attack from the equatorial side.⁶



The oxazoline formation exercised in the present study shows an extended utility of SiF_4 as selective Lewis acid.⁷

REFERENCES AND NOTES

1. J. A. Frump, Chem. Rev., 1971, 71, 483; A. I. Meyers and E. D. Mihelick, Angew. Chem. Int. Ed. Engl., 1976, 15, 270; K. T. Potts in 'Comprehensive Heterocyclic Chemistry', vol. 5, ed. by A. R. Katritzky and C. W. Rees, Pergamon, Oxford 1984, pp. 112-166
2. R. Oda, M. Okano, S. Tokiura, and F. Misumi, Bull. Chem. Soc. Jpn., 1962, 35, 1219.
3. SiF_4 promotes opening of oxiranes to form fluorohydrins; M. Shimizu and H. Yoshioka, Tetrahedron Lett., in press.
4. Reactions were most conveniently run in an ordinary glassware apparatus fitted with a balloon filled with SiF_4 . After the completion of the reaction, an aqueous KF solution was added to the mixture, and normal workup followed by separation on SiO_2 TLC gave 2-oxazolines.
5. E. J. Corey and M. Chaykovsky, J. Am. Chem. Soc., 1965, 87, 1353.
6. The structures of 5, 6, and 7 were established based on their spectroscopic analysis. They afforded the following spectra: 5: ir(neat) 2950, 1670, 1475, 1440, 1385, 1235, 990, and 900 cm^{-1} ; pmr(CDCl_3) δ 0.88(9H, s), 1.20-1.86(9H, m), 1.96(3H, s), and 3.80(2H, s). 6: ir(neat) 2950, 1680, 1450, 1380, 1360, 1225, 990, 895, and 750 cm^{-1} ; pmr(CDCl_3) δ 0.86(9H, s), 1.16-1.88(9H, m), 1.96(3H, s), and 3.96(2H, s). 7: ir(neat) 3300, 2950, 1460, 1365, 1060, 880, and 735 cm^{-1} , pmr(CDCl_3) δ 0.88(9H, s), 1.00-2.12(10H, m, including an OH proton), 3.54(2H, d, $J=19.7$ Hz); ^{19}F nmr ($\text{CDCl}_3/\text{CFCl}_3$) -172 ppm Detailed explanation for the formation of fluorohydrin, see ref. 3.
7. SiF_4 is a good catalyst for glycosylation, see; S. Hashimoto, M. Hayashi, and R. Noyori, Tetrahedron Lett., 1984, 25, 1379.

Received, 24th June, 1988