

## Synthesis of (-)-3-Butyl-4-hydroxyphthalide



Yoshimitsu Ogawa\*, Kunio Hosaka, Masao Chin (Chen Zhengxiong) and  
Hiroshi Mitsuhashi

Research Institute for Biology & Chemistry, Tsumura & Co.


3586 Yoshiwara, Ami-machi, Inashiki-gun, Ibaraki 300-11, Japan

**Abstract** — (-)-3-Butyl-4-hydroxyphthalide 1 was first synthesized enantioselectively by using a chiral aryllithium reagent and its absolute stereochemistry at C-3 was determined to be S-configuration.

(-)-3-Butyl-4-hydroxyphthalide 1 was isolated from the rhizome of Ligusticum wallichii Franch (Japanese name 'senkyu') of which phenolic constituents was known to have the effect of increasing coronary flow<sup>1)</sup>. Although the structure of this compound was shown to possess structure 1, its stereochemistry hasn't been clear yet. In this paper we wish to describe the first total synthesis of (-)-1 in a high optical yield<sup>2)</sup> and the absolute stereochemistry of this phthalide.

1(-)-1

We considered two approaches for the asymmetric synthesis of (-)-1 as shown in Scheme 1. In the first approach(A) a chiral center may be introduced by the asymmetric reduction of the ketone 2 with chiral reductant. In the second approach(B), the asymmetric induction would be performed by the reaction of chiral aryllithium reagent 4 with aldehyde, followed by oxidation.




R=Alkyl group  
Z\*=Chiral ortho-directing group

Scheme 1

It is known that asymmetric reduction of the prochiral aryl ketone with (S)-BINAL-H 5 reported by Noyori et al.<sup>3)</sup> provides an alcohol of S-configuration with high enantioselectivity. The asymmetric synthesis of (-)-1 was examined as shown in Scheme 2.

The reduction of the ester 6<sup>4)</sup> with (S)-BINAL-H afforded (-)-3-butylphthalide 7 containing S-configuration<sup>5)</sup> at C-3 in 83% ee<sup>2)</sup>. On the other hand, in the case of ester 8<sup>6)</sup> no (S)-3-butyl-4-methoxyphthalide 9 was obtained. This result suggested that fairly bulky (S)-BINAL-H 5 couldn't approach to the acyl group by steric interference with two ortho-substituents in the ester 8.




Scheme 2

Previously the asymmetric synthesis of (S)-(-)-7 was accomplished via the lithiation with the halogen-metal exchange of chiral aminal obtained from o-bromobenzaldehyde with (S)-2-(anilinomethyl)pyrrolidine 10 by Asami and Mukaiyama<sup>7,8)</sup>. As the regioselective directed lithiation of aromatic compounds by the chelation effect was well known<sup>9)</sup>, the directed ortho lithiation of the

chiral aminal 12 containing a methoxy group at m-position was examined as shown in Scheme 3.

The chiral aminal 12 was easily obtained by condensation of m-methoxybenzaldehyde 11 with 10. Then the directed ortho lithiation of the aminal 12 with n-BuLi, followed by reaction with valeraldehyde and then hydrolysis under mild conditions afforded the optically active lactol (-)-13. The desired compound (-)-1 was obtained in 84% ee<sup>2)</sup> by the demethylation<sup>10)</sup> of (-)-9 obtained by the oxidation of the lactol 13 with Ag<sub>2</sub>O<sup>9)</sup>.



Scheme 3

It was speculated that (-)-1 has S-configuration at C-3 according to the mechanism proposed by Asami and Mukaiyama<sup>7a)</sup>. That is, the directed ortho lithiation of aminal first occurred to give a rigid tricyclic five membered ring (Figure 1) and then the aldehyde approaches from the less hindered front side in the manner shown in Figure 2 to avoid steric repulsion between the alkyl group of the aldehyde and the pyrrolidine ring illustrated with arrow in Figure 3. Further, the phthalide (-)-1 was converted to 7 via the triflate 14<sup>7b)</sup> to confirm the absolute configuration of (-)-1. The optical rotation of the resultant phthalide 7 was identical with that of the authentic phthalide (-)-7. Consequently, it was possible to determine the absolute stereochemistry of 1 to be S-configuration at C-3 position.




Figure 1




Figure 2




Figure 3

#### EXPERIMENTAL

Melting points were measured on a YANACO micro melting point apparatus and are uncorrected. Infrared spectra were recorded on a Hitachi 270-300 spectrophotometer. Nuclear magnetic resonance spectra were obtained with a JEOL FX-200 spectrometer using tetramethylsilane as an internal standard. Mass spectra were determined on a JEOL DX-300 spectrometer. Optical rotations were taken on a JASCO DIP-360 using a 100mm cell.

#### Methyl 2-valerylbenzoate 6

The esterification<sup>4b)</sup> of 2-valerylbenzoic acid<sup>4a)</sup> prepared from phthalic anhydride with dibutylcadmium gave methyl 2-valerylbenzoate 6:  $\nu_{\text{max}}$  1726, 1700  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  0.94 (3H, t,  $J=7.3\text{Hz}$ ), 1.40 (2H, tq,  $J=7.3, 7.3\text{Hz}$ ), 1.72 (2H, tt,  $J=7.5, 7.3\text{Hz}$ ), 2.80 (2H, t,  $J=7.5\text{Hz}$ ), 7.35 (1H, dd,  $J=7.3, 1.5\text{Hz}$ ), 7.48 (1H, ddd,  $J=7.3, 7.3, 1.5\text{Hz}$ ), 7.56 (1H, ddd,  $J=7.3, 7.3, 1.5\text{Hz}$ ), 7.89 (1H, dd,  $J=7.3, 1.5\text{Hz}$ );  $m/z$  220 ( $M^+$ ), 163 ( $M-C_4H_9$ ).

#### Methyl 3-methoxy-2-valerylbenzoate 8

A solution of diazomethane in ether was gradually added to a stirred solution of 3-methoxy-2-valerylbenzoic acid<sup>5)</sup> (2.0g, 8.46mmol) in ether at 0 °C. The mixture

was continued to stir at 0 °C for 1 h, then at room temperature for 1 h and evaporated to give a residue. The residue was purified by flash column chromatography (SiO<sub>2</sub>; eluent, AcOEt:n-hexane=1:3) to give a colorless oil of methyl 3-methoxy-2-valerylbenzoate 8 (1.65g, 78%) : ir (KBr)  $\nu_{max}$  1726, 1466, 1280, 1060 cm<sup>-1</sup>; <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  0.94 (3H, t, J=7Hz), 1.12-1.95 (4H, m), 2.81 (2H, t, J=7Hz), 3.81 (3H, s), 3.84 (3H, s), 7.09 (1H, dd, J=8, 1Hz), 7.39 (1H, dd, J=8, 8Hz), 7.57 (1H, dd, J=8, 1Hz); ms m/z 250 (M<sup>+</sup>), 193 (M-C<sub>4</sub>H<sub>9</sub>).

(S)-(-)-3-Butylphthalide 7

A solution of (S)-BINAL-H 5 (0.4M in THF) was prepared in situ from LiAlH<sub>4</sub> (39mg, 1.03mmol), abs.EtOH (1.05mmol) and (S)-(-)-2,2'-dihydroxy-1,1'-binaphthyl<sup>11)</sup> (300mg, 1.05mmol) under argon atmosphere<sup>3)</sup>. 1.0M THF solution of methyl 2-valerylbenzoate 6 (68mg, 0.31mmol) was injected into the stirred solution of (S)-BINAL-H 5 at -80 °C. The reaction mixture was stirred at this temperature for 3 h and quenched by addition of 2N HCl at -80 °C and then extracted with ether 2 times. The combined organic layer was washed with brine, dried (MgSO<sub>4</sub>) and evaporated to a residue. After (S)-(-)-2,2'-dihydroxy-1,1'-binaphthyl was removed from the residue by recrystallisation (n-hexane-CHCl<sub>3</sub>), the residue was purified by flash column chromatography (SiO<sub>2</sub>; eluent, AcOEt:CHCl<sub>3</sub>=2:3) to give a colorless oil of (S)-(-)-3-butylphthalide 7 (18mg, 30%, optical yield 83%<sup>ee</sup>) : [ $\alpha$ ]<sub>D</sub> -49.3° (c=0.4, CHCl<sub>3</sub>), [lit.<sup>12)</sup> [ $\alpha$ ]<sub>D</sub> -59.5° (c=0.2, CHCl<sub>3</sub>)] ; ir (KBr)  $\nu_{max}$  1764 cm<sup>-1</sup>; <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  0.91 (1H, t, J=7.1Hz), 1.20-2.31 (6H, m), 5.48 (1H, dd, J=7.8, 4.2Hz), 7.44 (1H, dd, J=7.4, 1.0Hz), 7.52 (1H, dd, J=7.6, 7.4Hz), 7.67 (1H, ddd, J=7.6, 7.6, 1.0Hz), 7.93 (1H, d, J=7.6Hz); ms m/z 190 (M<sup>+</sup>), 133 (M-C<sub>4</sub>H<sub>9</sub>).

3-(3-Methoxyphenyl)-2-phenyl-1,5,6,7-tetrahydro-3H-pyrro[1,2-c]imidazole 12

A solution of m-methoxybenzaldehyde 11 (2.0g, 14.69mmol) and (S)-(+)-2-(anilinomethyl)pyrrolidine 10<sup>13)</sup> (2.59g, 14.69mmol) in benzene (20ml) was refluxed for removal of water azeotropically under argon atmosphere for 3 h. The mixture was evaporated and then recrystallized from ether to give 3-(3-methoxyphenyl)-2-phenyl-1,5,6,7-tetrahydro-3H-pyrro[1,2-c]imidazole 12 (3.95g, 91%) : mp 78-80°C; [ $\alpha$ ]<sub>D</sub> +38.5° (c=0.3, CH<sub>2</sub>Cl<sub>2</sub>); ir (KBr)  $\nu_{max}$  1602, 1504, 1368, 1278, 1042; <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  1.70-2.20 (4H, m), 2.78 (1H, dd, J=18, 9Hz),

3.21 (1H, dd, J=9, 9Hz), 3.33 (1H, m), 3.72 (1H, dd, J=7, 7Hz), 3.76 (3H, s), 3.89 (1H, m), 5.28 (1H, s), 6.45 (2H, d, J=8Hz), 6.65 (1H, dd, J=7, 7Hz), 6.78 (1H, dd, J=8, 2Hz), 6.88 (1H, d, J=2Hz), 6.91 (1H, d, J=8Hz), 7.12 (1H, d, J=7Hz), 7.18 (2H, dd, J=8, 8Hz); ms m/z 294 (M<sup>+</sup>), 189.

(-)-3-Butyl-1-hydroxy-4-methoxy-2-oxaindane 13

A solution of n-BuLi (1.6M in hexane, 2.13ml) was injected into a stirred solution of (-)-3-(3-methoxyphenyl)-2-phenyl-1,5,6,7-tetrahydro-3H-pyrro[1,2-c]imidazole 12 (1.0g, 3.40mmol) in dry ether (10ml) under argon atmosphere at room temperature. The mixture was stirred at room temperature for 4 h and then a solution of valeraldehyde (0.54ml, 5.10mmol) in dry ether (2ml) was injected at -80 °C. The mixture was stirred at -80 °C for 3 h and quenched with sat. NH<sub>4</sub>Cl at this temperature. The ethereal layer was hydrolyzed with 2% HCl at 0 °C for 1 h and then was extracted with ether 2 times. The ethereal layer was washed with water, dried (MgSO<sub>4</sub>) and evaporated to give a residue. The residue was purified by flash column chromatography (SiO<sub>2</sub>; eluent, AcOEt:benzene=1:5) to give a colorless oil of (-)-3-butyl-1-hydroxy-4-methoxy-2-oxaindane 13 (250mg, 33%, a mixture of diastereoisomers): [α]<sub>D</sub> -45.9° (c=0.1, CHCl<sub>3</sub>); ir (KBr) ν<sub>max</sub> 3416, 1604, 1486, 1266 cm<sup>-1</sup>; <sup>1</sup>H-nmr (CDCl<sub>3</sub>) δ 0.88 and 0.90 (3H, t, J=7.3Hz and t, J=7.3Hz, respectively), 1.10-1.55 (4H, m), 1.55-1.80 (1H, m), 1.90-2.15 (1H, m), 3.52 and 3.58 (1H, d, J=8.3 and d, J=7.6Hz, respectively, D<sub>2</sub>O exchangeable), 3.82 and 3.83 (3H, s and s), 5.22 and 5.49 (1H, dd, J=7.8, 2.9Hz and m, respectively), 6.36 and 6.46 (1H, d, J=7.6Hz and m, respectively), 6.82 (1H, d, J=7.8Hz), 7.00 (1H, d, J=7.3Hz), 7.30 (1H, dd, J=7.8, 7.3Hz); ms m/z 222 (M<sup>+</sup>), 166.

(-)-3-Butyl-4-hydroxyphthalide 1

A solution of (-)-3-butyl-1-hydroxy-4-methoxy-2-oxaindane 13 (137mg, 0.61mmol) in MeOH (2.5ml) was added to a stirred solution of Ag<sub>2</sub>O<sup>9</sup> (667mg, 2.88mmol) in H<sub>2</sub>O (3ml) -MeOH (0.5ml) at room temperature. The mixture was stirred for 1 h and filtered through a celite. The filtrate was evaporated to give a residue. After 2N H<sub>2</sub>SO<sub>4</sub> (3ml) was added to the residue at 0 °C, the mixture was extracted with ether 2 times. The ether layer was washed with brine, dried (MgSO<sub>4</sub>) and evaporated to give a residue. The residue was purified by flash column

chromathography (SiO<sub>2</sub> ; eluent, AcOEt:n-hexane=1:1) to give (-)-3-butyl-4-methoxyphthalide 9 (103mg, 76%) :  $[\alpha]_D -69.3^\circ$  (c=0.2, CHCl<sub>3</sub>) ; ir (KBr)  $\nu_{max}$  1770, 1492, 1274 ; <sup>1</sup>H-nmr (Acetone-d<sub>6</sub>)  $\delta$  0.88 (3H, t, J=6.8Hz), 1.10-1.45 (4H, m), 1.57-1.83 (1H, m), 2.10-2.37 (1H, m), 3.97 (3H, s), 5.53 (1H, dd, J=7.6, 3.1Hz), 7.32 (1H, d, J=7.8Hz), 7.38 (1H, d, J=7.8Hz), 7.56 (1H, dd, J=7.8, 7.8Hz) ; ms m/z 220 (M<sup>+</sup>), 163 (M-C<sub>4</sub>H<sub>9</sub>).

A solution of BBr<sub>3</sub><sup>10</sup> (0.8M in CH<sub>2</sub>Cl<sub>2</sub>, 0.9ml) was added to a solution of (-)-3-butyl-4-methoxyphthalide 9 (75mg, 0.34mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (1ml) at -30 °C. The mixture was stirred at -30 °C for 1 h and then at room temperature for 2 h. The mixture was poured into ice-water and extracted with CHCl<sub>3</sub> 2 times. The combined organic layer was washed with brine, dried (MgSO<sub>4</sub>) and evaporated to give a residue. The residue was recrystallized from benzene to give a colorless needle of 3-butyl-4-hydroxyphthalide 1 (61mg, 87%, optical yield 84%ee) : mp 188-190°C ;  $[\alpha]_D -88.7^\circ$  (c=0.38, EtOH), [lit.<sup>11</sup>  $[\alpha]_D -105.5^\circ$  (EtOH) ] ; ir (KBr)  $\nu_{max}$  3224, 1720 ; <sup>1</sup>H-nmr (Acetone-d<sub>6</sub>)  $\delta$  0.89 (3H, t, J=7.3Hz), 1.20-1.50 (4H, m), 1.65-1.90 (1H, m), 2.15-2.40 (1H, m), 5.57 (1H, dd, J=7.6, 2.9Hz), 7.18 (1H, dd, J=7.8, 1.0Hz), 7.31 (1H, dd, J=7.6, 1.0Hz), 7.42 (1H, dd, J=7.8, 7.6Hz) ; ms m/z 206 (M<sup>+</sup>), 149 (M-C<sub>4</sub>H<sub>9</sub>).

#### Conversion of (-)-1 to (-)-7

Trifluoromethanesulfonic anhydride (0.2ml, 1.20mmol) was added to a solution of (-)-3-butyl-4-hydroxyphthalide (206mg, 1.00mmol) and dry pyridine (0.24ml, 3.00mmol) in dry-CH<sub>2</sub>Cl<sub>2</sub> (1ml) at 0 °C. The mixture was stirred at 0 °C for 10min and then at room temperature for 1h. The mixture was poured into iced 2% HCl and extracted with ether. The organic layer was washed with water 2 times and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to give a yellow oil of 3-butyl-4-trifluoromethanesulfonyloxyphthalide 14 (283mg, 84%) : <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  0.91 (3H, t, J=7.1Hz), 1.15-1.50 (4H, m), 1.65-1.90 (1H, m), 2.10-2.35 (1H, m), 5.70 (1H, dd, J=8.1, 3.0Hz), 7.58 (1H, dd, J=8.1, 1.2Hz), 7.66 (1H, dd, J=8.1, 7.3Hz), 7.95 (1H, dd, J=7.3, 1.2Hz). Without purification of 14, formic acid (0.06ml, 1.67mmol) was added to a solution of the triflate 14 (283mg, 0.84mmol), bis(triphenylphosphine)-palladium(II) chloride (30mg, 0.04mmol) and tributylamine (0.6ml, 2.51mmol) in dry DMF (1.8ml) under argon atmosphere at room temperature<sup>12</sup>. The mixture was heated at 110 °C for 3h and then filtered. The filtrate was washed

with sat. NaHCO<sub>3</sub>, 2% HCl, water and brine, dried (MgSO<sub>4</sub>) and evaporated to give a residue. The residue was purified by flash column chromatography (SiO<sub>2</sub>; eluent, AcOEt:n-hexane=1:8) to give a colorless oil of (-)-3-butylphthalide 7 (128mg, 81%) :  $[\alpha]_D -49.1^\circ$  (c=0.2, CHCl<sub>3</sub>); the other spectral data was identical with those of the preceding (-)-7.

#### REFERENCES AND NOTES

- 1) Peking Institute of Pharmaceutical Industries, Yao Hsueh Hsueh Pao, 1979, 14, 670.
- 2) The optical yields (%) was calculated by the A/B ratio x100.  
A=Specific rotation of a synthesized product  
B=Specific rotation of a natural product
- 3) R.Noyori, I.Tomino, and Y.Tanimoto, J. Am. Chem. Soc., 1979, 101, 3129.
- 4) (a) T.Kariyone and S.Shimizu, Yakugaku Zasshi, 1953, 73, 336; P.L.De Benneville, J. Org. Chem., 1941, 6, 462.  
(b) G.E.Ullyot, H.W.Taylor, and N.Dawson, J. Am. Chem. Soc., 1948, 70, 542.
- 5) U.Nagai, T.Shishido, R.Chiba, and H.Mitsuhashi, Tetrahedron, 1965, 21, 1701; P.R.Jones and C.J.Jarboe, Tetrahedron Lett., 1969, 1849.
- 6) In submitting to Chem. Pharm. Bull..
- 7) (a) M.Asami and T.Mukaiyama, Chem. Lett., 1980, 17.  
(b) Q.Y.Chen, Y.B.He, and Z.Y.Yang, J.Chem.Soc., Chem. Commun., 1986, 1452.
- 8) (a) D.W.Slocum and C.A.Jennings, J. Org. Chem., 1976, 41, 3653.  
(b) A.I.Meyers and E.D.Mihelich, J. Org. Chem., 1975, 40, 3158; A.I.Meyers and W.B.Avila, ibid., 1981, 46, 3881; A.I.Meyers, M.A.Hanagan, L.M.Trefonas, and R.J.Baker, Tetrahedron, 1983, 39, 1991.
- 9) M.Rosenberger, D.Andrews, F.DiMaria, A.J.Duggan, and G.Saucy, Helv. Chim. Acta, 1972, 55, 249.
- 10) J.F.W.McOmie, M.L.Watts, and D.E.West, Tetrahedron, 1968, 24, 2289.
- 11) J.Jacques, C.Fouquey, and R.Viterbo, Tetrahedron Lett., 1971, 4617.
- 12) T.Yamagishi and H.Kaneshima, Yakugaku Zasshi, 1977, 97, 237.
- 13) T.Mukaiyama, M.Asami, J.Hanna, and S.Kobayashi, Chem. Lett., 1977, 783.
- 14) A.I.Meyers, D.L.Temple, D.Haidukewych, and E.D.Mihelich, J. Org. Chem., 1974, 39, 2787.

Received, 17th October, 1988