

BAKUCHICIN, A NEW SIMPLE FURANOCOUMARIN
FROM PSORALEA CORYLIFOLIA L.


Yoshikazu Kondo*, Atsushi Kato, Yoshiyuki Kubota,
and Shigeo Nozoe

Pharmaceutical Institute, Tohoku University
Aobayama, Aoba-ku, Sendai 980, Japan

Abstract — A new simple furanocoumarin, bakuchicin, has been isolated from the hexane-extract of seeds of Psoralea corylifolia L. (Leguminosae) alone with stigmasterol, psoralen and bakuchiol. The structure of bakuchicin has been shown as 8-oxo-8H-furo[2,3-f][1]benzopyran (1) by spectral means.

The seeds of Psoralea corylifolia L. (Leguminosae, Sanskrit name: Bakuchin) is used as a tonic or an aphrodisiac against impotence and menstruation disorders. The seed-oil of this plant also is used externally for the treatment of leucodermy, psoriasis, and leprosy in Indian folkloric remedy.¹ Many studies on the constituents of the seeds have been reported.² Now the presence of a new simple furanocoumarin was proved.

The *n*-hexane extract of the powdered seeds of P. corylifolia yielded an oily fraction which on silica gel chromatography gave a 0.1% of yield of a crystalline component (1), mp 138 °C, besides the known compounds, stigmasterol, psoralen (2), and bakuchiol (3). 1 is a new compound, for which we propose the name bakuchicin.



	δ (ppm)	J (Hz)
C2-H	7.70 (d)	2.0
C3-H	7.14 (dd)	2.0, 1.25
C4-H	4.44 (dd)	8.75, 1.25
C5-H	7.38 (d)	8.75
C6-H	7.82 (d)	9.5
C7-H	6.40 (d)	9.5





Table 1. ¹H Nmr spectral data of 1

Figure 1. ¹³C Nmr spectral data of 1



Bakuchicin (1) analyzed for $C_{11}H_6O_3$, a formula which confirmed by high-resolution mass spectrum molecular ion peak 186.0334 (Calcd 186.0317). The ultraviolet absorption spectrum (in 95% EtOH) showed maxima at 242 nm (infl.) ($\log \epsilon$ 4.47), 248 nm ($\log \epsilon$ 4.48) and 297 nm ($\log \epsilon$ 4.13), while the infrared spectrum (in $CHCl_3$) displayed bands at 1722 and 1619 cm^{-1} , suggesting the presence of a furano-coumarin skeleton. This suggestion is supported by spectral data of 1H nmr (Table 1), ^{13}C nmr (Figure 1), and 2D nmr.




Figure 2. Double resonance 1H nmr spectra of 1

The COSY spectrum of bakuchicin showed that the signal at δ 7.14 (1H, dd, $J=2.0$ Hz, 1.25 Hz) is correlated with the signals at δ 7.44 (1H, dd, $J=8.75$ Hz, 1.25 Hz) and δ 7.70 (1H, d, $J=2.0$ Hz), but no correlation between the signals δ 7.82 and other aromatic protons. These findings, combined with a double-resonance experiment (Figure 2), showed that another structure (1a) can be ruled out and the structure of bakuchicin is elucidated as 8-oxo-8H-furo[2,3-f][1]benzopyran (1). Interestingly, previous investigators have been reported isolation of angelicin (4) from seeds of *P. corylifolia*,³ however, we proved the presence of bakuchicin (1), a new isomeric simple furanocoumarin, in this paper.

EXPERIMENTAL

Melting points were taken on a Yamato MP-2 melting point apparatus and are uncorrected. Ultraviolet spectra were recorded with a Hitachi U-3200 spectrophotometer. Infrared spectra were determined as CHCl_3 solutions on a Jasco A-100S infrared spectrophotometer. ^1H Nmr and ^{13}C nmr spectra were recorded in CDCl_3 on a JEOL FX-500 spectrometer. TMS was used as an internal standard; chemical shifts are reported in δ ppm units. Mass spectra were determined with a JEOL DX-303 double focusing mass spectrometer operating at 70 eV.

Isolation of substances Dried and powdered seeds of *P. corylifolia* (1000 g) were exhaustively extracted with *n*-hexane (2 l x 4) at room temperature for 24 h. The *n*-hexane solution was evaporated and then subjected to column chromatography on silica gel (6 x 50 cm). Elution was accomplished with *n*-hexane and increasing quantities of AcOEt. Fractions of about 50 ml were collected and monitored by thin-layer chromatography (tlc). The four major compounds, bakuchiol 3 (52.127 g), stigmasterol (0.050 g), bakuchicin 1 (0.992 g), and psoralen 2 (0.884 g), were obtained from the corresponding fractions 10-16, 28-30, 31-35, and 37-41, respectively.

Bakuchiol (3) Colorless oil, bp 146-147 °C/0.8 mmHg. Ms m/z : 256 (M^+) (Calcd 256). The identity of this compound was established by comparison of ^1H nmr spectral data.⁴ **Bakuchiol 3,5-dinitrobenzoate**: Pale yellow plates (acetone-MeOH), mp 130 °C (lit.,⁴ mp 136 °C). **Anal.** Calcd for $\text{C}_{25}\text{H}_{26}\text{N}_2\text{O}_6$: C, 66.56; H, 5.82; N, 6.22. Found: C, 66.61; H, 5.97; N, 6.14. **Fdms** m/z : 450 (M^+).

Stigmasterol Colorless plates (MeOH), mp 170 °C. **Anal.** Calcd for $\text{C}_{29}\text{H}_{48}\text{O}$: C, 84.40; H, 11.72. Found: C, 84.59; H, 11.77. Ms m/z : 412 (M^+).

Bakuchicin (1) Colorless needles (acetone-*n*-hexane), mp 138 °C. **Anal.** Calcd for

$C_{11}H_6O_3$: C, 70.97; H, 3.25. Found: C, 70.97; H, 3.46. Ms m/z : 186.0334 (Calcd 186.0317). Ir ($CHCl_3$) cm^{-1} : 1722 (α, β -unsaturated lactone), 1619 (aromatic). Uv (95% EtOH) nm ($\log \epsilon$): 242 infl. (4.47), 248 (4.48), 297 (4.13). 1H Nmr: Table 1. ^{13}C Nmr: Figure 1.

Psoralen (2) Colorless needles (MeOH), mp 155 °C. Anal. Calcd for $C_{11}H_6O_3$: C, 70.97; H, 3.25. Found: C, 71.11; H, 3.29. Ms m/z : 186 (M^+). The identity of this compound with an authentic sample of psoralen was confirmed by comparison of 1H nmr, ir, uv spectra, and tlc behavior.

REFERENCES

1. E. F. Steinmetz and G. Lindermann, Riechstoffe, Aromen, Koerpepflegemittel, 1964, **14**, 409.
2. G. K. Gupta, J. L. Suri, B. K. Gupta, and K. L. Dhar, Phytochemistry, 1982, **21**, 2149, and references cited therein.
3. T. R. Seshadri and C. Venkatarao, Proc. Indian Acad. Sci., 1937, **5A**, 351.
4. G. Mehta, U. R. Nayak, and S. Dev, Tetrahedron, 1973, **29**, 1119.

Received, 15th November, 1989