

SYNTHESIS OF (+)-TRANS-WHISKY LACTONE, (+)-TRANS-COGNAC LACTONE AND
(+)-ELDANOLIDE


Takashi Ebata*, Katsuya Matsumoto, Hajime Yoshikoshi^{a)}, Koshi
Koseki, Hiroshi Kawakami, and Hajime Matsushita

Life Science Research Laboratory, Japan Tobacco Inc., 6-2,
Umeagaoka, Midori-ku, Yokohama, Kanagawa 227, Japan

a) Tokyo Research Laboratory, Yuki Gosei Kogyo Co., Ltd., 3-37-1,
Sakashita, Itabashi-ku, Tokyo 174, Japan

Abstract — A new and useful synthesis of (+)-trans-whisky
lactone (**3**), (+)-trans-cognac lactone (**4**) and (+)-eldanolide (**5**)
starting from levoglucosenone (**1**) is described.


Levoglucosenone (**1**)¹ (1,6-anhydro-3,4-dideoxy- β -D-glycero-hex-3-enopyranos-2-ulose) is widely known as a pyrolytic product of cellulose. It is a very useful chiral source for synthesizing natural products² because of its highly functionalized structure, which contains one chiral center. We have recently reported an efficient method for the convenient preparation of (S)-5-Hydroxypenten-4-olide (**2**) from levoglucosenone (**1**)³ by using the Baeyer-Villiger oxidation as a key step. (Scheme 1)



Scheme 1

In this paper, we describe a general method that permits easy access to 3,4-disubstituted γ -lactones, such as trans-whisky lactone (**3**) [(3*S*,4*R*)-3-methyl-4-octanolide], the key flavor of whisky and wine,^{4,5} trans-cognac lactone (**4**) [(3*S*,4*R*)-3-methyl-4-nonanolide], the key flavor of cognac,⁶ and the natural eldanolide (**5**) [(3*S*,4*R*)-3,7-dimethyl-6-octen-4-olide], the pheromone produced by the male *Eldana saccharina* (Wlk.),⁷ in an enantioselective manner using the chirality of levoglucosenone (**1**).

Our synthesis of **3**, **4** and **5** was straightforward, as illustrated in Scheme 2. Treatment of levoglucosenone (**1**) with Me_2CuLi gave **6** in 84.3 % yield ($[\alpha]_{\text{D}}^{23} -293^\circ$ (Et_2O), lit.^{2a} $[\alpha]_{\text{D}}^{23} -299^\circ$ (Et_2O)). It was shown to be 100 % diastereomerically pure when analyzed by glc and ^1H nmr. It was oxidized with AcOOH to give **7** in 86.0 % yield. The corresponding tosylate (**8**) was treated with K_2CO_3 in MeOH to give epoxide (**9**) in 66.4 % yield from **7**. Finally, treatment of **9** with $n\text{-Pr}_2\text{CuLi}$ gave trans-whisky lactone (**3**)⁸ in 76.2 % yield ($[\alpha]_{\text{D}}^{23} +79.5^\circ$ (MeOH), lit.^{9b} $[\alpha]_{\text{D}}^{20} +72.8^\circ$ (MeOH)). And similar treatment of **9** with $n\text{-Bu}_2\text{CuLi}$ gave trans-cognac lactone (**4**)¹⁰ in 78.3 % yield ($[\alpha]_{\text{D}}^{23} +79.5^\circ$ (CH_2Cl_2), lit.¹¹ $[\alpha]_{\text{D}}^{15} +48.3^\circ$ (CH_2Cl_2)). And further, eldanolide (**5**)¹² was also synthesized in a similar manner using $\text{Me}_2\text{C}=\text{CHMgBr}$ and CuBr in 73.8 % yield ($[\alpha]_{\text{D}}^{23} +57.8^\circ$ (EtOH), lit.^{13b} $[\alpha]_{\text{D}}^{21} +55.9^\circ$ (EtOH)).



Scheme 2

In conclusion, we developed a new and useful synthesis of (+)-trans-whisky lactone (3), (+)-trans-cognac lactone (4) and (+)-eldanolide (5) starting from levoglucosenone (1).

REFERENCES

1. This compound is available from Yuki Gosei Kogyo Co., Ltd.
2. a) M. Mori, T. Chuman, and K. Kato, *Carbohydr. Res.*, 1984, **129**, 73.
b) M. Isobe, N. Fukami, and T. Goto, *Chemistry Lett.*, 1985, 71.
3. K. Koseki, T. Ebata, H. Kawakami, H. Matsushita, Y. Naoi, and K. Itoh, *Heterocycles*, 1990, **31**, 423.
4. K. Otsuka, Y. Zenibayashi, M. Itoh, and A. Totsuka, *Agric. Biol. Chem.*, 1974, **38**, 485 and refs cited therein.
5. M. Masuda and K. Nishimura, *Chemistry Lett.*, 1981, 1333.
6. R. ter Heide, P. J. deValois, J. Visser, P. P. Jaegers, and R. Timmer, 'Analysis of Food and Beverages', ed. G. Charalambous, Academic Press, New York, 1978, p. 275.
7. G. Kunesch, P. Zagatti, J. Y. Lallemand, A. Debal, and J. P. Vigneron, *Tetrahedron Lett.*, 1981, **22**, 5271.
8. Physical data for our synthetic **3** are as follows; bp 123-125°C/16 torr; n_D^{23} 1.4402; ir (film) $\nu_{cm^{-1}}$ 2964, 2938, 1783, 1214, 1174; 1H -nmr (CDCl₃, 300 MHz): δ 0.92 (3H, t, J=7.2 Hz), 1.14 (3H, d, J=6.4 Hz), 1.30~1.75 (6H, m), 2.12~2.31 (2H, m), 2.60~2.75 (1H, m), 4.01 (1H, dt, J=4.0 and 7.7 Hz).
9. Optically active trans-whisky lactone (**3**) has been prepared;
 - a) C. Gunther and A. Mosandl, *Liebigs Ann. Chem.*, 1986, 2112
 - b) R. Bloch and L. Gilbert, *J. Org. Chem.*, 1987, **52**, 4603.
 - c) D. Hoppe and O. Zschage, *Angew. Chem. Int. Ed. Engl.*, 1989, **28**, 69.
 - d) M. Beckmann, H. Hildebrandt, and E. Winterfeldt, *Tetrahedron Asym.*, 1990, **1**, 1335.
10. Physical data for our synthetic **4** are as follows; bp 101-103°C/6 torr; n_D^{23} 1.4431; ir (film) $\nu_{cm^{-1}}$ 2962, 2936, 1779, 1212, 1172; 1H -nmr (CDCl₃, 300 MHz): δ 0.90 (3H, t, J=6.8 Hz), 1.14 (3H, d, J=6.3 Hz), 1.21~1.75 (8H, m), 2.12~2.30 (2H, m), 2.60~2.75 (1H, m), 4.01 (1H, dt, J=4.0 and 7.7 Hz).
11. Only one synthesis of optically active trans-cognac lactone (**4**) has been reported; R. M. Ortuno, R. Merce, and J. Font, *Tetrahedron*, 1987, **43**, 4497.

12. Physical data for our synthetic **5** are as follows; bp 115~117°C/21 torr; n_D^{23} 1.4606; ir (film) $\nu_{cm^{-1}}$ 2972, 2920, 1783, 1214, 1195; 1H -nmr (CDCl₃, 300 MHz): δ 1.14 (3H, d, J=6.5 Hz), 1.64 (3H, s), 1.73 (3H, s), 2.10~2.50 (4H, m), 2.68 (1H, dd, J=7.5 and 16.6 Hz), 4.06 (1H, dd, J=6.5 and 12.2 Hz), 5.10~5.22 (1H, m).
13. Optically active eldanolide (**5**) has been prepared;
- a) J. P. Vigneron, R. Meric, M. Larcheveque, A. Debal, G. Kunesch, P. Zagatti, and M. Gallois, Tetrahedron Lett., 1982, **23**, 5051.
 - b) T. Uematsu, T. Umemura, and K. Mori, Agric. Biol. Chem., 1983, **47**, 597.
 - c) Y. Yokoyama and M. Yunokihara, Chemistry Lett., 1983, 1245.
 - d) K. Suzuki, T. Ohkuma, and G. Tsuchihashi, Tetrahedron Lett., 1985, **26**, 861.
 - e) R. M. Ortuno, R. Merce, and J. Font, Tetrahedron, 1987, **43**, 4497.

Received, 22nd June, 1990