

ACCENTUATION OF THE DI- π -METHANE REACTIVITY BY CENTRAL CARBON
SUBSTITUTION IN THE 4-(PHENYLMETHYL)-2(5H)-FURANONE SYSTEM

Osamu Muraoka,* Genzoh Tanabe,# and Takefumi Momose##,*

Faculty of Pharmaceutical Sciences, Kinki University#

Kowakae 3-4-1, Higashi-osaka, Osaka 577, Japan and


Faculty of Pharmaceutical Sciences, Toyama Medical and

Pharmaceutical University,## Sugitani 2630, Toyama 930-01, Japan.

Abstract - The effect of 'central methane' substitution on the di- π -methane rearrangement in 4-(phenylmethyl)-2(5H)-furanones (1b-d) was investigated. Significant enhancement of efficiency in the reaction leading in high combined yields to two isomeric products (endo-2 and exo-2) was discussed in terms of both the substituent effects at the allylic methane carbon and the restrained feature of the ring-enrolled π -system.

There are three known versions of the di- π -methane rearrangement:¹ the divinyl methane variety, the arylvinylmethane type and the oxa-di- π -methane^{1c,1d} variation. These differ in many aspects. The marked differences between the first two are as follows: the divinylmethane rearrangement requires the 'methane carbon' substitution while the arylvinyl type does not;² the arylvinylmethane singlet tends to react more slowly than does its divinyl counterpart, and substituent effects are reversed for these two cases.²


In a previous paper³ dealing with photo-reactivity of 4-(phenylmethyl)-2(5H)-furanone (1a), an arylvinylmethane example of the di- π -methane rearrangement, we obtained



an expected photo-rearranged product (2a) in a poor yield (~23%). It was of considerable interest to study the photochemical behavior of the system with substituents on the central methano-carbon from a mechanistic standpoint in order to determine and understand the difference between these two versions of the di- π -methane rearrangement. Thus we synthesized some 'central methane'-substituted

4-(phenylmethyl)-2(5*H*)-furanones (1b-d), and investigated their photochemical reactivities and found significant enhancement of efficiency of the reaction under singlet and acetone-photosensitized conditions.

The reactants (1b-d) were prepared as follows. The selective reduction of 1-ethyl 4-hydrogen succinates (3b-d)⁴ with lithium aluminum hydride at 0°C, and subsequent acid-catalyzed lactonization of the resulting hydroxy acids gave 3-alkylidenedihydro-2(3*H*)-furanones (4b-d), which were isomerized into the desired 2(5*H*)-furanones (1b-d) in 62-82% yields by the action of tosic acid in dimethyl sulfoxide at 150°C.⁵



Direct irradiation⁶ of (1b-d) in methanol through a Pyrex filter gave two isomeric photo-rearranged products, *endo*- and *exo*-6-substituted 5-phenyl-3-oxabicyclo[3.1.0]hexan-2-ones (*endo-2* and *exo-2*).^{7,8} The configuration of the photo-products was determined on examination of the vicinal coupling constants across the cyclopropane ring in the ¹H-nmr spectra.⁹ The acetone-sensitized irradiation of 1b-d through a Pyrex filter also resulted in the formation of the same members of products as those obtained in the run carried out in methanol. The results of the irradiation are summarized in Table 1.

The efficiency of the rearrangement was greatly increased by the substitution on the central carbon. The substituent effect on central methane has mostly been examined





Table 1

| R | solvent | reaction time(hr) | <i>endo-2</i> isolated yield (%) | <i>exo-2</i> yield (%) |
|-----------------|----------|-------------------|----------------------------------|------------------------|
| Ph | acetone | 4 | 70 | 15 |
| | methanol | 11 | 70 | 13 |
| c-Hex | acetone | 8 | 85 | 8 |
| | methanol | 11 | 74 | 19 |
| Me ¹ | acetone | 6 | 90 | trace |
| | methanol | 11 | 70 | 7 |
| H ² | acetone | 3 | 47 (recovered 7%) | |
| | methanol | 12 | 39 (recovered 33%) | |

1. The product distribution was determined by 500 MHz ¹H-nmr spectra.
2. The yields were optimized in the present study.

on disubstituted substrates.^{2,10} Present findings reveal that the monosubstitution also exerts the effect in a similar fashion and enhances the efficiency of the di- π -methane rearrangement to a large extent irrespectively of substituent types. Most interestingly, triplet sensitization in an acetone solution led to highly efficient formation of the cyclopropane. Hammond¹¹ reported the photoirradiation of a 4,4-dimethylated 4-phenylbutenoate (**5**), an acyclic version of functionality similar to **1**, where the di- π -methane reaction to yield **6** could only be effected upon direct irradiation, but not on acetone-photosensitized irradiation. Hixon¹² rationalized the non-occurrence of the di- π -methane reaction of the triplet excited state of **5** in terms of preferential dissipation of the triplet excitation energy by "free-rotor" effects (facile *E-Z* isomerization)¹³ about the double bond. The efficiency of the rearrangement in the present study should be ascribed to both the substituent effects on the 'central methane' carbon in favor of its facile migration *via* radical fission and the inherent feature of the 2(5H)-furanone system where this alternative photo-process (*E-Z* isomerization) is precluded by incorporating the double bond in a furanone ring. In addition, the relatively consistent product ratios and increased efficiency of the reaction in the triplet-sensitized conditions are thought to indicate that the products resulted from the triplet state. The prolonged reaction periods observed in direct irradiations would be responsible for low efficiency of the intersystem crossing in methanol.



REFERENCES AND NOTES

1. a) S. S. Hixon, P. S. Mariano, and H. E. Zimmerman, *Chem. Rev.*, 1973, 73, 531;
 b) H. E. Zimmerman, 'Rearrangements in Ground and Excited States', ed. P. De Mayo, Academic Press, Inc., New York, 1980, Vol. 3, p. 131; c) R. S. Givens and W. F. Oettle, *Chem. Commun.*, 1969, 1184; d) W. G. Dauben, M. S. Kellogg, J. I. Seeman, and W. A. Spitzer, *J. Am. Chem. Soc.*, 1970, 92, 1786.
2. H. E. Zimmerman, M. G. Steinmetz, and C. L. Kreil, *J. Am. Chem. Soc.*, 1978, 100, 4146.
3. T. Momose, G. Tanabe, H. Tsujimori, M. Higashiura, I. Imanishi, and K. Kanai, *Heterocycles*, 1989, 29, 257.
4. Compounds 3b and 3d were prepared according to the described method (G. H. Daub and W. S. Johnson, *J. Am. Chem. Soc.*, 1948, 70, 418). Compound 3c was prepared from cyclohexyl phenyl ketone in the same manner as that described for the preparation of 3b and 3d.
5. The acid-catalyzed *exo* → *endo* isomerization of the double bond in 2-benzylidenecycloalkanones has been reported. Cf, A. Hassner and T. C. Mead, *Tetrahedron*, 1964, 20, 2201.
6. The photochemical reaction was carried out in an immersion apparatus fitted with an Ishii UV-HT 200 W high-pressure mercury lamp.
7. Independent synthesis of endo-2b and exo-2b was reported: T. E. Nalesnik, J. G. Fish, S. W. Horgan, and M. Orchin, *J. Org. Chem.*, 1981, 46, 1987.
8. Both isomers are interconvertible under the conditions of irradiation for 1. Cf, P. C. M. van Noort and H. Cerfontain, *J. Chem. Soc., Perkin Trans. II*, 1978, 757.
9. The one with a larger coupling constant was usually found to correspond to the *endo*-isomer.
10. H. E. Zimmerman, R. J. Boettcher, and W. Braig, *J. Am. Chem. Soc.*, 1973, 95, 2155.
11. D. DeKeukeleire, E. C. Sanford, and G. S. Hammond, *J. Am. Chem. Soc.*, 1973, 95, 7904.
12. S. S. Hixon and J. C. Tausta, *Tetrahedron Lett.*, 1974, 2007.
13. H. E. Zimmerman and P. S. Mariano, *J. Am. Chem. Soc.*, 1969, 91, 1718.

Received, 9th July, 1990