



REDUCTIVE PHOTOCYCLIZATION OF α -METHYLTHIO- AND
 α -ARYLTHIOENAMIDES

Takeaki Naito,* Hiromi Tanada, Toshiko Kiguchi, and
Ichiya Ninomiya
Kobe Women's College of Pharmacy, Motoyamakita, Higashinada,
Kobe 658, Japan


Abstract--Reductive photocyclization of α -methylthio- and
 α -arylthioenamides (**2a-c**) gave six-membered lactams (**3a-d**)
and five-membered lactams (**4a,b**).

Enamide photocyclization¹ has been established as one of the most useful cyclization reactions for constructing six-membered lactams from the readily available enamides. Particularly, reductive photocyclization² of enamide in the presence of a hydride reagent has provided a promising and potentially versatile approach to the formation of six-membered lactams possessing a wide variety of functionality. As an extension of our research on reductive photocyclization of enamide, we have investigated the photochemical reaction of the α -methylthio- and α -arylthioenamides (**2a-c**) in the presence of sodium borohydride and found that these enamides undergo reductive photocyclization to give not only six-membered lactams but also five-membered lactams. Phenylthio- and 2-naphthylthioimidates (**1b,c**) were readily prepared from the known³ methylthioimide (**1a**) by treatment with either thiophenol or 2-naphthalenethiol. Acylation of three thioimidates (**1a-c**) with methacryloyl chloride gave three types of enamides (**2a-c**) in 75-85% yield. Irradiation of the methylthioenamide (**2a**) with a high pressure mercury lamp through a Pyrex filter in the presence of sodium borohydride in acetonitrile-methanol (9:1) gave many products which were carefully separated by column chromatography to give two cyclized products (**3a**)⁴ and (**3b**)⁵ in 48 and 8% yields respectively.

Both lactams (3a,b) are the expected six-membered lactams and their stereochemistries were deduced from their spectral data.^{4,5} Reductive photocyclization of the phenylthioenamide (2b) afforded three products (3c),⁶ (4a),⁷ and (5)⁸ in 6, 20, and 9% yields respectively. The product (3c) is an expected six-membered lactam while the product (4a) was found to be a five-membered lactam based on the analysis of their spectral data.⁷ Analysis of the spectral data⁸ of the product (5) established its structure as a tricyclic product which would be formed by the double cyclization (Domino cyclization) of the α -phenylthioenamide (2b) via an intermediary phenyl vinyl sulfide (A)⁹ which would be formed in situ by the cyclization of the enamide (2b). Very interestingly, reductive photocyclization of the (2-naphthylthio)enamide (2c) proceeded smoothly to give two lactams (3d)¹⁰ and (4b)¹¹ in 7 and 46% yields respectively, both of which have no naphthylthio group. Desulfurization reaction of the lactams (3a-c) and (4a) by treatment with tributyltin hydride in the presence of 2,2'-azobisisobutyronitrile gave the known lactams, 1,3-dimethyl-2-piperidinone¹² and 1,3,3-trimethyl-2-pyrrolidinone¹³ respectively. Since all these lactams (3-5) were not formed under non-photochemical condition (with exclusion of light) or in the absence of sodium borohydride, the above finding provided a new photochemical cyclization of enamide which affords not only six-membered lactam but also five-membered lactam depending upon the substituent at the α -position of the enamide. The newly found reductive photocyclization of the (2-naphthylthio)enamide giving the pyrrolidinone derivatives has been successfully applied to a facile synthesis of 2-azaspiro-[4.4]nonane derivative which is a skeleton existing in polyzonimine.^{14,15} Reductive photocyclization of the enamide (6), prepared from the imidate (1c) and cyclopentene-1-carbonyl chloride, gave the spiro compound (7)¹⁶ and the six-membered lactam (8)¹⁷ in 21 and 13% yields respectively. This new aspect of reductive photocyclization of the α -arylthioenamide would provide a potential method for the construction of the five-membered lactams. Investigation for the reaction mechanism and further application have now been extensively continued.



Entry	R ¹	R ²	R ¹	R ²	R ¹	R ²	R ¹
	2		3		4		5
			(%)		(%)		(%)
1	2a	Me Me	3a	Me —SMe (48)			
			3b	Me ...SMe (8)			
2	2b	Me Ph	3c	Me SPh (6)	4a	Me SPh (20)	5 Me (9)
3	2c	Bn 2-Naph	3d	Bn H (7)	4b	Bn H (46)	



REFERENCES AND NOTES

- I. Ninomiya and T. Naito, "The Alkaloids," ed. by A. Brossi, Academic Press, New York, 1983, Vol. XXII, pp. 189-279.
- T. Naito, Y. Tada, Y. Nishiguchi, and I. Ninomiya, *J. Chem. Soc., Perkin Trans. 1*, **1985**, 487 and cited references.
- B. Zeeh and H. Kiefer, *Liebigs Ann. Chem.*, **1975**, 1984.
- 3a**: colorless oil; ν 1630 cm⁻¹; m/z 173 (M⁺); δ (500 MHz) 4.38 (ddd, J=4, 3, 1 Hz, 6-Heq), 2.37 (ddq, J=11, 8, 7 Hz, 3-Hax), 1.96 (tdd, J=13, 11, 4 Hz, 4-Hax), 1.24 (d, J=7 Hz, 3-Me).
- 3b**: colorless oil; ν 1632 cm⁻¹; m/z 173 (M⁺); δ (500 MHz) 4.38 (t, J=5.5 Hz,

- 6-Heq), 2.45 (br sextet, $J=7$ Hz, 3-Heq), 2.17 (dddd, $J=14, 9, 7, 3$ Hz, 4-Hax), 1.24 (d, $J=7$ Hz, 3-Me).
6. **3c**:pale yellow oil; ν 1632 cm^{-1} ; m/z 235 (M^+); δ (200 MHz) 4.75 (t-like, $J=3.5$ Hz, 6-Heq), 2.37 (ddq, $J=10, 8, 7$ Hz, 3-Hax), 1.22 (d, $J=7$ Hz, 3-Me).
7. **4a**:pale yellow oil; ν 1678 cm^{-1} ; m/z 235 (M^+); δ (200 MHz) 4.84 (dd, $J=8, 5$ Hz, 5-H), 2.37 (dd, $J=14, 8$ Hz, 4-H), 2.02 (dd, $J=14, 5$ Hz, 4-H), 1.14, 0.98 (each s, 3-Me $\times 2$).
8. **5**:pale yellow oil; ν 1702 cm^{-1} ; m/z 233 (M^+); δ (200 MHz) 4.87 (d, $J=12$ Hz, 8a-H), 3.38 (br d, $J=12$ Hz, 3a-H), 2.98 (s, NMe), 1.52, 1.18 (each s, 3-Me $\times 2$). The nuclear Overhauser effect spectroscopy (NOESY) spectrum of **5** showed cross peaks due to the NOE between 3 α -methyl and 3a-H and 3 β -methyl and 8a-H, indicating the stereochemistry as trans.
9. A. G. Schultz, W. Y. Fu, R. D. Lucci, B. G. Kurr, K. M. Lo, and M. Boxer, J. Am. Chem. Soc., 1978, **100**, 2140.
10. M. Mori, Y. Washioka, T. Urayama, K. Yoshiura, K. Chiba, and Y. Ban, J. Org. Chem., 1983, **48**, 4058. **3d**:colorless oil; ν 1622 cm^{-1} ; m/z 203 (M^+); δ (200 MHz) 2.52 (br sextet, $J=7$ Hz, 3-H), 1.33 (d, $J=7$ Hz, 3-Me).
11. **4b**:colorless oil; ν 1674 cm^{-1} ; m/z 203 (M^+); δ (200 MHz) 3.14 (t, $J=7$ Hz, 5-H₂), 1.84 (t, $J=7$ Hz, 4-H₂), 1.18 (s, Me $\times 2$).
12. P. Deslongchamps, U. O. Cheriyan, and D. R. Patterson, Can. J. Chem., 1975, **53**, 1682.
13. P. Hullot, Th. Cuvigny, M. Larcheveque, and H. Normant, Can. J. Chem., 1976, **54**, 1098.
14. J. Meinwald, J. Smolanoff, A. T. McPhail, R. W. Miller, T. Eisner, and K. Hicks, Tetrahedron Lett., 1975, 2367 and cited references.
15. T. Sugahara, Y. Komatsu, and S. Takano, J. Chem. Soc., Chem. Commun., 1984, 214.
16. **7**:pale yellow oil; ν 1670 cm^{-1} ; m/z 229 (M^+); δ (200 MHz) 4.50 (s, NCH_2Ph), 3.17 (t, $J=7$ Hz, 3-H₂), 1.89 (t, $J=7$ Hz, 4-H₂).
17. **8**:pale yellow oil; ν 1620 cm^{-1} ; m/z 229 (M^+); δ (200 MHz) 4.74, 4.54 (ABq, $J=14$ Hz, NCH_2Ph), 3.20 (t-like, $J=7$ Hz, 3-H₂), 2.80 (q, $J=8$ Hz, 7a-H).

Received, 20th May, 1991