

CONCISE AND VERSATILE SYNTHESSES OF
N-ARYLALKYLPYPERIDINES AS POTENTIAL INTERMEDIATES FOR
 4-ANILIDOPYPERIDINE ANALGESICS


Young-Ger Suh,* Dong-Yun Shin, Kyung-Ho Cho, and Jae-Sang Ryu

College of Pharmacy, Seoul National University, San 56-1, Shinrim-Dong,
 Kwanak-Gu, Seoul 151-742, Korea

Abstract - *N*-Arylalkylpiperidones and *N*-arylalkylspiroepoxypiperidine as potential intermediates for 4-anilidopiperidine analgesics and their structural analogues have been efficiently synthesized from the simple arylalkylamines by two and three step sequences respectively.


4-Anilidopiperidine represents a class of potent analgesics.¹ Particularly fentanyl² and sufentanil³ which are typical structures of this series, have been the recent subjects due to their potent analgesic properties. Consequently, some extensive synthetic studies focused on fentanyl, sufentanil and their analogues have been carried out although few synthetic studies on sufentanil have been reported.⁴⁻⁶ We have recently been searching for a synthetic method for easy access to the potential intermediates of widespread utility in syntheses of diverse anilidopiperidine analgesics. Herein we report our recent studies on concise and efficient synthetic method for potential *N*-arylalkylpiperidine intermediates.

Scheme 1



Our synthetic approach shown in Scheme 1 involves a facile construction of the initial *N*-arylalkylpiperidone intermediate (**3**) by the expedient two step sequence of aminomethano desilylation-cyclization⁷ followed by oxidation. The *N*-arylalkylpiperidone (**3**) has been also directly converted to the spiroepoxypiperidine (**1**) as a second potential intermediate which reacted with aniline to afford 4-anilinopiperidine or 4-anilinomethylpiperidine as its regioisomer. The arylalkylpiperidones (**3a**)⁴ and (**3b**) were efficiently prepared by subjection of phenylethylamine (**4a**) or thiophenylethylamine(**4b**)⁸ to Grieco's conditions⁷ followed by Swern oxidation⁹ of the resulting hydroxypiperidine (**5**) as shown in Scheme 2. It is noteworthy that only Swern oxidation was effective for oxidation of hydroxypiperidine (**5**). The known intermediate (**3a**) was straightforwardly transformed into fentanyl in 58% overall yield by analogy of the reported procedure.⁶


Scheme 2



i) allylsilane, 37% HCHO (2.5 eq), H₂O, 58 °C, 63% for **5a**, 45% for **5b** ii) (COCl)₂, DMSO, Et₃N, CH₂Cl₂, -78 °C, 84% for **3a**, 83% for **3b**

The one step conversion of thiophenylethylpiperidone (**3b**) to the spiroepoxide (**6**) as a second potential intermediate was achieved by dimethylsulfonium ylide¹⁰ treatment as shown in Scheme 3. The synthetic utility of spiroepoxide (**6**) was demonstrated by transformation to anilinopiperidines (**7**) and anilinomethylpiperidine (**8**). The variety of reaction conditions for regioselective ring opening of the epoxide (**6**) at more substituted carbon by aniline¹¹ were examined and triethyloxonium tetrafluoroborate (Et₃O⁺BF₄⁻) as Lewis acid in methylene chloride below -78 °C turned out to be the best choice for the highest regioselection (1.8 : 1) in favor of (**7**). Use of other Lewis acids afforded the regioisomer (**8**) as a predominant product or only byproducts by retro-Mannich type reaction. Although the regioselectivity and yields are not satisfactory yet, the direct introduction of aniline nucleophile to spiroepoxide at more substituted carbon enables two step conversion of the arylalkylpiperidone (**3b**) to the highly advanced sufentanyl intermediate (**8**) which has been prepared from *N*-benzylpiperidone by seven steps.⁴ The complete regioselection in epoxide ring opening at less substituted carbon could be also achieved by reaction of epoxide (**6**) with aniline in the absence of Lewis acid. The anilinomethylpiperidine (**8**) was further transformed into the sufentanil analogue (**2**)¹³ which is presently under biological evaluation.

Scheme 3



i) $\text{Me}_3\text{S}^+\text{I}^-$, NaCH_2SOMe , THF, 78% ii) $\text{Et}_3\text{O}^+\text{BF}_4^-$, PhNH_2 , -78°C , CH_2Cl_2 , 23% based on 60% conversion iii) PhNH_2 , 35°C , 82% after recycle iv) $(\text{EtCO})_2\text{O}$, 110°C then NaOMe , MeOH , 60%

In conclusion, arylalkylpiperidone (3) and spiroepoxypiperidine (1) have been prepared from arylalkylamine by two and three steps respectively. One step conversion of piperidone (3b) to the potential spiroepoxypiperidine intermediate (6) followed by direct regioselective epoxide opening with aniline envisions easy access to sufentanyl as well as variety of its structural analogues.

ACKNOWLEDGEMENT

We are grateful to the Research Center for New Drug Development at Seoul National University and TAI YANG Chemical Co. for support of this research.

REFERENCES

1. A. Korolkovas, 'Essentials of Medicinal Chemistry,' Wiley Inc., New York, 1988, pp. 236-257; P. A. J. Janssen and C. A. M. Van der Eycken, 'Drugs Affecting the Central Nervous System,' ed. by A. Burger, Marcel Dekker, New York, 1968, pp. 51-54.
2. P. A. J. Janssen, *Br. J. Anaesth.*, 1962, **34**, 260; J. F. Gardocki, J. Yelnosky, W. F. Kuehn, and M. J. C. Gunster, *Toxicol. Appl. Pharmacol.*, 1964, **6**, 593.
3. P. G. H. Van Daele, M. F. L. De Bruyn, J. M. Boey, S. Sanczuk, J. T. M. Agten, and P. A. J. Janssen, *Arzeim.-Forsch.*, 1976, **26**, 1521.; W. F. M. Van Bever, C. J. E. Niemegeers, K. H. L. Schellekens, and P. A. J. Janssen, *Arzeim.-Forsch.*, 1976, **26**, 1548.
4. D. Landicer and L. A. Mitscher, 'The Organic Chemistry of Drug Synthesis,' John Wiley & Sons, New York, 1977, Vol. 1, pp. 286-311 and 1984, Vol. 3, pp. 116-121.
5. R. F. Borne, E. K. Fifer, and I. W. Waters, *J. Med. Chem.*, 1984, **27**, 1271; B. E. Maryanoff, D. F. McComsey, R. J. Taylor, Jr., and J. F. Gardocki, *J. Med. Chem.*, 1981, **24**, 79.

6. W. F. M. Van Bever, C. J. E. Niemegeers, and P. A. J. Janssen, *J. Med. Chem.*, 1974, **17**, 1047.
7. P. A. Grieco and W. F. Fobare, *Tetrahedron Lett.*, 1986, **27**, 5067; S. D. Larsen, P. A. Grieco, and W. F. Fobare, *J. Am. Chem. Soc.*, 1986, **108**, 3512.
8. Thiophenylethylamine was conveniently prepared from the commercially available thiophene-acetonitrile by LAH reduction.
9. W. R. Roush, J. A. Straub, and M. S. VanNieuwenhze, *J. Org. Chem.*, 1991, **56**, 1636.
10. E. Borredon, M. Delmas, and A. Gaset, *Tetrahedron Lett.*, 1987, **28**, 1877 and references cited therein.
11. Regioselective ring opening of unsymmetrical spiroepoxide by aniline has not been reported yet. For the related references, see; M. Onaka, K. Sugita, and Y. Izumi, *Chem. Lett.*, 1986, 1327; H. B. Meryala and B. Frei, *Helv. Chim. Acta*, 1986, **69**, 415.
12. The low yields for (7) is mainly attributed to decomposition of the substrate as the reaction proceeds. Currently, intensive studies for yield optimization are in progress.
13. Spectral data for the sufentanyl analogue (2): IR 3368, 1637 cm^{-1} ; $^1\text{H-NMR}$ (400 MHz) 7.35 (dd, $J=7.3, 7.3$ Hz, 2H), 7.26 (dd, $J=7.3, 7.3$ Hz, 1H), 7.13 (d, $J=7.3$ Hz, 2H), 7.03 (dd, $J=4.9, 1.0$ Hz, 1H), 6.83 (dd, $J=4.9, 3.7$ Hz, 1H), 6.73 (d, $J=3.7$ Hz, 1H), 4.89 (br s, 1H), 3.72 (s, 2H), 2.93 (t, $J=7.8$ Hz, 2H), 2.59 (br t, $J=7.8$ Hz, 4H), 2.42 (t, $J=10.7$ Hz, 2H), 2.05 (q, $J=7.3$ Hz, 2H), 1.63 (d, $J=10.7$ Hz, 2H), 1.46 (t, $J=10.7$ Hz, 2H), 0.98 (t, $J=7.3$ Hz, 3H); MS (m/z) 373 ($\text{M}^+\text{+H}$), 355, 275; HRMS Found, 373.5415 Calcd for $\text{C}_{21}\text{H}_{29}\text{N}_2\text{O}_2\text{S}(\text{MH})$, 373.5321.

Received, 7th November, 1997