

***ENT-KAURANE DITERPENOIDS FROM ISODON RUBESCENS VAR. LUSHIENSIS***


**Quanbin Han, Shenghong Li, Liyan Peng, and Handong Sun\***

State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, Yunnan, P. R. China; hdsun@mail.kib.ac.cn


**Abstract** – Three new 6,7-seco-6,19:6,20-diepoxy-1 $\alpha$ ,7-olide-*ent*-kaurane diterpenoids, ludongnins C-E (**1-3**), along with six known analogs, macrocalyxoformins A (**4**) and B (**5**), sculponeatins A (**6**) and B (**7**), and ludongnins A (**8**) and B (**9**), were isolated from *Isodon rubescens* var. *lushiensis*. The structures of the new compounds were determined on the basis of the spectroscopic evidences. Compound (**5**) exhibited significant inhibitory activities against the Bcap37 and K562 cell lines with IC<sub>50</sub> = 0.82 and 0.62  $\mu$ g/mL, respectively.

**INTRODUCTION**

The plants of the genus *Isodon* (Labiatae) are well known as a rich source of *ent*-kaurane diterpenoids.<sup>1</sup> About twenty *ent*-kaurane diterpenoids have been reported from *Isodon rubescens*,<sup>2</sup> which was the first studied species of this genus plants in China. Among them, oridonin and ponidicin which belong to



	<b>R<sub>1</sub></b>	<b>R<sub>2</sub></b>	<b>R<sub>3</sub></b>
<b>1</b>	H	= O	$\alpha$ -CH <sub>3</sub>
<b>2</b>	H	= O	$\beta$ -CH <sub>3</sub>
<b>3</b>	$\beta$ -OH	= O	$\alpha$ -CH <sub>3</sub>
<b>4</b>	H	$\alpha$ -OH	= CH <sub>2</sub>
<b>5</b>	H	= O	= CH <sub>2</sub>
<b>6</b>	$\beta$ -OH	= O	= CH <sub>2</sub>
<b>7</b>	$\beta$ -OH	$\alpha$ -OH	= CH <sub>2</sub>



7,20-epoxy type *ent*-kauranoids were regarded as the main diterpene constituents of this plant.<sup>3</sup> However, there were only two 6,7-seco-7,20-olide-*ent*-kaurane diterpenoids (ludongnins A and B) reported from *Isodon rubescens* var. *lushiensis* Gao et Li.<sup>4</sup> In our continuing search for biological active principles of *Isodon rubescens* and its varieties, we have reinvestigated on *Isodon rubescens* var. *lushiensis* collected in Lushi Prefecture, Henan Province of China. As a result, three new 6,7-seco-6,19:6,20-diepoxy-1 $\alpha$ ,7-olide-*ent*-kauranoids named ludongnins C-E (**1-3**) together with six known diterpenoids identified as macrocalyxofornins A and B (**4** and **5**),<sup>5</sup> sculponeatins A and B (**6** and **7**)<sup>6</sup>, and ludongnins A and B (**8** and **9**) were isolated. In this paper, we wish to present the structural elucidation of these new compounds and the biological testing results of compounds (**5**) and (**7**) against the Bcap37 and K562 tumor cell lines.

## RESULTS AND DISCUSSION

Compound (**1**), obtained as colorless quadrate lumpish crystals with  $[\alpha]_D^{20} -110.0^\circ$  ( $c$  0.82, MeOH), was determined to possess the molecular formula  $C_{20}H_{26}O_5$  by the HREIMS (found 346.1762, calcd 346.1780). The  $^{13}C$  and DEPT NMR spectra of **1** showed 20 carbon signals which were indicated to be composed of one carbonyl carbon, one lactonic carbonyl carbon, one acetal carbon, three quaternary carbons, five methines including one oxygenated, five methylenes besides two oxygenated ones, and two methyls, which suggested **1** as an *ent*-kauranoid, combined with the consideration of the structures of diterpenoids previously isolated from this plant.<sup>4</sup> Furthermore, the basic skeleton of 6,7-seco-6,19:6,20-diepoxy-1 $\alpha$ ,7-olide-*ent*-kaurane was deduced on the basis of the presence of the characteristic

Table 1:  $^{13}C$  NMR Spectral Data ( $C_5D_5N$ ) of **1-4**.

C	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
1	76.5 (d)	76.3 (d)	79.1 (d)	76.3 (d)
2	23.0 (t)	23.0 (t)	23.4 (t)	23.2 (t)
3	29.1 (t)	28.9 (t)	29.5 (t)	29.1 (t)
4	41.4 (s)	41.3 (s)	41.7 (s)	41.3 (s)
5	53.0 (d)	53.1 (d)	54.2 (d)	53.3 (d)
6	111.3 (d)	111.3 (d)	111.8 (d)	111.4 (d)
7	171.9 (s)	171.7 (s)	172.3 (s)	171.2 (s)
8	56.6 (s)	56.7 (s)	56.6 (s)	56.5 (s)
9	43.7 (d)	42.9 (d)	47.0 (d)	42.9 (d)
10	50.6 (s)	50.9 (s)	50.4 (s)	50.9 (s)
11	19.2 (t)	19.6 (t)	64.9 (d)	19.3 (t)
12	19.2 (t)	29.3 (t)	30.7 (t)	29.6 (t)
13	32.7 (d)	35.2 (d)	32.9 (d)	35.0 (d)
14	33.9 (t)	32.1 (t)	35.6 (t)	32.5 (t)
15	215.0 (s)	214.9 (s)	215.9 (s)	200.3 (s)
16	49.4 (d)	51.0 (d)	49.8 (d)	151.0 (t)
17	10.3 (q)	15.9 (q)	11.0 (q)	118.0 (t)
18	30.3 (q)	30.3 (q)	30.7 (q)	30.3 (q)
19	77.0 (t)	77.0 (t)	77.2 (t)	77.0 (t)
20	73.1 (t)	72.9 (t)	73.1 (t)	72.8 (t)

signals of a  $\delta$ -lactone group ( $\delta_C$  171.9, C-7;  $\delta_C$  76.5, C-1), two oxymethylene carbons ( $\delta_C$  77.0, C-19;  $\delta_C$  73.1, C-20), and one acetal group ( $\delta_C$  111.3, C-6). Comparison of the  $^{13}C$  NMR spectrum (Table 1) of **1** with that of macrocalyxofornin B (**5**), a known 6,7-seco-6,19:6,20-diepoxy-1,7-olide-*ent*-kauranoid isolated from the same plant this time, revealed that the olefinic signals [ $\delta_C$  151.0 (s, C-16) and 118.2 (t, C-17)] in **5** were replaced by a methine signal ( $\delta_C$  49.4, C-16) and a methyl signal ( $\delta_C$  10.3, C-17) in **1**,

which was approved by the  $^1\text{H}$ - $^1\text{H}$  COSY correlations of a methine proton ( $\delta_{\text{H}}$  2.43, H-16) with a methyl group ( $\delta_{\text{H}}$  0.99, H<sub>3</sub>-17) and a methine proton ( $\delta_{\text{H}}$  2.41, H-13), and by the HMBC correlations (Figure 1) of this methyl group with a ketonic carbon ( $\delta_{\text{C}}$  215.0, C-15) and a methine carbon ( $\delta_{\text{C}}$  32.7, C-13). The  $\alpha$ -orientation of 16-methyl group was determined by the significant upfield shift of C-12 ( $\Delta\delta$  10.4, from  $\delta_{\text{C}}$  29.6 in **5** to  $\delta_{\text{C}}$  19.2 in **1**) caused by a steric compression effect between Me-16 $\alpha$  and H-12 $\alpha$ , which was confirmed by the ROESY correlation (Figure 1) between H<sub>3</sub>-17 and H-12 $\alpha$ . Therefore, compound (**1**) was elucidated as 16(*R*<sup>\*</sup>)-6,7-seco-6,19;6,20-diepoxy-1,7-olide-*ent*-kaur-15-one, named ludongnin C.




Figure 1 Key HMBC (from H to C) and ROESY Correlations of **1**

Compound (**2**) exhibited the same HREIMS experimental result (found 346.1779, calcd 346.1780 for C<sub>20</sub>H<sub>26</sub>O<sub>5</sub>) as **1**. A general analysis of all spectra obtained revealed that the structure of **2** was quite similar to that of **1** except for two obvious downfield shifts of C-12 and C-17 (Table 1), suggesting **2** to be the 16(*S*<sup>\*</sup>)-epimer of **1** without the above-mentioned steric compression effect. This deduction was proved by the correlations between H<sub>3</sub>-17 with H-13 $\beta$  and H-14 $\alpha$  in the ROESY spectrum of **2**. Then, compound (**2**) was determined as 16(*S*<sup>\*</sup>)-6,7-seco-6,19;6,20-diepoxy-1,7-olide-*ent*-kaur-15-one, named ludongnin D.

Compound (**3**) showed the molecular ion peak at  $m/z$  362 in the EIMS spectrum with the molecular formula C<sub>20</sub>H<sub>26</sub>O<sub>6</sub> determined by the HREIMS (found 362.1747, calcd 362.1729). Comparison of the  $^{13}\text{C}$  NMR data of compounds (**1**) and (**3**) suggested that compound (**3**) was consistent with **1** except for a more hydroxyl group at C-11 of **3** which was approved by the HMBC correlations arising from H-11 ( $\delta_{\text{H}}$  4.43) with C-10 ( $\delta_{\text{C}}$  50.4) and C-13 ( $\delta_{\text{C}}$  32.9). The  $\beta$ -orientation of OH-11 was established by the small coupling constant ( $J = 4.0$  Hz) between H-11 $\alpha$  with H-9 $\alpha$  ( $\delta_{\text{H}}$  2.01) and the ROESY correlation between H-11 $\alpha$  with H-5 $\beta$  ( $\delta_{\text{H}}$  2.96), which was confirmed by the downfield shifts of H-1 $\beta$  ( $\delta_{\text{H}}$  5.51) and H-14 $\beta$  ( $\delta_{\text{H}}$  3.75),<sup>6</sup> and by the upfield shift of C-11 ( $\delta_{\text{C}}$  64.9) caused by the steric compress effect between H-11 $\alpha$

with H-5 $\beta$ . Hence, compound (3) was established as 16(*R*\*)-6,7-seco-11 $\beta$ -hydroxy-6,19:6,20-diepoxy-1 $\alpha$ ,7-olide-*ent*-kaur-15-one, named ludongnin E.

Six known diterpenoids isolated from *I. rubescens* var. *lushiensis* were identified as macrocalyxoformins A and B (4 and 5),<sup>5</sup> sculponeatins A and B (6 and 7),<sup>6</sup> and ludongnins A and B (8 and 9),<sup>4</sup> respectively, by comparing their spectral data with those reported in the literatures. The <sup>13</sup>C NMR spectral data of macrocalyxoformins B (5) were reported for the first time. And the chemical shifts of C-19 and C-20 in compounds (6) and (7) were revised to exchange according to the HMBC correlations between H-19 with C-18, and H-20 with C-1.

Compounds (5) and (7) were tested for their inhibitory activities against the Bcap37 and K562 cell lines using a previously described method with cis-platinum as a positive reference substance.<sup>7</sup> Compound (5) showed significant inhibitory activities against the Bcap37 and K562 cell lines with IC<sub>50</sub> = 0.82 and 0.62  $\mu$ g/mL (cis-platinum IC<sub>50</sub> = 1.54 and 3.84  $\mu$ g/mL), respectively, while compound (7) was inactive against either Bcap37 or K562 cell lines, which confirmed that the cyclopentanone conjugated with an exo-methylene group was the active center of the inhibitory effect.<sup>8</sup>

## EXPERIMENTAL

**General:** Melting points were measured on an XRC-1 micro melting point apparatus and uncorrected. Optical rotations were taken on a JASCO DIP-370 digital polarimeter. IR spectra were determined on Bio-Rad FtS-135 spectrophotometer with KBr pellets. UV spectra were obtained on a UV 210A spectrophotometer. MS were recorded on a VG Auto Spec-3000 spectrometer. 1D- and 2D- NMR spectra were run on a Bruker AM-400 and a DRX-50 instrument with TMS as internal standard, respectively.

**Plant material:** The leaves of *I. rubescens* var. *lushiensis* were collected in Lushi prefecture of Henan Province in August 2000, and air-dried. The identity of the plant material was verified by Prof. Zhong-Wen Lin, and a voucher specimen (KIB-2000-10 Lin) is deposited in the Herbarium of the Department of Taxonomy, Kunming Institute of Botany, Chinese Academy of Science.

**Extraction and Isolation:** The dried and powdered leaves (7.6 kg) were extracted with 70% Me<sub>2</sub>CO (3 $\times$ 10 L) at rt for 72 h and filtered. And then, the filtrate was concentrated and extracted with petroleum ether and EtOAc in turn. The EtOAc extract (350 g) was applied to column chromatography over Silica gel (100-200 mesh, 3.0 kg) column eluting with a gradient system of CHCl<sub>3</sub>-Me<sub>2</sub>CO (1:0-1:1). The CHCl<sub>3</sub>-Me<sub>2</sub>CO (8:2) fraction (40 g) was further chromatographed

repeatedly over Silica gel (cyclohexane: EtOAc = 4:1) to afford **4** (1.0 g), **5** (300 mg), **6** (150 mg), and **7** (120 mg). The CHCl<sub>3</sub>-Me<sub>2</sub>CO (7:3) fraction (23 g) was chromatographed on Silica gel (CH<sub>2</sub>Cl<sub>2</sub>: 2-propanol, from 150:1 to 25:1), then fined with preparative TLC (developed with CH<sub>2</sub>Cl<sub>2</sub>: 2-propanol = 100:1 for four times) and recrystallization to yield compounds (**1**) (6.0 mg), (**2**) (2.0 mg), and (**3**) (20.0 mg).

**Ludongnin C (1):** colorless quadrate lumpish crystals; mp > 250°C; [ $\alpha$ ]<sub>D</sub><sup>20</sup> -110.0° (*c* 0.82, MeOH); UV (MeOH): end absorption; IR (KBr)  $\nu_{\max}$ : 3436, 2947, 2875, 1760 (C=O), 1720 (C=O), 1392, 1268, 1195, 1113, 1037 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, C<sub>5</sub>D<sub>5</sub>N): 6.03 (1H, d, *J* = 5.2 Hz, H-6 $\beta$ ), 4.64 (1H, dd, *J* = 5.3, 11.6 Hz, H-1 $\beta$ ), 4.23 and 4.06 (each 1H, d, *J* = 9.0 Hz, H<sub>2</sub>-20), 3.96 and 3.42 (each 1H, d, *J* = 8.7 Hz, H<sub>2</sub>-19), 2.59 (1H, d, *J* = 12.0 Hz, H-14 $\beta$ ), 2.43 (1H, overlapped, H-16 $\beta$ ), 2.41 (1H, overlapped, H-13 $\beta$ ), 2.37 (1H, d, *J* = 5.2 Hz, H-5 $\beta$ ), 2.10 (1H, overlapped, H-12 $\beta$ ), 2.07 (1H, overlapped, H-9 $\alpha$ ), 1.86 (1H, overlapped, H-14 $\alpha$ ), 1.86-1.78 (2H, overlapped, H<sub>2</sub>-2), 1.72 (1H, m, H-12 $\alpha$ ), 1.70-1.34 (2H, overlapped, H<sub>2</sub>-11), 1.62 and 1.40 (each 1H, overlapped, H<sub>2</sub>-3), 1.04 (3H, s, Me-18), 0.99 (3H, d, *J* = 6.9 Hz, Me-17); <sup>13</sup>C NMR (C<sub>5</sub>D<sub>5</sub>N) see Table 1; EIMS *m/z*: 346 [M]<sup>+</sup> (32), 328 (36), 318 (40), 300 (20), 289 (44), 272 (18), 258 (32), 165 (77), 136 (42), 121 (39), 107 (60), 91 (65), 55 (100); HREIMS *m/z*: 346.1762 (calcd for C<sub>20</sub>H<sub>26</sub>O<sub>5</sub>, 346.1780).

**Ludongnin D (2):** white amorphous powder; [ $\alpha$ ]<sub>D</sub><sup>20</sup> -60.0° (*c* 0.30, MeOH); UV (MeOH): end absorption; <sup>1</sup>H NMR (500 MHz, C<sub>5</sub>D<sub>5</sub>N): 6.02 (1H, d, *J* = 5.0 Hz, H-6 $\beta$ ), 4.62 (1H, dd, *J* = 5.3, 11.4 Hz, H-1 $\beta$ ), 4.21 and 4.04 (each 1H, d, *J* = 9.0 Hz, H<sub>2</sub>-20), 3.96 and 3.42 (each 1H, d, *J* = 8.7 Hz, H<sub>2</sub>-19), 2.52 (1H, d, *J* = 11.8 Hz, H-14 $\beta$ ), 2.50 (1H, overlapped, H-16 $\alpha$ ), 2.16 (1H, overlapped, H-13 $\beta$ ), 2.39 (1H, d, *J* = 5.0 Hz, H-5 $\beta$ ), 2.10 (1H, overlapped, H-12 $\alpha$ ), 2.00 (1H, overlapped, H-12 $\beta$ ), 2.11 (1H, overlapped, H-9 $\alpha$ ), 1.82 (1H, overlapped, H-14 $\alpha$ ), 1.88-1.77 (2H, overlapped, H<sub>2</sub>-2), 1.68 and 1.32 (each 1H, overlapped, H<sub>2</sub>-11), 1.60 and 1.39 (each 1H, overlapped, H<sub>2</sub>-3), 1.04 (3H, s, Me-18), 1.07 (3H, d, *J* = 6.9 Hz, Me-17); <sup>13</sup>C NMR (C<sub>5</sub>D<sub>5</sub>N), see Table 1; EIMS *m/z*: 346 [M]<sup>+</sup> (32), 328 (60), 318 (90), 300 (24), 289 (70), 272 (35), 259 (60), 165 (100), 107 (70), 91 (75), 55 (100); HREIMS *m/z*: 346.1779 (calcd for C<sub>20</sub>H<sub>26</sub>O<sub>5</sub>, 346.1780).

**Ludongnin E (3):** white amorphous powder; [ $\alpha$ ]<sub>D</sub><sup>20</sup> -122.5° (*c* 0.20, acetone); UV (MeOH): end absorption; <sup>1</sup>H NMR (500 MHz, C<sub>5</sub>D<sub>5</sub>N): 6.11 (1H, d, *J* = 5.0 Hz, H-6 $\beta$ ), 5.51 (1H, dd, *J* = 5.5, 12.0 Hz, H-1 $\beta$ ), 4.43 (1H, t, *J* = 4.0 Hz, H-11 $\alpha$ ), 4.26 and 4.11 (each 1H, d, *J* = 10.0 Hz, H<sub>2</sub>-20), 4.01 and 3.45 (each 1H, d, *J* = 8.5 Hz, H<sub>2</sub>-19), 3.75 (1H, d, *J* = 11.0 Hz, H-14 $\beta$ ), 2.59 (1H, m, H-13 $\beta$ ), 2.52 (1H, m, H-16 $\beta$ ), 2.96 (1H, d, *J* = 5.0 Hz, H-5 $\beta$ ), 2.09 (1H, overlapped, H-14 $\alpha$ ), 2.05 (1H, overlapped, H-12 $\beta$ ),

2.01 (1H, d,  $J = 4.0$  Hz, H-9 $\alpha$ ), 1.96-1.74 (2H, overlapped, H<sub>2</sub>-2), 1.74 (1H, dd,  $J = 4.0, 15.5$  Hz, H-12 $\alpha$ ), 1.68-1.58 (2H, overlapped, H<sub>2</sub>-3), 1.07 (3H, s, Me-18), 1.03 (3H, d,  $J = 7.0$  Hz, Me-17); <sup>13</sup>C NMR (C<sub>5</sub>D<sub>5</sub>N), see Table 1; EIMS  $m/z$ : 362 [M]<sup>+</sup> (32), 346 (36), 332 (95), 314 (100), 298 (24), 286 (48), 270 (10), 260 (22), 220 (14), 165 (26), 107 (39), 91 (56), 83 (64), 55 (62); HREIMS  $m/z$ : 362.1747 (calcd for C<sub>20</sub>H<sub>26</sub>O<sub>6</sub>, 362.1729).

## REFERENCES (AND NOTES)

1. E. Fujita and M. Node, *Prog. Chem. Org. Nat. Prod.*, 1984, **46**, 77; Y. Takeda and H. Otsuka, *Stud. Nat. Prod., Chem.*, 1995, **15**, 111; H. D. Sun, Y. L. Xu, and B. Jiang, 'Diterpenoids from Isodon Species,' Sciences Press, Beijing, 2001, 5.
2. H. D. Sun, Y. L. Xu, and B. Jiang, 'Diterpenoids from Isodon Species,' Sciences Press, Beijing, 2001, 122.
3. Henan Institute of Medical Science, Henan Medical College, Yunnan Institute of Botany, and Zhengzhou Chemicopharmaceutical Plant. *Chin. Sci. Bull.*, 1978, **23**, 53; T. M. Zhang, Z. Y. Chen, J. H. Chao, Q. Z. Zhao, H. D. Sun, and Z. W. Lin, *Chin. Sci. Bull.*, **1980**, 25, 1051; H. D. Sun, Z. W. Lin, C. Q. Qin, J. H. Cao, and Q. Z. Zhao, *Acta Bot. Yunnan.*, 1981, **3**, 95.
4. X. R. Zheng, Z. Y. Gao, H. D. Sun, and Z. W. Lin, *Acta Bot. Yunnan.*, 1984, **6**, 316; X. R. Zheng, Z. Y. Gao, J. Q. Tang, H. D. Sun, and Z. W. Lin, *Acta Bot. Yunnan.*, 1986, **8**, 161; Z. Y. Gao and Y. R. Li, *Acta Phytoaxonomica Sinica*, 1986, **24**, 15; Q. B. Han, Q. S. Zhao, S. H. Li, L. Y. Peng, and H. D. Sun, submitted to *Acta Chim. Sin.*.
5. Z. Q. Wang, X. R. Wang, and J. G. Dong, *Chinese Traditional and Herbal Drugs*, 1983, **14**, 1.
6. H. D. Sun, Z. W. Lin, Y. L. Xu, Y. Minami, T. Marunaka, T. Togo, Y. Takeda, and T. Fujita, *Heterocycles*, 1986, **24**, 1.
7. A. J. Hou, M. L. Li, B. Jiang, Z. W. Lin, S. Y. Ji, Y. P. Zhou, and H. D. Sun, *J. Nat. Prod.*, 2000, **63**, 599.
8. E. Fujita, Y. Nagao, M. Node, K. Kaneko, S. Nakazawa, and H. Kuroda, *Experientia*, 1976, **32**, 203; M. Node, M. Sai, K. Fuji, E. Fujita, S. Takeda, and M. Ozaki, *Chem. Pharm. Bull.*, 1981, **29**, 3208; M. Node, M. Sai, E. Fujita, and K. Fuji, *Heterocycles*, 1984, **22**, 1701.