


CONVENIENT SYNTHESIS OF PULCHELLALACTAM, A CD45 PROTEIN TYROSINE PHOSPHATASE INHIBITOR FROM THE MARINE FUNGUS *COROLLOSPORA PULCHELLA*, AND ITS RELATED COMPOUNDS

Jun-ichiro Bessho, Yasuhiko Shimotsu, Sachiko Mizumoto, Nobuyuki Mase, Hidemi Yoda, and Kunihiko Takabe*

Department of Molecular Science, Faculty of Engineering, Shizuoka University, 3-5-1 Johoku, Hamamatsu 432-8561, Japan. e-mail: tcktaka@ipc.shizuoka.ac.jp


Abstract – Convenient synthesis of pulchellalactam (**1a**) and its related compounds was accomplished. Total yields of (*Z*)-pulchellalactam and (*E*)-pulchellalactam were 55% for 5 steps and 43% for 8 steps from commercially available and inexpensive citraconic anhydride, respectively.

Pulchellalactam (**1a**), (5*Z*)-1,5-dihydro-4-methyl-5-(2-methylpropylidene)pyrrol-2(2*H*)-one, was isolated from the marine fungus *Corollospora pulchella* in 1998 by Alvi *et al.*¹ Its bioactivity is a dose-dependent inhibitor of CD45 protein tyrosine phosphatase with an IC₅₀ of 124 μg/mL; interestingly, this inhibition is specific for CD45. Since CD45 participates in activation of both B and T cells *via* their antigen-specific receptors, pulchellalactam (**1a**) might be expected as a drug intervention in various auto-immune and/or inflammatory diseases.^{1,2} Quite recently, the synthesis of **1a** has been reported by Li *et al.*³ The key step of the sequence involves addition and elimination of an enolic lactam with an organocuprate reagent. Their total synthesis afforded (*Z*)-**1a** in 6 steps and 32% overall yield from Boc-glycine. Independently, we have recently accomplished convenient synthesis of **1a** and its related compounds (**1b**) from commercially available and quite inexpensive citraconic anhydride (**2**), employing regioselective reduction of citraconimide (**3**) (Scheme 1).



Scheme 1. Synthetic plan of (*Z*)-**1a** from citraconic anhydride.

A conventional method for the preparation of citraconimide (**3**) is heating the anhydride (**2**) under reflux in the presence of $\text{NH}_4\text{OAc}/\text{AcOH}$,⁴ however, giving **3** in 41% yield. A higher yield was achieved by using 1,1,1,3,3,3-hexamethyldisilazane (HMDS) as ammonia equivalent in DMF at 100 °C as shown in Scheme 2. In 1999, we succeeded in synthesis of an antitumor alkaloid (jatropham) through regioselective reduction of **3**, in which NaBH_4 reduction afforded 1,5-dihydro-5-hydroxy-4-methyl-2*H*-pyrrol-2-one (**4a**), whereas $\text{NaBH}_4/\text{CeCl}_3$ or DIBAL-H reduction gave the regioisomer (**4b**) with extremely high regioselectivities.⁵ We obtained **4a** as a single regioisomer by recrystallization. Reductive deoxygenation⁶ of **4a** following Boc-protection⁷ furnished *N*-*tert*-butoxycarbonyl-1,5-dihydro-4-methyl-pyrrol-2(2*H*)-one (**5a**)⁸ in 91% yield. The pyrrol-2(2*H*)-one (**5a**) was treated with LDA at -78 °C, and the resulting anion was reacted with isobutyraldehyde. Interestingly, the product highly depended on the reaction time and temperature. Longer reaction time and higher reaction temperature gave the desirable (*Z*)-**1a** as a single geometrical isomer in a single step. On the other hand, the aldol reaction at -78 °C for 15 min gave the aldol product (**6a**) in 80% yield, which was easily transformed into (*E*)-**1a** in an *E/Z* ratio of 96/4 by elimination and deprotection.



Scheme 2. Synthesis of pulchellalactam (**1a**) and its analogue (**1b**).

Structural determinations of both isomers, (*Z*)- and (*E*)-**1a**, were performed by X-Ray crystallographic analysis as shown in Figure 1.¹⁰ We also tried to determine the structure of synthetic pulchellalactam (**1a**) by NMR spectral analyses compared with the reported value of the natural product. ¹H and ¹³C NMR chemical shifts are given in Table 1. Most of the signals were in agreement with the reported values of (*E*)-pulchellalactam; however, unfortunately signals for H-6 proton and C-6 carbon were found to differ.




Table 1. ¹H and ¹³C NMR chemical shifts (ppm).

No.	Natural ^{a)}	(<i>Z</i>)- 1a	(<i>E</i>)- 1a
H-1	9.41 (br s)	7.65 (br s)	8.10 (br s)
H-3	5.94 (br s)	5.86 (br s)	5.91 (br s)
H-6	5.40 (d, 9.87 Hz)	5.11 (d, 9.80 Hz)	5.30 (d, 10.60 Hz)
H-7	2.60-2.67 (m)	2.51-2.64 (m)	2.89-3.10 (m)
H-8,9	1.12 (d, 6.63 Hz)	1.11 (d, 6.80 Hz)	1.08 (d, 6.60 Hz)
H-10	2.11 (d, 1.16 Hz)	2.08 (s)	2.28 (s)
C-2	173.1	173.0	171.7
C-3	119.7	120.7	125.7
C-4	150.2	148.5	146.9
C-5	137.0	137.7	136.6
C-6	125.5	120.7	124.5
C-7	28.0	27.3	26.6
C-8,9	22.7	22.8	23.4
C-10	11.9	11.7	16.1

a) Revised spectrum data provided by Dr. Khisal A. Alvi.

(*Z*)-Pulchellalactam is thermodynamically quite stable at least for a half-year, while the (*E*)-isomer is relatively unstable. An *E/Z* ratio of 99/1 of (*E*)-**1a** decreased to 70/30 after 1 week, then 24/76 after 3 months, even stored in a freezer at -20 °C. It took one week or more to isolate natural pulchellalactam from the marine fungus; therefore, it seemed reasonable to assume that the stereochemistry of **1a** was *E*.

One principal advantage of our synthetic method is a convenient approach to the related compounds, 1,5-dihydro-3-methyl-5-(2-methylpropylidene)pyrrol-2(*2H*)-one (**1b**). In a similar way to the synthesis of (*Z*)-**1a**, **1b** was obtained as a mixture of geometrical isomers in an *E/Z* ratio of 54/46 in 47% overall yield.

Refluxing in benzene with iodine afforded the thermodynamically more stable geometrical isomer ((*Z*)-**1b**) in an *E/Z* ratio of 22/78. Similarly, (*E*)-**1b** was obtained in an *E/Z* ratio of 97/3 in 34% overall yield, which was relatively unstable to show an *E/Z* ratio of 82/18 after being stored at $-20\text{ }^{\circ}\text{C}$ for 1 week. In conclusion, we have shown a synthesis of (*Z*)- and (*E*)-pulchellalactams (**1a**) from commercially available citraconic anhydride. Interestingly, the product ratio in the aldol reaction highly depended on the reaction time and temperature to give the (*E*)- and (*Z*)-isomers, regioselectively.

ACKNOWLEDGEMENTS

We thank Dr. Khisal A. Alvi of AMRI Bothell Research Center for providing the NMR spectra data of pulchellalactam and for scientific discussions on the stereochemistry of pulchellalactam. This work was supported in part by a Grant-in-Aid (No.14540491) for Scientific Research from Japan Society for the Promotion of Science.

REFERENCES AND NOTES

1. K. A. Alvi, A. Casey, and B. G. Nair, *J. Antibiotics*, 1998, **51**, 515.
2. A. M. S. Mayer and V. K. B. Lehmann, *The Pharmacologist*, 2000, **42**, 62.
3. W.-R. Li, S. T. Lin, N.-M. Hsu, and M.-S. Chern, *J. Org. Chem.*, 2002, **67**, 4702.
4. P. Kuehne and M. Hesse, *Tetrahedron*, 1993, **49**, 4575; G. B. Gill, G. D. James, K. V. Oates, and G. Pattenden, *J. Chem. Soc., Perkin Trans. 1*, 1993, 2567; K. R. Shah and C. D. Blanton, Jr., *J. Org. Chem.*, 1982, **47**, 502.
5. N. Mase, T. Nishi, Y. Takamori, H. Yoda, and K. Takabe, *Tetrahedron: Asymmetry*, 1999, **10**, 4469; N. Mase, T. Nishi, M. Hiyoshi, K. Ichihara, J. Bessho, H. Yoda, and K. Takabe, *J. Chem. Soc., Perkin Trans. 1*, 2002, 707.
6. C. Pedregal, J.-A. Escribano, M. C. Carreño, and J. L. G. Ruano, *Tetrahedron Lett.*, 1994, **35**, 2053.
7. D. L. Flynn, R. E. Zelle, and P. A. Grieco, *J. Org. Chem.*, 1983, **48**, 2424.
8. In a previous synthesis of the pyrrol-2(*2H*)-one (**5a**) was obtained as a by-product in low yield, see; I. Ojima, A. Korda, and W. R. Shay, *J. Org. Chem.*, 1991, **56**, 2024; I. Ojima and A. Korda, *Tetrahedron Lett.*, 1989, **30**, 6283; K. Clauss, *Tetrahedron Lett.*, 1974, **15**, 1271.
9. The geometrical selectivity was determined by ^1H NMR spectrometry before deprotection of the Boc group due to instability of (*E*)-**1a**.
10. (*Z*)-**1a**: monoclinic ($P2_1/c$); $a=11.9896(45)\text{\AA}$, $b=6.9290(34)\text{\AA}$, $c=12.5863(32)\text{\AA}$; $\beta = 115.698(20)^{\circ}$, $Z=4$, $R1(wR2)=0.0662(0.1679)$. (*E*)-**1a**: monoclinic ($P2_1/c$); $a=9.6464(158)\text{\AA}$, $b=8.4541(63)\text{\AA}$, $c=11.7772(54)\text{\AA}$; $\beta = 107.423(67)^{\circ}$, $Z=4$, $R1(wR2)=0.0591(0.0809)$.