

HETEROCYCLES, Vol. 65, No. 3, 2005, pp. 661 - 666

Received, 21st December, 2004, Accepted, 28th January, 2005, Published online, 1st February, 2005

A NEW FLAVONE FROM *LYCOPODIUM JAPONICUM*

Jian Yan,^{1,2} Lirong Sun,^{1,2} Xianming Zhang,¹ and Minghua Qiu^{1,2} *

¹ State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of sciences, Kunming, Yunnan 650204, China


*Corresponding author: E-mail: mhchiu@public.km.yn.cn; Tel: + 86-871-5223255; Fax: +86-871-5150227

² Graduate School of the Chinese Academy of Sciences, Beijing, 100039, China

Abstract — A new flavone, lycopodone (**1**), together with other three known flavonoids tricetin (**2**), tricetin 3',4',5'-OMe (**3**), 5,7,4'-trihydroxy-3'-methoxy flavone (**4**), were isolated from the fern *Lycopodium japonicum*. The structure of **1** was established on the basis of spectral measures. Compounds (**1**) and (**2**) showed moderate antitumor activity.

INTRODUCTION

Lycopodium japonicum is widely distributed in Guangdong, Guangxi, Yunnan, Guizhou Provinces.¹ This plant is one of the most commonly encountered traditional Chinese herbal medicines for treatments of arthritic pain, quadriplegia, dysmenorrhea, and contusion.²⁻³ The chemical constituents of the genus *Lycopodium* have been investigated previously.⁴⁻⁶ But flavones of the genus were hardly reported. Numerous biological activities of flavonoids have been reported. For example, tricetin (**2**) has been evaluated with respect to antimicrobial⁸ and cancer cell line activities.⁹⁻¹⁰ In the course of our screening of biologically active constituents, four compounds (**1**), (**2**)¹⁰⁻¹¹, (**3**)¹¹, (**4**)¹² (see **Scheme 1**) were isolated from *L. japonicum*. In this paper, we would like to present the isolation and identification of the new compound, lycopodone.



Scheme 1 the structures of compounds (**1-4**)

RESULTS AND DISCUSSION

Lycopodone (**1**) possessed the molecular formula $\text{C}_{27}\text{H}_{26}\text{O}_{11}$ by HRFAB-MS (neg.) (m/z 525.0867 $[\text{M}-\text{H}]^-$, calc. for $\text{C}_{27}\text{H}_{25}\text{O}_{11}$ 525.1396), which was consistent with its NMR data. The new compound exhibited UV maximum absorptions (273 and 331 nm) and IR absorptions spectral at 3417 and 1612 cm^{-1} , which were characteristic of a flavonoid. The ^{13}C NMR (DEPT) spectra presented nine carbon signals corresponding to a $\text{C}_6\text{-C}_3$ phenylpropane unit, and the frequencies were characteristic of a flavone skeleton. The ^1H NMR spectra showed signals (H-3, 6, 8, 2' and 6') typical of a tricin-like flavone skeleton. Comparing those of **2**, the ^{13}C NMR spectra of **1** displayed an additional aromatic ring and an additional glycerol chain and a methoxy group. In the aromatic part of the ^1H NMR spectrum, the signals at δ 6.95 (1H, d, $J = 1.8$ Hz, H-2''), 6.76 (1H, d, $J = 8.8$ Hz, H-5''), and 6.70 (1H, dd, $J = 1.8, 8.8$ Hz, H-6'') defined a 1, 3, 4 tri-substituted aromatic ring. This was confirmed in the ^{13}C NMR spectrum by three methane peaks at δ_{C} 111.2 (d, C-2''), 119.4 (d, C-5''), 114.7 (d, C-6''). Due to deshielded effect, two signal peaks δ_{H} 4.36 (1H, H-7''); δ_{C} 86.3 (d, C-7'') and δ_{H} 4.80 (1H, H-8''); δ_{C} 72.2 (d, C-8'') and a methylene peaks δ_{H} 3.51 (2H, d, $J = 9.3$, H-9''); δ_{C} 60.1 (t, C-9'') were observed in the ^1H NMR and ^{13}C NMR spectrum, which suggested glycerol chain. Thus, **1** was primarily determined to be a flavonophenylpropanetriol. The HMQC and HMBC 2D NMR spectral analyses confirmed that the three phenol hydroxyl groups were situated at C-5, C-7 and C-4'', and three methoxy groups at C-3', C-5' and C-3'' (see **Table 1**). From HMBC spectrum, the correlations H-7'' with C-1'' and C-4' were observed,

which suggested that C-7'' and C-4' were linked its *O* bond. The ¹H-¹H COSY spectra revealed the coupling patterns of the oxygenated methyne and methane protons of H-7'', H-8'', and H-9''. The relatively small constant between H-8'' and H-7'' (*J* = 5.0 Hz) suggested an erythro-configuration in the glycerol chain.¹¹⁻¹² Thus, lycopodone (**1**) was formulated as *rel*-(7''*R*, 8''*S*) - tricin-4'-*O*-(4''-hydroxy-3''-methoxyphenyl propanetriol).

EXPERIMENTAL

General Experimental Procedures. Column chromatography (CC): Silica gel (200-300 mesh, Qingdao Marine Chemical, China); and Lichroprep RP-18 (40-63μm, Merck, Darmstadt, German). All melting points: YANACO-MP-52 apparatus, uncorrected. Optical rotations: Horiba SEAP-300 spectropolarimeter. IR spectra: Shimadzu IR-450 instrument; in cm⁻¹; KBr pellets. UV spectral data: UV 210A spectrometer. FAB-MS and HRFAB-MS: VG-AUTOSPEC-3000 spectrometer; in *m/z* (rel. int. in % of the base peak). NMR spectra: Bruker AV-400, or DRX-500 instruments; Chemical shifts (δ) in ppm; TMS as the internal standard. Fractions were monitored by TLC, and spots were visualized by heating TLC sprayed with 10% H₂SO₄.

Plant Material. The whole body of *L. japonicum* Thunb was obtained from the Chinese herbal market. It was identified by Prof. S. G. Wu. A voucher specimen (KUN No. 001143) was deposited at the Laboratory of Phytochemistry, Kunming Institute of Botany.

Extraction and Isolation. The dried, milled whole body of *L. japonicum* (19.0 kg) was exhaustively extracted with 90% MeOH (3×10 L) under reflux for 4, 4, 3 h. The MeOH extract was evaporated under reduce pressure to yield syrup (910 g), which was suspended in water: MeOH (9:1, 1500 mL) and extracted successively with EtOAc (3×2000 mL) and n-BuOH part (3×2000 mL) to give EtOAc-soluble (410 g) and n-BuOH-soluble fractions (101 g). The AcOEt extract was absorbed on 600 g silica gel and was fractionated by CC [silica gel (1.5 kg), CHCl₃ : Me₂CO 10:0, 9:1, 8:2, 7:3, 6:4 0:10, *v:v*] to afford six fractions (*Fr.*): 1 (oil), 2 (104 g), 3 (119 g), 4 (64 g), 5 (49 g), 6 (22 g).

Fr. 2 (104 g) was subjected to CC (silica gel, CHCl₃ : MeOH 40:1, 30:1, 20:1, *v:v*) to get four

subfractions. *Fr. 2.1* was further purified by repeated CC (silica gel, CHCl₃: MeOH 60:1, 50:1, *v:v*) to yield **3** (30 mg). *Fr. 2.2* was dissolved in CHCl₃ and yielded yellow powder of compound (**2**) (10 mg). *Fr.2.4* was further purified by CC (silica gel, CHCl₃ : MeOH 35:1, *v:v*) to give **4** (40 mg).

Table 1 ¹³C and ¹H NMR spectra data of compound of **1**

No	¹³ C	¹ H	HMBC	COSY
2	162.9 s	/	H-3, H-2', H-6'	
3	104.7 d	7.00 (s)		
4	181.7 s	/	H-3, H-6	
5	161.3 s	/	H-6, 5-OH	
6	98.8 d	6.70 (d, 2.0)	H-8	
7	164.2 s	/	H-6, H-8	
8	94.2 d	6.22 (d, 2.0)	H-6	
9	157.3 s	/	H-8	
10	103.8 s	/	H-6, H-8, H-3, 5-OH	
1'	125.2 s	/	H-3, H-2', H-6'	
2'	104.4 d	7.30 (s)	H-6'	
3'	152.9 s	/	3'-OMe, H-2'	
4'	139.5 s	/	H-2', H-6', H-7''	
5'	152.9 s	/	5'-OMe, H-6'	
6'	104.4 d	7.30 (s)	H-2'	
3',5'-OMe	56.3 q	3.76 (s, 6H)		
1''	133.1 s	/	H-2'', H-6''	
2''	111.2 d	6.95 (d, 1.8)	H-6'', H-8''	
3''	146.9 s	/	3''-OMe, H-2'', H-5''	
4''	145.4 s	/	4''-OH, H-5''	
5''	119.4 d	6.76 (d, 8.8)	H-6'', H-8'', 4''-OH	
6''	114.7 d	6.70 (dd, 1.8, 8.8)	H-2'', H-5'', 4''-OH	
7''	86.3 d	4.36 (br s)	H-8'', H-9''	H-8''
8''	72.2 d	4.80 (br s)	H-7'', H-9''	H-7'', H-9''

9''	60.1 t	3.51 (d, 9.3)	H-8'', H-7''	H-8''
3''-OMe	55.6 q	3.65 (s, 3H)		
5-OH		12.83 (br s, 1H)		
4''-OH		8.62 (br s, 1H)		
8''-OH		5.07 (br s, 1H)		
9''-OH		4.06 (br s, 1H)		

Recorded at 500 MHz in DMSO-d₆, at room temperature.

Fr. 3 (119 g) was purified by repeated CC (silica gel, CHCl₃ : MeOH 25:1, 15:1, 10:1, v:v) to afford five fractions. *Fr.3.2* (200.0 mg) was a mixture of compound (**1**) and other compounds. And the mixture was repeatedly subjected to HPLC (MeOH : H₂O 70:30, v:v) to yield compound (**1**) (50.0 mg).

Compound (**1**), yellow powder, mp>300°C; $[\alpha]_D^{23} - 11.76^\circ$ (*c* 1.7, MeOH); UV $\lambda_{\max}^{\text{MeOH}}$ nm(log ϵ): 204(4.16), 273(3.55), 306(3.53), 331(3.55); ¹³C and ¹H NMR spectrum, see **Table 1**; IR (KBr) ν_{\max} 3417 (br.s), 2941, 2842, 1656, 1612, 1591, 1556, 1455, 1356, 1250, 1170, 1048, 1031, 989, 918, 838, 786 cm⁻¹; FAB-MS (*m/z*, %): 525([M-1]⁻, 55), 329([M-C₁₀H₁₃O₄], 100); HRFAB-MS: 525.0867 (calcd for C₂₇H₂₅O₁₁ 525.1396)

Antitumor Activity: Human tumor A549, K562 cells line assay were performed at Kunming Medical College prof. Q. Chen and her coworkers and previously described bioassay methods were adopted.¹⁵⁻¹⁷ The IC₅₀ values for compounds (**1**) and (**2**) (10-100 µg/mL, 11.68 µg/mL, respectively, against human tumor K562 cells) indicated moderate antitumor activity.

ACKNOWLEDGEMENTS

This work was supported by NKIP Foundation of CAS (KSCZX-SW-301-08) and Foundation of Key State Lab of Phytochemistry.

REFERENCES

1. C. Y. Yang, *Zhongyao Tongbao*, 1981, **6**, 12.

2. K. M. Dai, D. J. Pan, Z. H. Chen, and X. Cai, *Journal of Plant Resource and Environment*, 1992, **1**, 36.
3. Jingsu Institute of Botany, *Outline of New China Herbal*, Shanghai: The Shanghai Science and Technology Press, 1990, p. 625
4. H. Zhou, C. H. Tan, S. H. Tan, and D. Y. Zhou, *J. Nat. Prod.*, 2003, **66**, 1328.
5. M. J. Kulshreshtha, D. K. Kushireshta, and R. P. Rastogi, *Phytochemistry*, 1972, **11**, 2369.
6. P. Pant and R. P. Rastogi, *Phytochemistry*, 1979, **18**, 1095.
7. F. R. Ansair, W. H. Ansari, W. Rahman, O. Seligman, V. M. Chari, H. Wagner, and B. G. Osterdahl, *Planta Medica*, 1979, **36**, 196.
8. W. F. Zhang, R. X. Tan, L. Yang, and Z. L. Liu, *Planta Medica*, 1996, **62**, 160.
9. K. H. Lee, K. Tagahara, H. Suzuki, R. Y. Wu, M. Haruna, and I. H. Hall, *J. Nat. Prod.*, 1981, **44**, 530.
10. G. R. Pettit, Y. H. Mei, C. A. Stevenson, D. L. Doubek, J. C. Knight, Z. Cichacz, R. K. Pettit, J. C. Chapuis, and J. M. Schmidt, *J. Nat. Prod.*, 2003, **66**, 259.
11. K. R. Markham, B. Ternal, R. Stanley, H. Gelger, and T. J. Mabry, *Tetrahedron*, 1978, **34**, 1389.
12. M. Fujita, M. Nagai, and T. Inoue, *Chem. Pharm. Bull.*, 1982, **30**, 1151.
13. J. M. Fang, C. K. Lee, and Y. S. Cheng, *Phytochemistry*, 1992, **31**, 3659.
14. M. L. Cardona, B. Carcia, J. R. Pedro, and J. F. Sinisterra, *Phytochemistry*, 1990, **29**, 629.
15. T. J. Mosmman, *Immunol. Meth.*, 1983, **65**, 55.
16. M. C. Alley, D. A. Scudiero, and A. Monks, *Cancer Res.*, 1988, **48**, 589.
17. J. J. Zhou, X. F. Yue, and J. X. Han, *Chin. J. Pharm.*, 1993, **24**, 455.