

HETEROCYCLES, Vol. 81, No. 10, 2010, pp. 2343 - 2350. © The Japan Institute of Heterocyclic Chemistry
Received, 16th June, 2010, Accepted, 10th August, 2010, Published online, 10th August, 2010
DOI: 10.3987/COM-10-11996

SUBSTITUENT EFFECT OF 7-HYDROXY-1-METHYLQUINOLINIUM DERIVATIVES: A PHOTOCHEMICAL APPROACH TO DEVELOPMENT OF NEW FLUORESCENT pH INDICATOR

Naoko Senda, Atsuya Momotake, and Tatsuo Arai*


Graduate School of Pure and Applied Sciences, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki, 305-8571, Japan. Fax: +81 29-853-6503; Tel: +81 29-853-4315; E-mail: arai@chem.tsukuba.ac.jp

Abstract – Substituent effect of *N*-methyl-7-hydroxyquinolinium derivatives on photochemical properties in aqueous solution has been investigated. The efficiency of fluorescence emission from excited zwitter ion enhanced significantly by introduction of methyl group(s). pH Dependence of photochemical behavior of the quinolinium derivatives is also investigated.

7-Hydroxyquinoline (**HQ**) is known to form a hydrogen-bonding chain with protic solvents, through which proton or hydrogen atom relay takes place in the excited singlet state.¹⁻³ The solvent assisted proton relay mechanism in this type of molecule has been of interest because the processes may serve as simple models for proton migration in biological systems. However the behavior is slightly complicated, since **HQ** can take four prototropic species including a normal form (**N**), an *N*-protonated cation form (**C**), a OH-deprotonated anion form (**A**), and an *N*-protonated and OH-deprotonated zwitter ion form (**Z**) (Scheme 1a), and these four species are in equilibrium in neutral aqueous solution.⁴⁻⁶ When the quinolinium nitrogen in **HQ** is functionalized by treatment with such methyl iodide, the equilibrium can be simplified: 7-hydroxy-1-methylquinolinium iodide (**1**) can take only two prototropic species, a cation form (**C**) at acidic pH and a zwitter ion form (**Z**) at neutral to basic pH (Scheme 1b).^{7,8} The 7-hydroxyl group of **1** was found to exhibit a strong photoacidity. For example, excited-state protolysis can occur even in the presence of 10 M perchloric acid.⁷ The electron-withdrawing quinolinium cation part probably amplify the photoacidity.

Quinolinium iodide **1** shows the biological activity. For example, applying a series of isomeric 1-methylhydroxyquinolinium ions to the electroplax, **1** was found to be a stronger receptor inhibitor compared to 1-methylquinolinium,⁹ where 7-hydroxyl group of **1** is expected for the formation of a

hydrogen bond to the receptor.¹⁰ Compound **1** is also used as a fluorescent probe, by monitoring the fluorescence intensity upon progressing of an acetylcholinesterase-catalyzed hydrolysis of either 7-acetoxy-1-methylquinolinium iodides¹¹⁻¹³ or 7-dimethylcarbamoy-1-methylquinolinium iodide.¹⁴



Scheme 1

Very recently, we have reported the new HQ-based fluorescent dye, Tsukuba-Green (**TG**), 4-ethoxycarbonylmethyl-7-hydroxy-2-methylquinoline (Figure 1).¹⁵ Photosensitivity, product of the molar extinction coefficient and fluorescence quantum yield ($\epsilon \cdot \Phi_F$) of **TG** is 4-5 fold greater than that of the parent compound **HQ**. During the course of the study we have found that both acylation of 7-OH and *N*-methylation of **TG** (**TG'-Ac**) is essential to penetrate the cell membrane. Since **TG'-Ac** is non-fluorescent, observation of immediate staining of HEP-2 cells after treatment of **TG'-Ac** indicates that **TG'-Ac** passed rapidly through the plasma membrane followed by hydrolysis via intracellular esterase to give quinolinium ion **TG'** which was highly fluorescent. In fact, $\epsilon \cdot \Phi_F$ value of the




Figure 1. Chemical structures of 7-hydroxy-1-methylquinolinium iodide **1** and its analogues **2** and **3**

quinolinium ion **TG'** ($\epsilon \cdot \Phi_F = 4200$) is more than 2-fold larger than that of quinolinium ion **1** ($\epsilon \cdot \Phi_F = 2000$). This significant enhancement of photosensitivity led us to assume the favorable effect of alkyl substituents on its photosensitivity of **1**.

In this paper we wish to report the substituent effect of 7-hydroxy-1-methylquinolinium ions on its photosensitivity. On this aspect, two methyl analogues, 7-hydroxy-1,2-dimethylquinolinium iodide **2** and 7-hydroxy-1,2,4-trimethylquinolinium iodide **3** (Figure 1) are prepared and their photochemical properties are compared to the parent compound **1**. In addition pH dependence of photochemical behavior of **1-3** is also investigated.

7-Hydroxy-1-methylquinolinium iodides **1**,¹⁵ **2**, **3**¹⁶ were prepared from **HQ**, 7-hydroxy-2-methylquinoline and 7-hydroxy-2,4-dimethylquinoline,¹⁷ respectively, by treatment with methyl iodide. The preparation of the starting materials and photochemical data of **1** have been reported elsewhere.¹⁵

Figure 2 shows the steady-state absorption (2a) and fluorescence (2b) spectra of **1-3** in water (1% DMSO). Compare to the absorption band of **1**, slight hypsochromic shift and a small shoulder at 360 nm were observed in **2**, and these features were prominent in **3**. The ϵ values at the maximum wavelength at pH




Figure 2. Absorption (a) and fluorescence spectra (b) of **1** (thin line), **2** (dash line), and **3** (solid line) in water (1% DMSO)

Table 1. Absorption maxima, Molar extinction coefficient, Fluorescence maxima, Fluorescence quantum yield, Fluorescence lifetime, Fluorescence rate constant, Non-radiative rate constant, and pK_a at 7-hydroxy group of *N*-methyl quinolinium iodides **1-3**

	$\lambda_{\text{abs}}/\text{nm}$	$\varepsilon/\text{M}^{-1}\text{cm}^{-1}$	$\lambda_{\text{fl}}/\text{nm}$	Φ_f	$\varepsilon \cdot \Phi_f$	τ/ns	$k_f/10^7\text{ s}^{-1}$	$k_d/10^8\text{ s}^{-1}$	pK_a
1 ^a	405	10300	517	0.19	2000	7.4	2.6	1.1	6.09
2	402	8400	500	0.36	3000	9.1	4.0	0.70	6.25
3	397	8100	496	0.45	3700	10.0	4.5	0.55	6.45

a) Photophysical data of **1** have been reported in ref. 15.

7.0 were 10300, 8400, and 8100 $\text{M}^{-1}\text{cm}^{-1}$, for **1**, **2**, and **3**, respectively. The spectral difference is caused by the difference of pK_a values at 7-hydroxyl group: pK_a values of **1**, **2** and **3** were estimated to be 6.09, 6.25, and 6.45, respectively, showing that the pK_a values increased by introduction of methyl group(s) (Table 1). At pH 7.0, **1** mostly takes **Z** form, while population of **C** form is not negligible for **2** and **3**. The formation of **C** form of **3** is more significant at pH 7.0 due to its higher pK_a value than those of **1** and **2**, resulted in the smaller ε value at the band for **Z** form. As in Figure 2b, a single fluorescence band around 500 nm appeared for each compound, ascribed to emission from the excited **Z** form.⁶ The fluorescence slightly shifted to shorter wavelength from 517 nm for **1** to 500 and 496 nm for **2** and **3**, respectively. In each compound, the fluorescence spectra by excitation at different wavelength were almost identical, and all three compounds showed a single fluorescence decay curve. It is noteworthy that the quantum yield of fluorescence of **2** ($\Phi_f = 0.36$) and **3** ($\Phi_f = 0.45$) are much higher than that of **1** ($\Phi_f = 0.19$). The fluorescence lifetime of **2** ($\tau = 9.1$ ns) and **3** ($\tau = 10.0$ ns) are also longer compared to that for **1** ($\tau = 7.4$ ns). One can calculate the fluorescence rate constant k_f and non-radiative rate constant k_d from the observed τ and Φ_f (Table 1). While the value of the fluorescence rate constant k_f increased in the order of **1**, **2**, and **3**, the other non-radiative rate constant k_d decreased in the same order. These facts indicate that the introduction of methyl group increased the k_f value and decreased the k_d value in 7-hydroxy-1-methylquinolinium iodides. This should be the reason why Φ_f value increased in the order of **1**, **2**, and **3**. The photosensitivity, the product of the molar extinction coefficient and fluorescence quantum yield ($\varepsilon \cdot \Phi_f$) for **2** ($\varepsilon \cdot \Phi_f = 3000$) and **3** ($\varepsilon \cdot \Phi_f = 3700$), respectively, is about 1.5-1.9-fold greater than that of **1** ($\varepsilon \cdot \Phi_f = 2000$) (Table 1).

The absorption spectra of all three compounds changed depending on pH, while the fluorescence spectra are almost identical at various pH values. The apparent absorption spectra of **1**, **2**, and **3** in aqueous buffered solution at various pH are shown in Figure 3. At pH 7.0 (solid line in Figure 3a-3c), the absorption spectra of each compound correspond with those in Figure 2a. The absorption spectra at pH




Figure 3. pH dependence of the absorption spectra of **1** (a), **2** (b), and **3** (c) at pH 2.0 (closed circle), pH 7.0 (solid line), and pH 12.0 (opened circle), respectively. Inset of each figure shows the normalized fluorescence spectra (excited at 350 nm) at pH 2.0 (closed circle), pH 7.0 (solid line), and pH 12.0 (opened circle), respectively

2.0 (closed circle) are dominated by **C** form, and those at pH 12 (opened circle) are due to **Z** form. The spectra of **2** and **3** at pH = 7.0 are composed of both **C** and **Z** forms. The pH dependence of absorption spectra of **2** and **3** are shown in Figure S1. Unlike the pH dependence observed in the absorption, there is practically no change in the fluorescence spectra of each compound (Inset of Figure 3) measured in acidic (pH = 2), neutral (pH = 7), and basic (pH = 12). In addition, the fluorescence quantum yield of all compounds did not depend on pH. Furthermore, the fluorescence excitation spectra of each compound are very similar to the corresponding absorption spectra, and are also changed with changing pH in similar manner (Figure S2). This indicates that the structure of the emission state does not depend on





Figure 4. pH dependence of the fluorescence spectra of **3**. Excitation wavelength is 400 nm

pH at least between pH = 2 and pH = 12, because the excitation of these compounds renders the 7-hydroxyl group to be more acidic than that in the ground state. The schematic representation of the structural change of **1** in different pH is shown in Scheme 2.

Since intracellular pH is generally between 6.8-7.4 in the cytosol, these dyes may be a potential candidate for pH indicator for inside live cells. Remarkably, the pKa values of **3** (Table 1) is appropriate for detection of intracellular pH change. In fact, excitation of **3** at 400 nm light displayed a dramatic change in the fluorescence especially between pH 6.0-7.5 (Figure 4) which is close to the cytosolic pH. This change in the fluorescence intensity probably is large enough to detect intracellular pH changes. In addition, the absorption wavelength of **3** is different from similar fluorescein-based fluorescence pH indicators, such as BCECF and SNAFL that are currently used for live cell imaging,¹⁸ indicating that **3**

could be used for multi-color imaging together with fluorescein-based probes. To load this dye into cells, however, further chemical modification, such as for enhancement of membrane permeability and retention inside cells, would be needed.

In summary, substituent effect of 7-hydroxy-1-methylquinolinium ions has been investigated. The efficiency of fluorescence emission from excited state **Z** form is enhanced significantly by introduction of methyl group(s). In aqueous solution, the products of the molar extinction coefficient and fluorescence quantum yield ($\epsilon \cdot \Phi_F$) for **2** and **3**, respectively, are 1.5- and 1.9-fold greater than that of **1**. The lifetime of fluorescence increased with increasing the fluorescence quantum yield from 7.4 ns in **1** to 9.1 ns and 10.0 ns in **2** and **3**. The pH dependence of the absorption spectra was observed and the pK_a values at 7-hydroxy group increased by the effect of methyl group(s). Spectral features in the fluorescence spectra or fluorescence quantum yield did not show pH dependence (from pH 2 to 12) suggesting the same structure in emission state despite the different ground-state structure at various pH. These results presented in this paper indicate that **2** and **3** are considered to be good candidates as fluorescent chromophores for bioimaging. Further investigation for biological use of these chromophores is underway.

EXPERIMENTAL

Measurement

^1H NMR spectra were measured with a JEOL EX-270 (270 MHz for ^1H NMR) or a Bruker ARX-400 (400 MHz for ^1H NMR) spectrometer using CD_3OD as a solvent. UV absorption and fluorescence spectra were recorded on a Shimadzu UV-1600 spectrophotometer and on a Hitachi F-4500 fluorescence spectrometer, respectively. Fluorescence decay measurements were performed using time-correlated single-photon counting.

Materials

Syntheses of **1** and **3** were written in elsewhere.^{15, 16}

7-Hydroxy-1,2-dimethylquinolinium iodide (2): A mixture of 2-methyl-7-hydroxyquiniline (107 mg, 0.67 mmol) and methyl iodide (1.0 mL, 0.70 mmol) was heated in a sealed tube at 65 °C for 23 h. The precipitate formed was collected by filtration, washed thoroughly with dichloromethane to give 94 mg of **2** in 47 % yield; ^1H NMR (270 MHz, CD_3OD) δ 8.78 (d, $J = 8.1$ Hz, 1H), 8.17 (d, $J = 8.6$ Hz, 1H), 7.71 (d, $J = 8.4$ Hz, 1H), 7.56 (s, 1H), 7.47 (dd, $J = 1.9, 8.9$ Hz, 1H), 4.34 (s, 3H), 3.04 (s, 3H) ^{13}C NMR (67.5 MHz, CD_3OD) δ 165.9, 160.2, 146.0, 143.8, 133.7, 124.4, 122.2, 122.1, 101.9, 39.9, 23.4. Anal. Calcd

for C₁₁H₁₂NOI: C, 43.87; H, 4.02; N, 4.65 %. Found: C, 43.68; H, 3.90; N, 4.55 %.

ACKNOWLEDGEMENTS

This work was supported by a Grant-in-Aid for Scientific Research in a Priority Area "New Frontiers in Photochromism (No. 471) and (No. 19550176) from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan.

REFERENCES

1. W.-H. Fang, *J. Am. Chem. Soc.*, 1998, **120**, 7568.
2. C. Tanner, C. Manca, and S. Leutwyler, *Science*, 2003, **302**, 1736.
3. H. Yu, O.-H. Kwon, and D.-J. Jang, *J. Phys. Chem. A*, 2004, **108**, 5932.
4. S. I. Lee and D. J. Jang, *J. Phys. Chem.*, 1995, **99**, 7537.
5. T. G. Kim, Y. R. Kim, and D.-J. Jang, *J. Phys. Chem. A*, 2001, **105**, 4328.
6. T. G. Kim and M. R. Topp, *J. Phys. Chem. A*, 2004, **108**, 10060.
7. E. Bardez, A. Chatelain, B. Larrey, and B. Valeur, *J. Phys. Chem.*, 1994, **98**, 2357.
8. E. Bardez, A. Fedorov, M. N. Berberan-Santos, and J. M. G. Martinho, *J. Phys. Chem. A*, 1999, **103**, 4131.
9. T. R. Podleski and D. Nachmansohn, *Proc. Nat. Acad. Sci.*, 1966, **56**, 1034.
10. T. R. Podleski, *Proc. Nat. Acad. Sci.*, 1967, **57**, 268.
11. A. K. Prince, *Arch. Biochem. Biophys.*, 1966, **113**, 195.
12. A. K. Prince, *Biochem. Pharmacol.*, 1966, **15**, 411.
13. F. Caturla, J. Enjo, M. C. Bernabeu, and S. L. Serre, *Tetrahedron*, 2004, **60**, 1903.
14. T. L. Rosenberry and S. A. Bernhardt, *Biochemistry*, 1971, **10**, 4114.
15. N. Senda, Y. Miwa, J. Tanaka, A. Momotake, and T. Arai, *Chem. Lett.*, 2010, **39**, 308.
16. H. Lempert and R. Robinson, *J. Chem. Soc.*, 1934, 1419.
17. O. D. Fedoryak and T. M. Dore, *Org. Lett.*, 2002, **4**, 3419.
18. R. P. Haugland, *Handbook of Fluorescent Probes and Research Products* (Molecular Probes, Eugene, OR, ed.9, 2002).