

HETEROCYCLES, Vol. 82, No. 1, 2010, pp. 775 - 790. © The Japan Institute of Heterocyclic Chemistry
Received, 22nd June, 2010, Accepted, 15th July, 2010, Published online, 16th July, 2010
DOI: 10.3987/COM-10-S(E)65

SCREENING THE STRUCTURAL SPACE OF BICYCLO-DNA: SYNTHESIS AND PROPERTIES OF BICYCLO-DNA FUNCTIONALIZED AT C(6')

Samuel Luisier and Christian J. Leumann*

Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3,
CH-3012 Bern, Switzerland, leumann@ioc.unibe.ch

This paper is dedicated to Professor Dr. Albert Eschenmoser on the occasion of his 85th birthday in full appreciation for his scientific achievements and his outstanding intellectual leadership in introducing us into science.

Abstract – The synthesis of a novel bicyclo-thymidine nucleoside bearing an ester functionality at C(6') (bc^{α-alk}-nucleosides) is reported. This nucleoside was incorporated into oligodeoxynucleotides via solid phase phosphoramidite chemistry, and the ester moiety was post-synthetically converted to an amide or a carboxy group, or was left unchanged. Thermal melting data (T_m) with complementary DNA and RNA were collected and compared to natural DNA and to bc- and bc^{ox}-DNA. It was found that single incorporations of bc^{α-alk}-nucleosides in DNA duplexes were destabilizing by 0.5 to 2.5 °C/mod, whereas two consecutive bc^{α-alk}-residues were less destabilizing, and in some cases even stabilizing by 0.5 °C/mod. In duplexes with complementary RNA, isolated bc^{α-alk}-residues destabilized the duplex by -1.0 to -4.0 °C/mod, depending on the chemical nature of the substituent, whereas two consecutive modifications were only destabilizing by 0.3-1.0 °C/mod. The pairing selectivity was similar to that of unmodified or bc-DNA.

INTRODUCTION

The concept of conformational restriction has successfully been applied to nucleic acids and resulted in several nucleic acid analogues, such as locked-nucleic acids (LNA),^{1,2} hexitol nucleic acids (HNA),³ and tricyclo-DNA (tc-DNA, Figure 1).^{4,5} These analogues typically exhibit increased affinity to RNA and feature higher nuclease resistance. Due to their properties, such modified nucleic acids are considered as

next generation therapeutic drugs.^{6,7} While the challenges linked to target affinity and nuclease resistance have largely been met in the past, several problems, the most prominent ones being cellular uptake and distribution, largely remained unsolved.

Bicyclo-DNA (bc-DNA),⁸ was developed as a first generation conformationally constrained oligonucleotide to enhance duplex stability with natural nucleic acids. It turned out, however, that bc-DNA base-pairs to natural nucleic acids only with about equal stability as DNA itself.⁹ Extensive structural and synthetic investigations suggested that the lack of increased affinity may be due to the structural misalignment of torsion angle γ (C(4')-C(5')) with respect to that occurring in duplexes of the A- and B-conformational type.^{10,11} One of the unique structural features of bc-DNA is the ethylene bridge between the centers C(3') and C(5') of the ribose unit which lends itself for further chemical substitution. We reasoned that additional substituents at position C(6') could be interesting e.g. for controlling the conformation of the carbocyclic ring, eventually correcting the misaligned torsion angle γ . In addition, incorporation of a post-synthetically transformable group could be used either to attach reporter groups or to improve the lipophilic character of bc-DNA, thus improving its cellular uptake properties. Earlier work on C(6')-oxime modified bc-DNA (bc^{ox}-DNA) was encouraging in this context.¹²

One candidate of interest to be synthesized and tested is bc ^{α -alk}-DNA (Figure 1). Here we report on the synthesis of the bc ^{α -alk}-T nucleoside carrying a C2-carboxyl substituent in α -position at C(6'), on its conformational preference as explored by molecular modeling, on its incorporation into oligonucleotides and on the base-pairing properties with DNA and RNA.




Figure 1. bicyclo-DNA derivatives that have been investigated so far, as well as bc ^{α -alk}-modified DNA the synthesis and pairing properties of which are subject of this paper.

RESULTS AND DISCUSSION

Modeling of nucleosides and oligonucleotides. A conformational search of $bc^{\alpha\text{-alk}}$ -T **9** using *Hyperchem* yielded two low energy conformers (Table 1) with virtually identical furanose pucker (C_1' -exo) but with different cyclopentane ring conformations. In the lowest energy conformer, both the $C(5')$ and $C(6')$ substituents were in a pseudo-equatorial arrangement giving rise to a 6'-endo-conformation of the cyclopentane unit (Figure 2a, left). In the second conformer (Figure 2a, right), which is higher in energy by 2.36 kcal/mol, both the $C(5')$ and $C(6')$ substituents were in a pseudo-axial arrangement resulting in a $C(6')$ -exo arrangement of the carbocyclic ring. The relevant backbone torsion angles γ and δ of the two conformers are summarized in Table 1.

Modeling of a DNA-duplex containing one $bc^{\alpha\text{-alk}}$ -T nucleotide in the $C(1')$ -exo/ $C(6')$ -exo conformation resulted in a structure with only minor distortions from an ideal B-conformation at the site of modification (Figure 2b). The $C(6')$ substituent is pointing away from the minor and major groove, directly into the solvent. Thus no interference of this substituent with base-pairing is to be expected which makes $C(6')$ a perfect position for the attachment of additional functional groups.





Figure 2. a) The two low energy conformers found for $bc^{\alpha\text{-alk}}$ -T nucleoside; b) stereo view of a DNA duplex model containing one $bc^{\alpha\text{-alk}}$ -T residue (yellow) in one DNA strand.

Table 1. conformational characteristics, including torsion angles γ and δ (Figure 1), of low energy $bc^{\alpha\text{-alk}}$ -T conformers.

$bc^{\alpha\text{-alk}}$ -T	ΔE (kcal/mol)	Bicyclic pucker	γ	δ
Conf 1	0	$C(1')$ -exo/ $C(6')$ -endo	147°	118°
Conf 2	2.36	$C(1')$ -exo/ $C(6')$ -exo	98°	118°

Synthesis of C(6')-alkyl bicyclonucleosides. We intended to synthesize nucleoside **9** starting from the already known ketone **1** via cyclopropanation of the corresponding silylenol ether **2** or **3** and subsequent ring opening (Scheme 1). Previous experiments showed that cyclopropanation of silylenol ethers derived from ketone **1** under Simmons-Smith conditions¹³ yielded cyclopropanes in good yields and excellent selectivity.¹⁴ We thus prepared the triethylsilyl (TES)-enol ether by treatment of the anomeric mixture of **1** (α : β 4:1) with LDA followed by TES-Cl. Besides the desired silylenol ether **2**, some bis-silylated products could be identified in minor quantities. Preliminary experiments towards cyclopropanation of **2** with ethyl diazoacetate (EDA) and Cu(II) catalysts¹⁵ showed that carbene insertion into the tertiary OH-function occurred to a non-negligible extent. To suppress this side reaction enol ether **2** was silylated to bis-silylether **3** by treatment with bis(trimethylsilyl) acetamide in pyridine. Compound **3** was then cyclopropanated with EDA, which lead to three isomeric cyclopropanes **4a-c**, whose relative configuration at C(5') and C(6') could not unambiguously be assigned. Desilylation of the major isomer, **4a**, induced cyclopropane ring opening leading to the mixture of isomers of ketone **5**. Thus, during desilylation keto-enol tautomerisation takes place leading to epimerization at C(6'). In later experiments we used **4a-c** as a mixture of isomers for this reaction, thus reducing the complexity of the mixture from 3 to 2 isomers on the level of ketone **5**. Reduction of the isomeric ketones **5** in the presence of excess CeCl₃ yielded **6** and **7** in acceptable yields that are separable by column chromatography.



Scheme 1. Conditions: a) Li-DIPA, Et₃SiCl, THF, -78 °C, 2h, 54%; b) BSA, pyridine, 17h, rt, 86%; c) N₂CHCOOEt, ClCH₂CH₂Cl, Cu(acac)₂, 90 °C, 9h, 80%; d) HF-pyridine, pyridine, rt, 1h, 79%; e) CeCl₃, NaBH₄, 15min, 0 °C, 57% (combined yield **6+7**).

To unambiguously assign the relative configuration at the centers C(6') and C(7') we performed ¹H-NMR difference-NOE experiments (Figure 3). The consistently large, mutual NOE-effects between the protons

at C(4') and C(5') indicated their *cis*-relationship in both isomers **6** and **7**. The relatively small NOE between C(5') and C(6') in the case of **6** allowed the assignment of a *trans* relationship for these protons which is corroborated by the relatively large NOE between these protons in isomer **7** indicating a *cis* relationship. We rationalize the stereoselectivity of the reduction step in **5** with the bicyclic nature of the carbon scaffold which favors hydride transfer to occur on the convex side of the system.





Figure 3. intensity of observed (mutual) $^1\text{H-NMR}$ difference NOE effects of the relevant protons used for assigning the *cis*- and *trans*-relationship of substituents in the carbocyclic ring in **6** and **7**.



Scheme 2. Synthesis of building block **11** and $^1\text{H-NMR}$ difference NOE effects observed for nucleoside **9**. Conditions: a) 2,6-lutidine, TMS-OTf, CH_2Cl_2 , 1h, rt, quant.; b) thymine, BSA, NIS, rt, 3h; c) AIBN, Bu_3SnH , toluene, 95°C , 2h; d) HF-pyridine, pyridine, rt, 4h, 13% from **6**; e) DMT-OTf, pyridine, rt, 13h, 72%; f) $(i\text{Pr}_2\text{N})\text{P}(\text{Cl})\text{OCH}_2\text{CH}_2\text{CN}$, CH_3CN , $(i\text{Pr})_2\text{NEt}$, rt, 1h, 60%.

The synthesis of nucleoside **9** and its building block for DNA synthesis **11** continued by treatment of **6** with 2,6-lutidine and trimethylsilyl triflate (TMS-OTf) to produce the glycal **8** in quantitative yields (Scheme 2). We envisaged to perform a β -selective nucleosidation via a two step procedure that had proven successful for tricyclo-nucleoside synthesis before.¹⁶ Thus we converted **8** to the corresponding





2'-iodinated nucleoside upon treatment with *in situ* silylated thymine and *N*-iodosuccinimide (NIS). The intermediate iodo-nucleoside was not isolated but directly reduced with Bu_3SnH . Desilylation of the crude product of this reaction finally yielded nucleoside **9** with only minor amounts <10% of the undesired α -anomer. The β -configuration of the anomeric center in **9** was confirmed by a strong NOE between the C(1')- and C(4')-protons in the $^1\text{H-NMR}$ NOE difference spectrum (Scheme 2). Likewise, the relative configurations of C(5') and C(6') could be confirmed again, as in the case of **6**. The relatively large coupling constant of 9.8 Hz together with the relatively low NOE between HC(5') and HC(6') speaks for a trans-diaxial arrangement which is in full agreement with the C(6')-endo conformation of the carbocyclic ring. Likewise. The coupling pattern of the HC(1') to the two HC(2') together with the relatively large NOE between HC(4') and HC(1') of 4.9% are very similar to that of the unmodified bicyclothymidine, for which the furanose ring was found to prefer the C(1')-exo conformation in solution and in the solid state.¹⁷ Thus the $^1\text{H-NMR}$ data are completely conform with a (C(1')-exo/C(6')-endo conformation which was also predicted to be the most stable one by modeling (Table 1). Selective tritylation of the sterically less hindered secondary OH in nucleoside **9** with 4,4'-dimethoxytrityl triflate (DMT-OTf) and subsequent phosphitylation of the tertiary alcohol of nucleoside **10** lead to the phosphoramidite building block **11**, ready for oligonucleotide synthesis.

Oligonucleotide synthesis. Two oligonucleotides, containing single or double, consecutive $\text{bc}^{\alpha\text{-alk}}\text{-T}$ nucleotides were synthesized on a 0.5 μmol scale by standard automated phosphoramidite chemistry. For incorporation of the modified nucleotides the standard coupling step was extended to 6 min. No further changes were necessary, and the coupling yields for the modified nucleosides were similar to those of unmodified nucleosides. After synthesis the solid supported crude oligonucleotides were split into 4 portions each, followed by separate deprotection and detachment from the solid support using different conditions. Standard ammonolysis with aqueous ammonia yielded oligonucleotides in which the ester function of the $\text{bc}^{\alpha\text{-alk}}\text{-T}$ residues were converted to the unsubstituted amides. Deprotection with 40% methylamine in an ethanol-water mixture (1:2) yielded *N*-methylamides. Hydrolysis with 1M KOH in water yielded the free carboxylic acid while treatment with the less nucleophilic benzylamine left the ester function unchanged. All oligonucleotides were purified and desalted by standard HPLC methods and analyzed by ESI-MS. Table 2 gives an overview over the isolated oligonucleotides and their determined masses. All were in agreement with the proposed structures. For comparison we also prepared the two reference oligonucleotides **Ref1** and **Ref2**, carrying unsubstituted bicyclo-T units.

T_m measurements. UV-melting curve analysis was performed at 260 nm with a cooling-heating-cooling cycle at a rate of 0.5 °C/min in standard saline buffer (10mM NaH_2PO_4 , 150 mM NaCl, pH 7.0). All curves within a cycle were superimposable thus ruling out non-equilibrium states. T_m data are

summarized in Table 3. Incorporation of single $bc^{\alpha\text{-alk}}\text{-T}$ nucleotides induce a destabilization of -2.5 to -0.6 °C per modification with complementary DNA and -0.7 to -3.9 °C with complementary RNA depending on the chemical nature of the substituent at C(6'). Increased lipophilicity of this substituent, as for the methylamide or ester function lowers the T_m slightly more than a less hydrophobic primary amide function. The carboxylate not unexpectedly destabilizes significantly due to increased charge repulsion in the duplex. However, incorporation of two consecutive $bc^{\alpha\text{-alk}}\text{-T}$ units slightly stabilize duplexes with complementary DNA and lead to less destabilization with complementary RNA, compared to the mono-substituted oligonucleotides. A comparison with oligonucleotides **Ref 1** and **Ref 2** shows that the substituents at C(6') are slightly depressing the T_m s with complementary DNA while they have negligible effects if RNA is the pairing complement. Taking into account that there is no major change in conformation between $bc\text{-T}$ and $bc^{\alpha\text{-alk}}\text{-T}$ we believe that the slight variations in T_m are not of structural origin but may reflect the altered hydration sphere between the phosphodiester functions as a consequence of the chemical nature of the substituent. Nevertheless, it appears that an α -substituent in position C(6') is generally well tolerated on the bicyclo-DNA skeleton.

Table 2. Deprotection conditions and mass spectrometric analysis of oligonucleotides **12-19**.

Oligonucleotides	Deprotection conditions	X	m/z calc	m/z found
Ref 1 d(GGATGTTCXCGA)	33% NH ₃ , 55°C, 8h		3702.5	3702.0
Ref 2 d(GGATGXXCTCGA)	33% NH ₃ , 55°C, 8h		3728.5	3728.3
12 d(GGATGTTCXCGA)	33% NH ₃ , 55°C, 8h		3759.5	3759.4
13 d(GGATGXXCTCGA)	33% NH ₃ , 55°C, 8h		3862.6	3843.2






14	40% MeNH ₂ , d(GGATGTTTCXCGA)	MeOH/H ₂ O 60°C, 8h		3773.6	3773.8
15	40% MeNH ₂ , d(GGATGXXCTCGA)	MeOH/H ₂ O 60°C, 8h		3870.7	3870.5
16	1M KOH, H ₂ O, d(GGATGTTTCXCGA)	60°C, 8h		3760.5	3760.8
17	1M KOH, H ₂ O, d(GGATGXXCTCGA)	60°C, 8h		3844.6	3844.2
18	25% BnNH ₂ , d(GGATGTTTCXCGA)	MeOH/H ₂ O, 60 °C, 8h		3788.6	3787.9

Table 3. T_m values [°C] from UV melting curves (260 nm) for modified dodecamer duplexes with complementary DNA and RNA. Conditions: 10 nM NaH₂PO₄, 150 nM NaCl, pH 7, c = 2 μM total strands.

	T_m vs DNA (°C, 260 nm) ^{a,b}	T_m vs RNA (°C, 260 nm) ^{a,b}
Ref1	49.0 (+1.5)	48.0 (-1.5)
Ref2	48.7 (+0.6)	49.0 (-0.3)
12	46.9 (-0.6)	46.3 (-3.2)
13	48.5 (+0.5)	48.2 (-0.7)
14	45 (-2.5)	46.0 (-3.5)
15	47.9 (+0.2)	n.d.
16	45.5 (-2.0)	46.0 (-3.5)
17	48.0 (+0.3)	47.5 (-1.0)
18	45.9 (-1.6)	45.6 (-3.9)

a) T_m of unmodified oligodeoxynucleotide: 47.5 °C (vs. DNA), 49.5 °C (vs RNA)

b) values in parenthesis are ΔT_m per modification

To get information on the selectivity of base recognition with $bc^{\alpha\text{-alk}}$ -DNA, T_m data of duplexes containing a mismatched base opposite to the modification were recorded (Table 4). All mismatched duplexes were destabilized compared to the matched duplexes. However, the degree of destabilization was lower than that observed for bc -DNA. The order of stability of mismatched base pairs is as follows: $bc^{\alpha\text{-alk}}\text{-T}\cdot\text{dT} < bc^{\alpha\text{-alk}}\text{-T}\cdot\text{dC} < bc^{\alpha\text{-alk}}\text{-T}\cdot\text{dG}$. Interestingly $bc\text{-T}\cdot\text{dT}$ and $bc^{\alpha\text{-alk}}\text{-T}\cdot\text{dT}$ are particularly destabilizing, in contrast to the natural $\text{dT}\cdot\text{dT}$ mismatch. The influence of the nature of the substituent at C(6') was negligible on mismatch destabilization with the exception of the carboxylate substituent that showed less discriminative power.

Table 4. T_m values ($^{\circ}\text{C}$) from UV melting curves (260 nm) for modified duplexes carrying a mismatched base opposite the modified nucleotides. Conditions: 10 mM NaH_2PO_4 , 150 mM NaCl , pH 7, $c = 2\mu\text{M}$.

Oligonucleotide ^{a,b}	C·T	G·T	T·T
DNA	36.0 (-11.5)	39.6 (-7.8)	38.0 (-9.5)
Ref 1	35.0 (-13.0)	37.0 (-11.0)	32.0 (-16.0)
12	36.0 (-10.3)	38.3 (-8.0)	33.3 (-13.0)
14	35.6 (-9.4)	39.0 (-7.0)	32.9 (-13.1)
16	35.6 (-9.9)	38.0 (-7.5)	33.9 (-11.6)
18	34.9 (-10.7)	37.2 (-8.4)	32.9 (-12.7)

a) values for matched duplex: see Table 1

b) values in parenthesis are ΔT_m values relative to matched duplex.

CONCLUSIONS

We have synthesized a bicyclic nucleoside having a post-synthetically transformable ester function attached to its carbocyclic core. This bicyclic nucleoside was successfully incorporated into oligonucleotides by standard phosphoramidite chemistry. Depending on the deprotection conditions, the original ester function could be transformed in amides and carboxylates or left unchanged. Single incorporations of this modification in oligodeoxynucleotides lead to a slight drop in duplex stability. However, duplexes containing two consecutive modifications exhibited similar or slightly enhanced stabilities compared with duplexes containing parent bc -DNA units or natural deoxynucleotides. No critical differences in stability caused by the different chemical nature of the substituents could be observed. This novel convertible building block **11** may be of interest in the future for attaching fluorescent reporter groups onto oligonucleotides or for post-synthetic derivatization with chemical entities that enhance cellular uptake.

EXPERIMENTAL

All reactions were performed under Ar in dried glassware. Anhydrous solvents for reactions were obtained by filtration through activated aluminium oxide, or by storage over 4 Å molecular sieves. Column chromatography was performed on silica gel (Fluka) with an average particle size of 40 µm. All solvents for CC were of technical grade and distilled prior to use. Thin-layer chromatography (TLC) was performed on silica gel plates (Macherey-Nagel, 0.25 mm, UV254). Visualization was performed either by UV or by staining in dip solution (10.5 g Cer(IV)-sulfate, 21 g phosphomolybdic acid, 60 mL conc. sulfuric acid, 900 mL H₂O) followed by heating with a heat gun. NMR spectra were recorded on a Bruker DRX-400 or a Bruker AC-300 spectrometer at 400 MHz or 300 MHz (¹H-NMR) or 100 MHz (¹³C-NMR) in either CDCl₃ or CD₃OD. δ are given in ppm relative to residual undeuterated solvent (CHCl₃: 7.26 ppm (¹H) and 77.0 ppm (¹³C); CHD₂OD: 3.35 ppm (¹H) and 49.3 ppm (¹³C)), *J* in Hz. ¹³C-multiplicities were determined from DEPT-spectra and signal assignments are based on ¹³C/¹H-HMBC spectra. Proton signal assignments were based on COSY and HMBC. ¹H-NMR difference-NOE spectra were recorded on a Bruker DRX-500 instrument at 500 MHz. High resolution electrospray ionization (ESI) mass spectra (MS, *m/z*) were recorded on an Applied Biosystems Sciex QSTAR Pulsar instrument.

Molecular modeling

Conformational search of the nucleosides were performed using *Hyperchem* software, using the *Amber 2* force field and a *Polak-Riebere* gradient. All torsion angles on the bicyclic core were modulated and the bicyclic ring system was doubly defined: first as 8-membered ring, then as two five membered rings. A geometry optimization preceded the conformational search, and the 1000 lowest energy conformations were screened for relevant structures. Oligodeoxynucleotide duplexes were setup in B-conformation using the standard *Hyperchem* dataset. Modified nucleosides were constructed from the parent natural nucleosides. Geometry optimization was run using the *Amber 2* force field with a *Polak-Riebere* gradient.

Oligonucleotides synthesis

Oligonucleotide syntheses were performed with solid-phase phosphoramidite methodology on a Polygen DNA-synthesizer (1-µmol-slider). Ethyl thiotetrazole (0.25 M in MeCN) was used as activator in the coupling step. Commercial dA-CPG (50 µmol/mg) solid support was used. After deprotection, the crude oligonucleotide solutions were evaporated and dried on a *Savant Speed-Vac SC 110*.

HPLC

All oligonucleotides were purified by ion-exchange HPLC using an *ÄktaTMbasic 10/100* system (*Amersham Pharmacia Biotech*) on a *DNAPAC PA200* column (4 x250 mm, *Dionex*). All oligonucleotides were desalted after chromatography using *Sep-Pak Classic C18 Cartridges* (*Waters*) and were routinely analyzed by ESI mass spectrometry (see Table 3).

UV-Melting curves

UV-melting curves were carried out on a *Varian Cary 100Bio UV/Vis* spectrophotometer. Absorbances were monitored at 260 nm and the heating rate was set to 0.5 °C/min. A heating-cooling-heating cycle in the temperature range 15-80 °C was applied. The absorbance melting curves were smoothed and the first derivative curves obtained using the *Varian WinUV* software was used to determine the T_m . To avoid evaporation of the solution, the samples in the cells were covered with a layer of dimethylpolysiloxane. All measurements were carried out in phosphate buffered saline (150 mM NaCl, 10 mM NaH₂PO₄, pH 7.0). Measurements were carried out at a total strand concentration of 2 μM.

2-Methoxy-6-triethylsilyloxy-2,3,4,6a-tetrahydrocyclopenta[*b*]furan-3a-ol (2)

A solution of ketone **1** (α,β ca. 4:1, 494 mg, 2.87 mmol) in THF (5 mL) was added at -78 °C in 5 min to a solution of BuLi (3.6 mL 1.56 M, hexane, 6 mmol, 2 eq) and diisopropylamine (0.83 mL, 6 mmol, 2 eq) in THF (5 mL), followed by triethylchlorosilane (1.05 mL, 6.4 mmol, 2 eq) and TEA (0.5 mL) in THF (5 mL) within 10 min. After stirring for 2 h at -78 °C and 30 min at 0 °C, NH₄Cl (1 g) was added. After filtration over *Celite*, removal of the solvents followed by column chromatography (hexane/EtOAc 3:1), enol ether **2** (441 mg, 54%) was obtained as a colorless oil; TLC (hexane/EtOAc 3:1) R_f = 0.58; ¹H-NMR (400 MHz, CDCl₃) δ 5.06 (1H, d, J = 4.16 Hz), 4.66 (1H, s), 4.59 (1H, t, J = 2.44 Hz), 3.39 (3H, s), 3.19 (1H, s), 2.42 (2H, t, J = 1.84 Hz), 2.20 (1H, d, J = 13.32 Hz), 2.01 (1H, dd, J = 13.32, 4.04 Hz), 0.98 (9H, t, J = 8.08 Hz), 0.74-0.67 (6H, m); ¹³C-NMR (100 MHz, CDCl₃) δ 152.28 (s), 105.79 (d), 103.15 (d), 92.37 (d), 85.14 (s), 54.95 (q), 47.29 (t), 37.99 (t), 6.85 (q), 5.04 (t); HRMS (ESI⁺) m/z 309.1492 (M+Na⁺, C₁₄H₂₆O₄NaSi requires 309.1498).

2-Methoxy-6-triethylsilyloxy-3a-trimethylsilyloxy-,3a,4,6a-tetrahydro-2H-cyclopenta[*b*]furan (3)

BSA (0.56 mL, 2.3 mmol, 1eq) was added at rt to a solution of **2** (655 mg, 2.3 mmol) in pyridine (5 mL). After stirring for 12h at rt, a second portion of BSA (0.56 mL, 2.3 mmol, 1 eq) was added. After stirring for another 5h at rt, the solvents were removed by evaporation, yielding **3** (707 mg, 86%) as a slightly brownish oil; TLC (hexane/EtOAc 9:1) R_f = 0.9; ¹H-NMR (CDCl₃, 400MHz) δ 5.01 (1H, dd, J = 4.40, 0.88 Hz); 4.73 (1H, s), 4.60 (1H, t, J = 2.44, Hz), 3.37 (3H, s), 2.49 (1H, dt, J = 15.61, 2.09 Hz), 2.41 (1H, dd, J = 15.50, 1.45 Hz), 2.29 (1H, dd, J = 13.58, 0.71 Hz); 2.03 (1H, dd, J = 13.52, 4.57 Hz), 0.98 (9H, t, J = 7.77 Hz), 0.70 (6H, q, J = 7.96 Hz); 0.14 (9H, s); ¹³C-NMR (CDCl₃, 100MHz) δ 152.09 (s), 105.77 (d), 102.89 (d), 91.53 (d), 86.29 (s), 54.97 (q), 48.59 (t), 42.53 (t), 6.55 (q), 4.77 (t), 1.76 (q); HRMS (ESI⁺) m/z 381.1897 (M+Na⁺, C₁₇H₃₄O₄NaSi₂ requires 381.1893).

3-Methoxy-1a-triethylsilyloxy-4a-trimethylsilyloxyoctahydro-2-oxacyclopropa[*a*]pentalene-1-carboxylic acid ethyl ester (4)

A solution of ethyl diazoacetate (0.6 mL, 5.9 mmol, 3 eq) in CH₂Cl₂ (100 mL) was dropped at 90 °C over a period of 9h to a solution of **3** (0.7 g, 1.95 mmol) and Cu(acac)₂ (19 mg, 0.08 mmol, 0.04 eq) in CH₂Cl₂ (50 mL). During the addition the solvent was partly evaporated. Filtration over Celite and evaporation of the solvents followed by column chromatography (hexane/EtOAc 15:1 + 0.1 % TEA) yielded a 1.3 : 1 : 1 mixture of isomers **4a-c** (**4a**, 286 mg, 33%; **4b**, 217 mg, 25%; **4c**, 192 mg, 22%), all as colorless oils.

Data of **4a**: TLC (EtOAc) R_f = 0.88; ¹H-NMR (CDCl₃, 400MHz) δ 5.12 (1H, dd, J = 5.72, 2.20 Hz), 4.58 (1H, s), 4.23 - 4.12 (2H, m), 3.40 (3H, s), 2.27 (1H, dd, J = 13.96, 5.52 Hz), 2.22 (1H, d, J = 1.96 Hz), 2.17 (1H, dd, J = 13.56, 7.00 Hz), 2.06 (1H, d, J = 9.68 Hz), 1.95 (1H, dd, J = 13.72, 1.44 Hz), 1.88-1.83 (1H, m), 1.27 (3H, t, J = 7.07 Hz), 1.00 - 0.91 (9H, m), 0.72-0.67 (6H, m), 0.12 (9H, s); ¹³C-NMR (CDCl₃, 100MHz) δ 168.69 (s), 106.26 (d), 91.49 (s), 88.28 (d), 60.10 (t), 55.14 (q), 47.75 (t), 34.96 (t), 33.50 (d), 30.70 (d), 14.37 (q), 6.73 (q), 5.11 (t), 2.01 (q); HRMS (ESI⁺) m/z 467.2260 (M+Na⁺, C₂₁H₄₀O₆NaSi₂ requires 467.2261). Data of **4b**: TLC (EtOAc) R_f = 0.83; ¹H-NMR (CDCl₃, 400MHz) δ 5.03 (1H, dd, J = 5.80, 3.80 Hz), 4.16 - 4.11 (3H, m), 3.39 (3H, s), 2.41 (1H, dd, J = 13.40, 5.80 Hz), 2.23 - 2.20 (2H, m), 2.14 - 2.04 (2H, m), 1.89 (1H, d, J = 13.80 Hz), 1.27 (3H, t, J = 11.23 Hz), 0.96 (9H, t, J = 8.04 Hz), 0.66 (6H, q, J = 7.89 Hz), 0.12 (9H, s); ¹³C-NMR (CDCl₃, 100MHz) δ 165.32 (s), 105.50 (d), 88.18 (d), 86.63 (s), 61.62 (t), 55.74 (q), 48.48 (t), 41.86 (t), 31.33 (d), 29.56 (d), 14.63 (q), 7.15 (q), 5.76 (t), 2.23 (q); HRMS (ESI⁺) m/z 467.3166, (M+Na⁺, C₂₁H₄₀O₆NaSi₂ requires 467.2261). Data of **4c**: TLC (EtOAc) R_f = 0.71; ¹H-NMR (CDCl₃, 400MHz) δ 5.17 (1H, dd, J = 5.38, 0.83 Hz), 4.83 (1H, s), 4.12 (2H, dq, J = 7.21, 1.85 Hz), 3.36 (3H, s), 2.42 - 2.35 (2H, m), 2.08 (1H, dd, J = 14.27, 0.87 Hz), 1.90 (1H, d, J = 13.3 Hz), 1.84 (1H, dd, J = 14.15, 4.82 Hz), 1.79 (1H, d, J = 3.99 Hz), 1.27 (3H, t, J = 7.11 Hz), 0.98 (9H, t, J = 7.94 Hz), 0.73-0.67 (6H, m), 0.15 (9H, s); ¹³C-NMR (CDCl₃, 100MHz) δ 107.56 (d), 98.19 (d), 60.51 (t), 54.88 (q), 47.19 (t), 43.80 (t), 35.01 (d), 30.75 (d), 14.31 (q), 7.12 (q), 6.74 (t), 1.83 (q); HRMS (ESI⁺) m/z 467.2274 (M+Na⁺, C₂₁H₄₀O₆NaSi₂ requires 467.2261).

(3a-Hydroxy-2-methoxy-6-oxohexahydrocyclopenta[b]furan-5-yl)acetic acid ethyl ester (5a,b)

HF-py (0.01 mL 70% HF, ca 1 mmol, 10 eq) was added at 0 °C to a solution of **4a** (39 mg, 0.09 mmol) in py (0.5 mL). After stirring for 1h at rt the reaction was quenched by the addition of silica gel (ca 0.5 g), followed by filtration over *Celite* after 15 min. Evaporation of the solvents and column chromatography (EtOAc) yielded an inseparable 2:1 mixture of isomers **5a,b** (18.4 mg, 79%) as a colorless oil; TLC (hexane/EtOAc 3:1) R_f = 0.20; ¹H-NMR (CDCl₃, 400MHz) δ 5.10-5.09 (0.4H, d, J = 3.60 Hz), 5.08 (0.6H, d, J = 3.48 Hz), 4.50 (0.4H, d, J = 1.16 Hz), 4.22 (0.5H, s), 4.09 - 4.05 (2H, m), 3.35 (3H, s), 3.35 - 3.00 (1H, m), 2.70 - 2.52 (3H, m), 2.46 - 2.31 (1H, dd, J = 13.56, 5.96 Hz); 2.18 (0.6H, d, J = 11.08 Hz), 2.14 (0.4H, d, J = 11.08), 2.04 - 1.97 (1H, m), 1.91 - 1.85 (1H, m), 1.28 - 1.23 (3H, m), ¹³C-NMR (CDCl₃, 100MHz) δ 171.56 (s), 107.40 (d), 107.00 (d), 89.58 (d), 88.50 (d), 82.43 (s), 61.05 (t), 60.82 (t), 55.24

(q), 55.19 (q), 46.66 (t), 45.29 (t), 44.02 (d), 43.53 (d) 34.04 (t), 34.01 (t), 14.14 (q); HRMS (ESI⁺) *m/z* 281.0993, (M+Na⁺, C₁₂H₁₈O₆Na requires 281.1001).

3a-6-Dihydroxy-2-methoxyhexahydrocyclopenta[*b*]furan-5-yl)acetic acid ethyl ester **6** and **7**

A solution of CeCl₃ (638 mg) and **5a,b** (133 mg, 0.52 mmol ca 1:1 mixture) in MeOH was stirred 15 min at rt. Then NaBH₄ (85 mg) was added at 0 °C and the reaction was quenched after 5 min with sat. NaHCO₃ (10 mL). After washing with sat. NaHCO₃ (2 x10 mL) and extraction with tBuOMe (3 x25 mL) and EtOAc (30 mL), the combined organic phases was dried over MgSO₄, filtered and the solvents evaporated. Column chromatography (hexane/EtOAc 1:4) yielded **6** (35 mg, 0.13 mmol, 26%), and **7** (41 mg, 0.16 mmol, 31%) both as colorless oils. Data of **6**: TLC (EtOAc) *R_f* = 0.64; ¹H-NMR (CDCl₃, 400MHz) δ 5.26 (1H, d, *J* = 4.28 Hz), 4.30 (1H, d, *J* = 4.28 Hz), 4.15 - 4.09 (3H, m), 3.39 (3H, s), 2.79 - 2.73 (1H, m), 2.62 (1H, dd, *J* = 15.76, 7.68 Hz), 2.38 (1H, dd, *J* = 15.76, 6.96 Hz), 2.16 (1H, d, *J* = 13.60 Hz), 2.03 (1H, dd, *J* = 13.44, 7.60 Hz), 1.97 (1H, dd, *J* = 13.68, 4.40 Hz), 1.77 (1H, t, *J* = 12.32 Hz), 1.26 (3H, t, *J* = 7.17); ¹H-NMR-difference-NOE (CDCl₃, 400MHz) δ 5.25 → 3.40 (9.04%), 1.97 (9.32%); 4.34 → 4.09 (6.62%), 2.27 (9.11%); 2.76 → 5.25 (0.63%), 4.34 (5.22%), 4.10 (6.49%), 2.38 (2.34%), 2.15 (1.29%), 2.02 (2.71%); 2.62 → 5.25 (0.93%), 4.33 (2.62%), 4.10 (4.77%), 2.37 (5.27%), 1.76 (1.64%); 2.37 → 5.25 (0.71%), 4.33 (1.85%), 4.10 (1.74%), 2.62 (7.29%); 2.17 → 5.25 (1.69%), 4.33 (2.62%), 4.10 (1.65%), 1.96 (14.68%); 2.05-1.94 → 5.25 (4.11%), 4.33 (2.27%), 4.10 (1.86%), 2.15 (10.52%), 1.77 (9.70%); 1.77 → 4.33 (1.82%), 4.10 (1.14%), 2.60 (2.52%), 2.38 (2.22%), 2.03 (12.36%), 1.99 (1.59%); ¹³C-NMR (CDCl₃, 100MHz) δ 172.93 (s), 109.69 (d), 94.01 (d), 87.48 (s), 72.71 (d), 60.67 (t), 55.15 (q), 48.31 (t), 41.74 (t), 41.23 (d), 34.08 (t), 14.50 (q); HRMS (ESI⁺) *m/z* 283.1152, (M+Na⁺, C₁₂H₂₀O₆Na requires 283.1157). Data of **7**: TLC (EtOAc) *R_f* = 0.45; ¹H-NMR (CDCl₃, 400MHz) δ 5.12 (1H, d, *J* = 4.25 Hz), 4.17-4.11 (3H, m), 3.72-3.67 (1H, m), 3.38 (3H, s), 2.65 (1H, dd, *J* = 15.64, 5.11 Hz), 2.33 (1H, dd, *J* = 15.66, 8.09 Hz), 2.18 (1H, d, *J* = 13.93 Hz), 2.09-2.00 (3H, m), 1.57 (1H, t, *J* = 5.35 Hz), 1.26 (3H, t, *J* = 5.37 Hz); ¹H-NMR-NOE (CDCl₃, 400MHz) δ 5.14 → 3.40 (5.31%), 2.04 (3.09%); 3.72 → 4.17 (7.48%), 2.69 (12.11%), 2.33 (3.01%) 1.39 (2.24%); 2.66 → 4.16 (2.58%), 3.71 (1.52%), 2.32 (12.56%), 2.19 (1.55%), 2.09 (2.41%), 1.38 (0.90%); 2.32 → 4.15 (1.09%), 3.71 (2.58%), 2.66 (1.52%), 2.06 (1.65%), 1.38 (2.02 %); 2.19 → 5.12 (0.67%), 2.06 (5.62%); 2.06 → 5.25 (0.60%), 2.67 (1.58%), 2.19 (1.67%), 1.38 (9.16%); ¹³C-NMR (CDCl₃, 100MHz) δ 173.20 (s), 108.32 (d), 87.49 (d), 84.97 (s), 75.94 (d), 60.91 (t), 54.99 (q), 48.91 (t), 41.38 (t), 39.58 (d), 37.02 (t), 14.51 (q); HRMS (ESI⁺) *m/z* 283.1151, (M+Na⁺, C₁₂H₂₀O₆Na requires 283.1157).

(3a,6-Bistrimethylsilyloxy-4,5,6a-tetrahydro-3aH-cyclopenta[*b*]furan-5-yl)acetic acid ethyl ester (**8**)

Lutidine (0.1 mL, 0.6 mmol, 5 eq) was added at 0 °C to a solution of **6** (29 mg, 0.11 mmol) in CH₂Cl₂

(0.4 mL), followed by TMS-OTf (0.1 mL, 0.72 mmol, 4 eq) after 5 min. After 15 min at 0 °C and 1 h at rt, the mixture was diluted with EtOAc (5 mL), washed with sat. NaHCO₃ (2 x 10 mL), extracted with EtOAc (3 x 10 mL) and evaporated to yield crude **8** (50 mg, >100%) as a brown oil, which was used for the next reaction without further purification; TLC (hexane/EtOAc 7:2) R_f = 0.94; ¹H-NMR (CDCl₃, 400 MHz): 6.34 (1H, d, J = 2.72 Hz), 5.12 (1H, d, J = 2.66 Hz), 4.41 (1H, d, J = 4.90 Hz), 4.34 (1H, dd, J = 4.80, 2.92 Hz), 4.14 (2H, q, J = 14.28 Hz), 2.50 - 2.47 (1H, m), 2.45 (1H, m), 2.30 - 2.25 (1H, m), 2.02 (1H, dd, J = 12.63, 6.26), 1.81 (1H, t, J = 12.74 Hz), 1.26 (3H, t, J = 7.09 Hz), 0.09 (9H, s), 0.07 (9H, s); ¹³C-NMR (CDCl₃, 100 MHz) δ 173.03 (s), 148.26 (d), 108.05 (d), 93.99 (d), 90.65 (s), 75.36 (d), 60.21 (t), 44.61 (t), 38.20 (d), 33.24 (t), 14.27 (q), 1.86 (q), 0.13 (q); HRMS (ESI⁺) m/z 395.1686, (M+Na⁺, C₁₇H₃₂O₅NaSi₂ requires 395.1686.

[3a-6-Dihydroxy-2-(5-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)hexahydrocyclopenta[b]-furan-5-yl]acetic acid ethyl ester (9)

A suspension of BSA (0.14 mL, 0.48 mmol, 4.2 eq) and thymine (61 mg, 0.48 mmol, 4.2 eq) in CH₂Cl₂ (0.2 mL) was stirred for 1 h at rt. Then a solution of crude **8** (50 mg, ca. 0.11 mmol) in CH₂Cl₂ (0.2 mL) was added and stirring was continued for 30 min. N-Iodosuccinimide (41 mg, 0.18 mmol, 1.5 eq) was added and the mixture stirred for 30 min at 0 °C and for 3 h at rt. The reaction mixture was then diluted with EtOAc (5 mL), washed with sat. Na₂CO₃ (2 x 5 mL) and sat. NaHCO₃ (5 mL) and extracted with EtOAc (3 x 10 mL) to yield the crude idonucleoside (54 mg, 0.09 mmol, 79%). This intermediate was dissolved in toluene (0.5 mL) and AIBN (10 mg, 0.06 mmol, 0.5 eq) and Bu₃SnH (ca 0.05 mL, 0.16 mmol, 1.5 eq) was added. After 2 h at reflux the solvents were removed yielding the crude deiodinated nucleoside (20.7 mg) that was subsequently redissolved in pyridine (1 mL) and treated with a solution of HF-pyridine (ca 0.01 mL 70% HF, 0.2 mmol, 5 eq) for 4 h at rt. The reaction mixture was quenched by the addition of silica gel (ca 0.5 g) filtered over *Celite* and washed with EtOAc/MeOH 10:1 (5 mL). Evaporation and column chromatography (CH₂Cl₂/MeOH 10:1) yielded **9** (7 mg, 0.014 mmol, 13% from **8**) as a colorless foam; TLC (EtOAc) R_f = 0.27; ¹H-NMR (CDCl₃, 400 MHz): 7.65 (1H, d, J = 1.24 Hz), 6.13 (1H, dd, J = 9.80, 5.28 Hz), 4.13 - 4.04 (2H, m), 3.94 (1H, d, J = 5.96 Hz), 3.64 (1H, dd, J = 9.80, 5.96 Hz), 2.67 (1H, d, J = 11.12 Hz), 2.40 (1H, dd, J = 13.72, 5.12 Hz), 2.31 - 2.24 (2H, m), 2.14 (1H, dd, J = 12.84, 5.60 Hz), 1.97 (1H, dd, J = 13.80, 9.80 Hz), 1.86 (3H, d, J = 1.20), 1.32 (1H, t, J = 12.20 Hz), 1.21 (3H, dd, J = 8.80, 7.08 Hz); ¹H-NMR difference-NOE (CD₃OD, 400 MHz) δ 7.64 → 6.13 (3.4%), 2.29 (3.6%), 1.97 (4.6%), 1.86 (7.0%); 6.13 → 7.65 (3.1%), 3.94 (4.9%), 2.40 (5.7%), 1.86 (1.8%) 1.21 (1.8%); 3.94 → 6.13 (5.9%), 3.65 (10.3%), 1.86 (1.3%), 1.21 (1.4%); 3.64 → 3.94 (11.0%), 2.27 (4.6%), 1.31 (3.1%); 2.67 → 3.69 (2.5%), 2.26 (35.9%); 2.40 → 6.13 (11.7%), 1.97 (23.4%); 2.26 → 7.64 (2.7%), 3.64 (3.6%), 2.67 (14.7%), 1.96 (2.8%); 2.13 → 1.96 (2.3%), 1.31 (22.6%); 1.96 → 7.64 (7.6%), 6.13 (2.4%), 2.39 (20.0%), 2.27 (5.7%), 2.13 (3.9%), 1.21 (1.4%); 1.86 → 2.4 (7.6%); ¹³C-NMR (MeOD, 100

MHz) δ 136.71 (d), 110.78 (s), 88.04 (d), 84.53 (d), 83.47 (s), 75.44 (d), 60.67 (t), 46.91 (t), 41.52 (t), 40.68 (d), 36.40 (t), 13.52 (q), 11.38 (q); HRMS (ESI⁺) m/z 377.1323 (M+Na⁺, C₁₆H₂₂N₂O₇Na requires 377.1324).

{6-[Bis-(4-methoxyphenyl)phenylmethoxy]-3a-hydroxy-2-(5-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)hexahydrocyclopenta[b]furan-5-yl}acetic acid ethyl ester (10)

DMT-OTf (250 mg, 0.53 mmol, 3.8 eq) was added in two portions at rt to a solution of **9** (50 mg, 0.14 mmol) in pyridine (0.3 mL). After 13h at rt, the mixture was diluted with EtOAc (5 mL), washed with NaHCO₃ (2 x 10 mL) and extracted with EtOAc (3 x 15 mL), the combined organic layers were dried over MgSO₄, filtered and the solvents. Column chromatography (EtOAc + 1% TEA) yielded **10** (66 mg, 72%) as a yellow foam; TLC (EtOAc) R_f = 0.45; ¹H-NMR (CDCl₃, 400MHz) δ 8.25 (1H, s), 7.67 (1H, d, J = 1.20 Hz), 7.52 - 7.36 (10H, m), 6.82 (4H, m), 6.20 (1H, dd, J = 9.12, 5.32 Hz), 4.03 (2H, dq, J = 14.44, 0.88 Hz), 3.80 (7H, m), 3.62 (1H, d, J = 5.48 Hz), 2.58 (1H, dd, J = 13.68, 5.20 Hz), 2.20 - 2.04 (4H, m), 1.96 - 1.89 (2H, m), 1.87 (3H, d, J = 1.20 Hz), 1.54 (1H, dd, J = 15.96, 10.32 Hz), 1.26 (1H, m), 1.20 (1H, t, J = 7.12 Hz); ¹³C-NMR (CDCl₃, 100 MHz) δ 172.94 (s), 163.65 (s), 159.22(s), 159.16 (s), 150.33 (s), 145.60 (s), 136.59 (s), 136.52 (s), 135.45 (d), 131.09 (d), 131.05 (s), 128.96 (d), 128.07 (d), 127.46 (d), 113.50 (s), 113.41 (d), 113.40 (d), 113.28 (s), 111.52 (s), 88.42 (d), 87.77 (s), 84.93 (d), 84.83 (s), 76.87 (d), 60.70 (t), 55.59 (q), 55.57 (q), 48.49 (t), 41.20 (d), 40.99 (t), 35.34 (t), 14.47 (q), 12.78 (q); HRMS (ESI⁺) m/z 679.2613 (M+Na⁺, C₃₇H₄₀N₂O₉Na requires 679.2632).

{6-[Bis-(4-methoxyphenyl)phenylmethoxy]-3a-[2-(cyanoethoxy)diisopropylaminophosphanyloxy]-2-(5-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)hexahydrocyclopenta[b]furan-5-yl}acetic acid ethyl ester (11)

To a solution of **10** (63 mg, 0.09 mmol) and diisopropylamine (0.07 mL, 0.4 mmol, 4.5 eq) in MeCN (0.3 mL) was added (iPr₂N)P(Cl)OCH₂CH₂CN (0.06 mL, 0.35 mmol, 4 eq) at rt. After stirring for 90min at rt, the mixture was diluted with EtOAc (5 mL), washed with sat NaHCO₃ (2 x 5 mL) and extracted with EtOAc (3 x 10 mL). The combined organic phases were dried over MgSO₄, evaporated and the crude product purified by chromatography (hexane/EtOAc 1:1) to give **11** (46 mg, 60%) as a colorless foam; TLC (EtOAc) R_f = 0.83, 0.74; ¹H-NMR (CDCl₃, 400MHz) δ 7.90 (1H, s), 7.66/7.65 (1H, 2d, J = 1.24/1.20), 7.51 - 7.48 (1H, m), 7.43 - 7.39 (4H, m), 6.84 - 6.80 (4H, m), 6.21 - 6.13 (1H,), 4.05 - 4.02 (2H, m), 3.79/3.78 (6H, 2s), 3.79 - 3.75 (1H, m), 3.73 - 3.61 (2H, m), 3.59 - 3.53 (3H, m), 3.00 - 2.91 (1H, m), 2.60-2.52 (2H, m), 2.18 - 2.04 (3H, m), 1.93 - 1.81 (3H, m), 1.31 - 1.08 (18H, m); ³¹P-NMR (CDCl₃, 100 MHz) δ 142.25, 141.80; HRMS (ESI⁺) m/z 879.3692 (M+Na⁺, C₄₆H₅₇N₄O₁₀NaP requires 879.3710).

ACKNOWLEDGEMENTS

Financial support by the Swiss National Science Foundation (grant-No: 200020-115913) is gratefully acknowledged.

REFERENCES

1. A. A. Koshkin, P. Nielsen, M. Meldgaard, V. K. Rajwanshi, S. K. Singh, and J. Wengel, *J. Am. Chem. Soc.*, 1998, **120**, 13252.
2. S. Obika, D. Nanbu, Y. Hari, J. I. Andoh, K. I. Morio, T. Doi, and T. Imanishi, *Tetrahedron Lett.*, 1998, **39**, 5401.
3. C. Hendrix, H. Rosemeyer, I. Verheggen, F. Seela, A. Van Aerschot, and P. Herdewijn, *Chem. Eur. J.*, 1997, **3**, 110.
4. R. Steffens and C. J. Leumann, *J. Am. Chem. Soc.*, 1997, **119**, 11548.
5. D. Renneberg and C. J. Leumann, *J. Am. Chem. Soc.*, 2002, **124**, 5993.
6. C. F. Bennett and E. E. Swayze, *Annu. Rev. Pharmacol. Toxicol.*, 2010, **50**, 259.
7. K. Tiemann and J. J. Rossi, *EMBO Molecular Medicine*, 2009, **1**, 142.
8. M. Tarköy, M. Bolli, and C. Leumann, *Helv. Chim. Acta*, 1994, **77**, 716.
9. M. Bolli, H. U. Trafelet, and C. Leumann, *Nucleic Acids Res.*, 1996, **24**, 4660.
10. M. Egli, P. Lubini, M. Bolli, M. Dobler, and C. Leumann, *J. Am. Chem. Soc.*, 1993, **115**, 5855.
11. J. C. Litten and C. Leumann, *Helv. Chim. Acta*, 1996, **79**, 1129.
12. S. Luisier and C. J. Leumann, *ChemBioChem*, 2008, **9**, 2244.
13. J. M. Conia, *Pure Appl. Chem.*, 1975, **43**, 317.
14. R. Steffens and C. Leumann, *Helv. Chim. Acta*, 1997, **80**, 2426.
15. E. J. Corey and A. G. Myers, *Tetrahedron Lett.*, 1984, **25**, 3559.
16. S. Luisier, P. Silhar, and C. J. Leumann, *Nucleic Acids Symp. Ser.*, 2008, 581.
17. M. Tarköy, M. Bolli, B. Schweizer, and C. Leumann, *Helv. Chim. Acta*, 1993, **76**, 481.