

## SYNTHESIS AND CHARACTERIZATION OF 6-SUBSTITUTED DIBENZO [d,f][1,3,2] DIOXA PHOSPHEPIN 6-OXIDES

K. Ananda Kumar, C. Suresh Reddy\*, L. Nagaprasada Rao and C. Naga Raju  
Department of Chemistry, Sri Venkateswara University, Tirupati - 517 502, India



**Abstract :** Several 6-trichloromethyl dibenzo[d,f] [1,3,2] dioxaphosphepin 6-oxide (3a), 2-chloroethoxy dibenzo[d,f] [1,3,2] dioxaphosphepin 6-oxide (3b) and alkylcarbamato dibenzo [d,f] [1,3,2] dioxaphosphepin 6-oxides (6a-c) have been synthesized from reactions of equimolar quantities of 2,2'-dihydroxybiphenyl (1) with trichloromethylphosphonic dichloride (2a), O-2-chloroethyl phosphoryldichloride (2b) and dichlorophosphinyl carbamates (5a-c) at 45-50°C for (3a & 3b) and 40-45°C for (6a-c) in dry toluene in the presence of triethylamine. Elemental, IR and NMR (<sup>1</sup>H & <sup>31</sup>P) data have been discussed and supported all structures.

### Introduction

A variety of organophosphorus compounds containing 1,3,2-dioxaphospholene and 1,3,2-dioxaphosphorinane have been used in cancer chemotherapy<sup>1,2</sup> and those of dibenzo[d,f] [1,3,2] dioxaphosphepins as stabilizers in polymer field<sup>3</sup>. Certain phosphorus carbamates possess antitumour, anticarcinogenic, pesticidal and bactericidal properties<sup>2,4-8</sup>. In view of their diverse applications, we have synthesized novel 6-trichloromethyl / chloroethoxy / alkylcarbamato dibenzo [d,f] [1,3,2] dioxaphosphepin 6-oxides and characterized them by elemental, IR and NMR (<sup>1</sup>H & <sup>31</sup>P) spectral analyses.

### Results and Discussion

When equimolar quantities of 2,2'-dihydroxybiphenyl (1) were condensed with trichloromethyl phosphonic dichloride (2a), O-2-chloroethyl phosphoryldichloride (2b) and dichloroisocyanatophosphinyl carbamates (5a-c), members of 3 and 6 were obtained (Scheme 1 & 2). The dichloroisocyanatophosphinyl carbamates (5a-c) have been obtained by the nucleophilic addition reaction of various dry alcohols to isocyanatophosphonic dichloride (4) at -10°C in dry toluene (scheme 2). Two equivalents of triethylamine as the base in dry toluene as the solvent and with



**SCHEME 2**

temperature ranging between 45-50°C for (3a & 3b) and 40-45°C for (6a-c) was found to be ideal to effect this cyclocondensations. Physical and spectral data of the compounds are found in tables 1-2.

The IR spectra<sup>9</sup> of 3a and 3b showed bands (in cm<sup>-1</sup>) at 1302 and 1297 respectively for (P=O) and 3b also exhibited band at 1030 for P-O-(C<sub>aromatic</sub>). Compounds 6a-c showed bands in the region 1247-1284 (P=O), 1743-1746 (C=O) and 3214-3271 (P-N-H) (Table 1). Proton NMR spectra<sup>10</sup> were simple for these systems.

The protons of aromatic moieties showed multiplets in the range of  $\delta$  7.19-7.52 for 3a and  $\delta$  7.18-7.65 for 3b, while on the chloroethyl group in 3b appeared as triplets at  $\delta$  4.39 and  $\delta$  3.66 for OCH<sub>2</sub> and CH<sub>2</sub>Cl groups respectively. Similarly the aromatic protons in 6a-c appeared as complex multiplets in the range of  $\delta$  7.22-7.61 (Table 2). The P-NH proton of the carbamate moiety in compounds 6a-c resonated as doublets in the region  $\delta$  6.03-6.67 (<sup>2</sup>J<sub>P-NH</sub>=14.8 Hz). It is interesting to note that the coupling is limited to P-NH only and not extended to the other protons of the carbamate moiety. It is also observed that the signals for the protons of the carbamate function appeared in slightly down field when compared to the signals of the corresponding protons in the free alcohols<sup>11</sup>.

Phosphorus resonance signals in 3a were observed at  $\delta$  8.32 and compounds 6a-c exhibited signals in two regions at  $\delta$  -0.37 to -4.19 and  $\delta$  -0.55 to 12.75. The two signals in <sup>31</sup>P NMR spectrum may be due to the presence of two conformers<sup>12</sup>.

#### Experimental

All melting points were determined in open capillary tubes on a Mel-Temp apparatus and are uncorrected. IR spectra ( $\delta_{\text{max}}$  in cm<sup>-1</sup>) were recorded as KBr pellets using a Perkin-Elmer 1000 unit. The <sup>1</sup>H and <sup>31</sup>P NMR spectra were recorded in CDCl<sub>3</sub> using varian AMX-400 MHz instrument and chemical shifts were referenced to TMS (<sup>1</sup>H) and 85% H<sub>3</sub>PO<sub>4</sub> (<sup>31</sup>P). Microanalytical data were obtained from Central Drug Research Institute, Lucknow, India.

#### Synthesis of 6-trichloromethyl dibenzo [d,f] [1,3,2] dioxaphosphepin 6-oxide (3a)

The general procedure to obtain the compounds 3a and 3b is illustrated for the preparation of 3a. A solution of trichloromethyl phosphonicdichloridate (2a, 2.36 g, 0.01 mol) in dry toluene (25 mL) was added dropwise to a stirred solution of 2,2'-dihydroxybiphenyl (1, 1.86 g, 0.01 mol) and triethylamine (2.02 g, 0.02 mol) in dry toluene (60 mL) at 0°C. After the addition was complete, the temperature was slowly raised to 45-50°C and was then maintained for 5-6 hours with stirring. The completion of the reaction was monitored by TLC analyses. After cooling to room temperature, the triethylamine hydrochloride was filtered off, and the filtrate was evaporated under reduced pressure. The gummy residue was washed with water, dried and treated with 2-propanol. A white solid formed and was recrystallised from 2-propanol to afford 3a, yield 0.168 gr(48%), mp 134-136°C. Physical and spectral data for compounds 3a and 3b are given in tables 1-2.

#### Synthesis of 6-ethylcarbamato dibenzo [d,f] [1,3,2] dioxaphosphepin 6-oxide (6b)

The general procedure for members of 6 is illustrated with that for 6b. A solution of ethanol (0.46 g, 0.01 mol) in dry toluene (20 mL) was added dropwise (20 min) to a cold (-10°C) solution of 4<sup>3</sup> (1.60 g, 0.01 mol) in dry toluene (20 mL). After the addition was complete, the reaction mixture was allowed to warm slowly to room temperature and stirring was continued for 2 hours. The new reaction mixture was then added dropwise to a cold (0°C)

**Table 1: Physical properties<sup>a</sup> and IR spectral(Cm<sup>-1</sup>) data of compounds (3a-b & 6a-c)**

Compd	Mol. Formula	M.P	Yield	Found (Calcd)%			IR(Cm <sup>-1</sup> )		
	(Mol.wt)	<sup>o</sup> C	%	C	H	C=O	P=O	P-NH	
3a	C <sub>11</sub> H <sub>8</sub> O <sub>3</sub> Cl <sub>1</sub> P (349.53)	134-136	48	44.51 (44.67)	2.27 (2.31)	-	1302	-	
3b	C <sub>14</sub> H <sub>12</sub> O <sub>3</sub> Cl <sub>1</sub> P (310.67)	70-73	90	54.26 (54.12)	3.79 (3.84)	-	1297	1030 (P-O- (C <sub>aliphatic</sub> ))	
6a	C <sub>14</sub> H <sub>12</sub> O <sub>3</sub> NP (305.23)	245-248	52	54.91 (55.09)	4.11 (3.96)	1745	1247	3249	
6b	C <sub>15</sub> H <sub>14</sub> O <sub>3</sub> NP (319.25)	250-252	60	56.52 (56.43)	4.39 (4.42)	1746	1248	3214	
6c	C <sub>16</sub> H <sub>16</sub> O <sub>3</sub> NP (333.28)	177-180	57	57.48 (57.66)	4.89 (4.84)	1743	1284	3271	

<sup>a</sup>Recrystallised from 2-Propanol

**Table 2: <sup>1</sup>H and <sup>31</sup>P NMR data <sup>a,b</sup> of compounds 3a-b and 6a-c**

Compd.	Ar-H	R-H	-NH	<sup>31</sup> P NMR <sup>c</sup>
3a	7.19 - 7.52 (m, 8H)	-	-	8.32
3b	7.18 - 7.65 (m, 8H)	4.39 (t, 2H, OCH <sub>2</sub> ) 3.66 (t, 2H, CH <sub>2</sub> Cl)	-	-
6a	7.27 - 7.58 (m, 8H)	3.64 (s, 3H, OCH <sub>3</sub> )	6.67 (brs)	-0.43 12.73
6b	7.28 - 7.61 (m, 8H)	1.15 (t, 3H, CH <sub>3</sub> ) 4.10 (q, 2H, OCH <sub>2</sub> )	6.22 (d, J=16Hz)	-0.37 12.76
6c	7.22-7.52 (m, 8H)	4.77 -4.85 (m, 1H, OCH) 1.06-1.08 (m, 6H, 2CH <sub>3</sub> )	6.03 (d, J=13.6Hz)	-4.19 -0.55

<sup>a</sup>Chemical shifts in  $\delta$  from TMS, J(Hz) given in parentheses

<sup>b</sup>Recorded in CDCl<sub>3</sub>

<sup>c</sup>Chemical shifts in  $\delta$  from 85% H<sub>3</sub>PO<sub>4</sub>

solution of 1 (1.86 g, 0.01 mol) and triethylamine (2.02 g, 0.02 mol) in dry toluene (50 mL). When the addition was complete, the mixture was stirred and allowed to warm slowly to 40-45°C and maintained at this temperature for 7 hours. The completion of the reaction was monitored by TLC analysis. The solid triethylamine hydrochloride was filtered off and the solvent was evaporated under reduced pressure. The gummy residue was washed with water, dried and treated with 2-propanol. A white solid obtained after recrystallization of the crude product from 2-propanol afforded 6b, yield 0.192 g (60%), mp 250-252°C. Physical and spectral data of 6a-c are given in tables 1-2.

#### Acknowledgements

The authors express their thanks to Professor C. Devendranath Reddy, Department of Chemistry, S.V. University, Tirupati for helpful discussions and also thankful to Directors SIF, IISc, Bangalore and CDRI, Lucknow for spectral recording.

#### References

1. L.A. Cates, *J. Med. Chem.*, **16**, 111(1973).
2. H. Arnold, F. Bourseaux and N. Brock, *Arzenimittel. Forsch.*, **11**, 143 (1961).
3. (a) S.D. Pastor, J.D. Spivack, P.L. Steinhuebel and C. Matzura, *Phosphorus, Sulfur and Silicon*, **15**, 253 (1983).  
(b) J.D. Spivack, *Brit Patent*, 1087399 (1982); *Chem. Abstr.*, **97**, 198374 (1982).
4. M. Kuranari, Chugai Pharmaceutical Co., *Jpn Patent* 26819 (1965); *Chem. Abstr.*, **64**, 9737 (1966).
5. (a) Z.B. Papanastassiou and T.J. Bardos, *J. Med. Chem.*, **5**, 1000 (1962);  
(b) R.I. Zhadanov, W.A. Buina, N. G. Kapitanova and I.A. Nuretdinov, *Synthesis*, 269 (1979).
6. O.M. Friedman, Z.B. Papanastassiou and R.S. Levi, *J. Med. Chem.*, **6**, 82 (1963).
7. H. Zimmer and A. Sill, *Progr. Drug. Res.*, **5**, 150 (1964).
8. P. Hsu, I.S. Kao, J.S. Tsai, C.H. Chou, M.C. Liu, M.L. Sheu, T.Y. Meug, Y.F. Jeu, R.C. Pan and S.W. Li, *Sci. Sinica (Perking)*, **13**, 789 (1964).
9. L.C. Thomas, *The Interpretation of the Infrared Spectra of Organophosphorus Compounds*, Heydon, London, 1974.
10. R.M. Silverstein, G.C. Bassler and T.C. Morrill, *Spectroscopic Identification of Organic Compounds*, Wiley Inc. New York, 1981.
11. Cymerman and Willis, *J. Chem. Soc.*, 1332 (1951).
12. D.G. Gorenstein, R. Rowell and J. Findlay, *J. Am. Chem. Soc.*, **102**, 5077 (1980).

Received on April 16, 2001