

Improved Synthesis of 4-(2-Substituted ethyl)-2-(4-Methylpiperazino)Quinazolines

Elizabeth Raux and Lucjan Streckowski*


Department of Chemistry, Georgia State University, Atlanta, Georgia 30302-4098, USA

ABSTRACT

An improved synthesis of the title compounds is reported.



Quinazoline derivatives show a wide variety of biological activities.¹⁻³ In particular, 2-amino substituted quinazolines exhibit activity as anti-inflammatory agents through the interaction with the histamine H₄ receptor,⁴ function as CCR4 antagonists,⁵ and are selective ligands for the 5-HT_{2A} receptor.^{6,7}

An improved procedure for the preparation of 4-(2-substituted ethyl)-2-(4-methylpiperazino)quinazolines **4-11** is the subject of this report.



* Email: Lucjan@gsu.edu

Route B

#	Nu	Yield (%)		#	Nu	Yield (%)	
		A	B			A	B
4	NMe ₂	27	50	8		23	50
5	HN-Bu ⁿ	31	55	9	OMe	27	52
6	NHCH ₂ Ph	50	62	10	SEt	35	54
7		40	53	11	SPh	50	63

The low-yield synthesis of quinazolines **4-11** (route A) was previously published.^{8,9} We now report an improved preparation of these products by simply reversing the order of the steps (route B). Thus, the first step of the more efficient synthesis involves nucleophilic displacement of chloride in 2-chloroquinazoline (**1**, 1.2 g, 7.2 mmol) with *N*-methylpiperazine (2.4 mL, 21.8 mmol) in toluene (10.0 mL) by heating the mixture at 80 °C for 4 h. The resulting compound **12** was purified by silica gel chromatography eluting with hexanes/ether (4:1). The subsequent treatment of **12** (0.77 mg, 3.4 mmol) with vinyl lithium¹⁰ (3.7 mmol in THF) followed by aromatization of the intermediate dihydroquinazoline (structure not shown) by potassium ferricyanide (2.6 g, 8.1 mmol) in saturated aqueous potassium hydroxide⁸ (1.5 mL) furnished compound **13**. Heating of **13** (0.16 g, 0.61 mmol) with a nucleophile (1.8 mmol) in toluene (5.0 mL) at 90 °C overnight gave the desired product **4-11** that, after purification (silica gel chromatography, hexanes/ether, 4:1) was identical in all respects with the corresponding products obtained by route A.⁸

REFERENCES

1. Y.A. Azev, S.V. Shorshnev and B.V. Golomolzin, *Tetrahedron Lett.*, **50**, 2899 (2009)
2. Y.Kabri, A. Gellis and P. Vanelle, *Green Chem.*, **11**, 201 (2009)
3. C.M. Dehnhardt, A.M. Venkatesan, Z. Chen, S. Ayral-Kaloustain, O. Dos Santos, E.D. Santos, K. Curran, M.T. Follettie, V. Diesl, J. Lucas, Y. Geng, S.Q. DeJoy, R. Petersen, I. Chaudhary, N. Brooijams, T.S. Mansour, K. Arndt and L. Chen, *J. Med. Chem.*, **53**, 897 (2010)
4. R. Smits, M. Adami, E.P. Istyastono, O.P. Zuiderveld, C.M.E. van Dam, F.J. de Kanter, A. Jongejan, G. Coruzzi, R. Leurs, I.J.P. de Esch, *J. Med. Chem.*, **53**, 2390 (2010)
5. K. Yokoyama, N. Ishikawa, S. Igarashi, N. Kawano, K. Hattori, T. Miyazaki, S. Ogino, Y. Matsumoto, M. Takeuchi, M. Ohta, *Bioorg. Med. Chem.*, **16**, 7021-2032 (2008)
6. J.L. Mokrosz, B. Duszyńska, S. Charakchieva-Minol, A.J. Bojarski, M.J. Mokrosz, R.L. Wydra, L. Janda, L. Strekowski, *Eur. J. Med. Chem.*, **31**, 973-980 (1996)
7. M. Avarado, M. Barceló, L. Carro, C.F. Masaguer, E. Raviña, *Chemistry & Biodiversity*, **3**, 106 (2006)
8. E. Raux, J. Klenc, A. Blake, J. Sączewski, L. Strekowski, *Molecules*, **15**, 1973-1984 (2010)
9. J. Sączewski, A. Paluchowka, J. Klenc, E. Raux, S. Barnes, S. Sullivan, B. Duszyńska, A. Bojarski, L. Strekowski, *J. Heterocycl. Chem.*, **46**, 1259-1265 (2009)
10. J.P. Quintard, B. Ellissondo, M. Pereyere, *J Organomet. Chem.*, **212**, C31 (1981)