

## High-Pressure Interaction of Sulfur Hexafluoride with Carbon Disulfide and Carbonyl Sulfide<sup>1</sup>

ARNULF P. HAGEN\* and BILL W. CALLAWAY

Received March 27, 1975


AIC502319

Sulfur hexafluoride has been found to react with carbon disulfide and carbonyl sulfide. At a minimum of 485° (1360 atm) with carbon disulfide the products are (CF<sub>3</sub>)<sub>2</sub>S, (CF<sub>3</sub>)<sub>2</sub>S<sub>2</sub>, carbon, and sulfur. Carbonyl sulfide reacts at 500° (270 atm) forming CF<sub>4</sub>, SOF<sub>2</sub>, and sulfur. No reaction takes place with carbon monoxide or carbon dioxide at conditions up to 500° (4000 atm). Graphite reacts at 500° (135 atm) forming CF<sub>4</sub> and SF<sub>4</sub> in a reaction which becomes nearly quantitative at 500° (4000 atm). When (CF<sub>3</sub>)<sub>2</sub>S<sub>2</sub> is combined with carbon disulfide at 540° (4000 atm) the products include carbon, sulfur, and fluoroalkanes. The observed interaction of CO<sub>2</sub>, COS, and CS<sub>2</sub> with SF<sub>6</sub> can be best explained by an initial reaction which results in the formation of SF<sub>4</sub>, COF<sub>2</sub> or CSF<sub>2</sub>, and sulfur.

Few reaction systems which include sulfur hexafluoride have been studied, even though many reactions are thermodynamically possible at standard conditions.<sup>2</sup> At moderate temperatures in a sealed glass ampoule it has been found to react with AlCl<sub>3</sub> and SO<sub>3</sub>. The former reaction at 200° formed sulfur chlorides and the latter at 250° led to a 20% conversion to form SO<sub>2</sub>F<sub>2</sub>.<sup>3</sup> In a previous study we have shown SF<sub>6</sub> to react with a series of oxides including MgO, NiO, SiO<sub>2</sub>, and water with a 10% conversion of SO<sub>2</sub>F<sub>2</sub> at 475° (130 atm) and 90% conversions at 500° (3300–4000 atm).<sup>4</sup>


### Results and Discussion

Sulfur hexafluoride has been found to react with carbon disulfide and carbonyl sulfide but not with CO<sub>2</sub> at conditions of temperature and pressure up to 500° (4000 atm). The results are summarized in Table I. The reaction



was observed at 485° (1350 atm) with a 19% conversion of SF<sub>6</sub>; when the pressure was increased to 4000 atm, a 73% conversion of SF<sub>6</sub> was obtained, and at 495° (4000 atm) 90% of the SF<sub>6</sub> was consumed. This reaction is analogous to the low-pressure reactions of CS<sub>2</sub> with IF<sub>5</sub> (195°),<sup>5</sup> HgF<sub>2</sub> (460°),<sup>6</sup> and UF<sub>6</sub> (25°),<sup>7</sup> and, in the presence of catalytic amounts of AsF<sub>3</sub> or BF<sub>3</sub>, SF<sub>4</sub> (200°).<sup>8</sup> The recovered (CF<sub>3</sub>)<sub>2</sub>S was formed from the thermal decomposition of (CF<sub>3</sub>)<sub>2</sub>S<sub>2</sub> which has been demonstrated to take place at low<sup>9</sup> and high pressures.

Since in this work CS<sub>2</sub> was found to undergo thermal decomposition at the temperatures and pressures where reaction took place, experiments were designed to determine whether the initial reaction of SF<sub>6</sub> was with carbon disulfide, carbon, or sulfur. Sulfur hexafluoride did not react with carbon at 450° (4000 atm), but at 500° (135 atm) a 20% reaction took place, and at 500° (4000 atm) the reaction



was nearly quantitative. At 500° (4000 atm) neither SF<sub>6</sub> nor CF<sub>4</sub> reacted with sulfur and CF<sub>4</sub> did not react with carbon or carbon disulfide. The lack of CF<sub>4</sub> in the reaction between SF<sub>6</sub> and carbon disulfide eliminated the formation of SF<sub>4</sub> by an initial reaction of SF<sub>6</sub> with carbon as the primary reaction.

Above 500° the reaction between carbon disulfide and SF<sub>6</sub> became more complex. In addition to thermal decomposition, the (CF<sub>3</sub>)<sub>2</sub>S<sub>2</sub> reacted with CS<sub>2</sub> or SF<sub>6</sub>. At 540° (270 atm) SF<sub>6</sub> and (CF<sub>3</sub>)<sub>2</sub>S<sub>2</sub> did not react, but when the pressure was increased to 4000 atm, CF<sub>4</sub> and SF<sub>4</sub> were formed. At 540° (270 atm) CS<sub>2</sub> and (CF<sub>3</sub>)<sub>2</sub>S<sub>2</sub> interacted forming CF<sub>4</sub>, SF<sub>4</sub>, and sulfur. When the pressure was increased to 4000 atm several fluoroalkanes up to C<sub>5</sub>F<sub>12</sub> were also isolated.<sup>9</sup>

No reaction was found between SF<sub>6</sub> and carbon dioxide at 500° (4000 atm). This lack of reaction is important in itself. It rules out the thermal decomposition of SF<sub>6</sub> to form SF<sub>4</sub> as



the primary reaction since CO<sub>2</sub> and SF<sub>4</sub> have been shown to form COF<sub>2</sub> and CF<sub>4</sub> at high temperatures.<sup>10</sup>

The reactions of SF<sub>6</sub> with carbonyl sulfide were observed at lower pressures than for the reaction of SF<sub>6</sub> with CS<sub>2</sub>. The results are summarized in Table II. No reaction took place at 450° (4000 atm) or at 500° (170 atm), but at 500° (270 atm) all of the SF<sub>6</sub> reacted according to the equation




The same stoichiometry was obtained whether the SF<sub>6</sub> or the COS was in excess.

Previous workers have shown that COS decomposed via two independent simultaneous equilibria




It was noticed that these reactions were sensitive to the vessel used for reaction.<sup>11</sup> For example more decomposition was noted when "Jena" glass vessels were used or silica was placed in a "Pyrex" vessel than was noted for a "clean Pyrex" reaction. In this laboratory no decomposition was noted at 500° and at 170, 270, or 4000 atm in gold tubing. Even though no thermal decomposition of the carbonyl sulfide had been observed in this study, SF<sub>6</sub> was combined with CO at 500° (4000 atm). No reaction took place in this experiment. Combining this experiment with the previously discussed reactions of CS<sub>2</sub> and the lack of reaction of SF<sub>6</sub> with sulfur and CO<sub>2</sub> indicated the primary reaction was with COS and not with its thermal decomposition products.


There are two reaction sequences which can describe the observed stoichiometry. The first sequence is analogous to the reaction of SF<sub>6</sub> with other oxides which lead to the formation of SO<sub>2</sub>F<sub>2</sub><sup>4</sup>



The SO<sub>2</sub>F<sub>2</sub> and CS<sub>2</sub> then react to form SOF<sub>2</sub> and sulfur



In a series of experiments with SO<sub>2</sub>F<sub>2</sub> the products of the second reaction were found not to include SOF<sub>2</sub> but did include (CF<sub>3</sub>)<sub>2</sub>S<sub>x</sub> (where x = 1, 2, 3) at 500° (4000 atm)



It was not possible to adjust the SO<sub>2</sub>F<sub>2</sub>:CS<sub>2</sub> ratio to eliminate the formation of (CF<sub>3</sub>)<sub>2</sub>S<sub>2</sub>. Since this material was not observed in any of the SF<sub>6</sub> experiments it is not unreasonable to rule out the initial formation of SO<sub>2</sub>F<sub>2</sub>.

The second reaction sequence has as its first step the formation of COF<sub>2</sub> and SF<sub>4</sub>




Table I. SF<sub>6</sub> and CS<sub>2</sub> System

Pressure, atm	Temp, °C	Time, hr	Conversion, <sup>a</sup> %	Amt of reactants, mmol		Amt of material out, mmol							
				SF <sub>6</sub>	CS <sub>2</sub>	SF <sub>6</sub>	CS <sub>2</sub>	(CF <sub>3</sub> ) <sub>2</sub> S <sub>2</sub>	(CF <sub>3</sub> ) <sub>2</sub> S	CF <sub>4</sub>	SF <sub>4</sub>	S <sup>b</sup>	
4000	280	18		0.65	0.68	0.65	0.68						
4000	430	18		1.26	1.09	1.26	1.09						
335	485	24		0.87	0.72	0.87	0.72						
1000	485	24		0.79	0.75	0.79	0.75						
1350	485	24	19	0.75	0.52	0.69	0.41	0.02	0.04				0.21
4000	485	18	73	0.82	1.80	0.17	0.48	0.52	0.13				2.01
4000	495	18	90	0.67	1.53		0.15	0.22	0.45				2.46
170	500	24		0.87	0.74	0.87	0.74						
270	520	18	100 <sup>c</sup>	1.27	1.07		0.33			0.64	1.27		1.27
4000	540	24	100	0.97	0.91	0.06			0.30	0.30	0.61		1.81


<sup>a</sup> Percent of CS<sub>2</sub> consumed. <sup>b</sup> Calculated. <sup>c</sup> Percent of SF<sub>6</sub> consumed.

Table II. SF<sub>6</sub> and COS System


Pressure, atm	Temp, C	Time, hr	Conversion, %	Amt of reactants, mmol		Amt of material out, mmol						
				SF <sub>6</sub>	COS	SF <sub>6</sub>	COS	CF <sub>4</sub>	SOF <sub>2</sub>	S <sup>a</sup>		
4000	300	24		1.26	1.26	1.26	1.26					
335	450	20		0.91	0.90	0.91	0.90					
4000	450	18		0.91	0.90	0.91	0.90					
170	500	18		1.15	0.95	1.15	0.95					
270	500	18	100 <sup>b</sup>	1.17	1.26		0.90	1.17	1.17			1.17
4000	500	24	100 <sup>c</sup>	1.60	1.24	0.36		1.24	1.23			1.24

<sup>a</sup> Calculated. <sup>b</sup> Percent of SF<sub>6</sub> consumed. <sup>c</sup> Percent of COS consumed.

This reaction is then followed by




a reaction that has been shown to take place readily without the need for pressure.<sup>10</sup> In related experiments at 500° (4000 atm), CO and COF<sub>2</sub> were found not to be fluorinated by SF<sub>6</sub>, and SF<sub>6</sub> was found not to react with sulfur. The reaction




was observed at 500° (4000 atm). In this reaction no CF<sub>4</sub> or SOF<sub>2</sub> was obtained which indicated that the SF<sub>6</sub> and carbonyl sulfide did not react. When COS, SF<sub>6</sub>, and COF<sub>2</sub> were combined at 500° (4000 atm), no CF<sub>4</sub> or SOF<sub>2</sub> were obtained and at the same conditions COS and COF<sub>2</sub> did not react.


The reactions of SF<sub>6</sub> with carbon disulfide and carbonyl sulfide are temperature and pressure dependent. These two interactions and the lack of reaction with carbon dioxide can be explained by the initial formation of SF<sub>4</sub> via a coordinated intermediate, SF<sub>6</sub>·CS<sub>2</sub>





This intermediate then decomposes forming sulfur tetrafluoride



Equations can be written for CS<sub>2</sub>, COS, and CO<sub>2</sub> which all involve the same type of coordinated intermediate



The different secondary reactions can then be explained by reactions of SF<sub>4</sub> with COF<sub>2</sub> or CS<sub>2</sub> and CSF<sub>2</sub>



The lack of any reaction between CO<sub>2</sub> and SF<sub>6</sub> is not surprising since the initial interaction would not be thermodynamically favorable as compared to the COS and CS<sub>2</sub> interactions.

## Experimental Section

All experiments were conducted under conditions at which the compounds were relatively thermally stable to minimize extraneous reactions. When an interaction was not observed at or near ambient conditions, the system was then examined at elevated conditions of temperature and pressure.

All work at reduced pressures was carried out in a borosilicate glass vacuum system constructed with Teflon stopcocks (Fischer & Porter Co., Warminster, Pa.). High pressures were generated using a gas pressure booster (High Pressure Equipment Co., Erie, Pa.) attached to a high pressure-high temperature hydrothermal research unit (Model HR-1B-4, Tem-Pres Research, State College, Pa.). The samples were contained in sealed ampoules made from 3-mm diameter thin-walled gold tubing which were placed in a high-pressure reactor. Nitrogen gas was used to generate the desired pressure. At the end of a reaction period the reactor was cooled to -196° before releasing the pressure. The ampoule was removed and placed into an opening device attached to the vacuum line. After warming of the system to room temperature, water, CO<sub>2</sub>, and other condensable materials on the surface of the gold tubing were pumped away. The opening device and vacuum line were pretreated with Me<sub>3</sub>SiCl or SF<sub>4</sub> to remove any remaining traces of H<sub>2</sub>O, and then the ampoule was opened. The substances which volatilized were transferred directly into the vacuum line. Solid material was recovered in a glove bag under an N<sub>2</sub> or Ar atmosphere.

All reaction mixtures were separated by using standard vacuum-line fractionation techniques, except for the removal of SF<sub>4</sub>, SOF<sub>2</sub>, COF<sub>2</sub>, or COS by condensation on powdered moist NaOH. All products were identified and confirmed by two or more techniques including infrared spectroscopy, mass spectroscopy, vapor pressure measurements, melting point determinations, or elemental analysis.

**Instrumentation.** Infrared absorption spectra were obtained in the 4,000–300 cm<sup>-1</sup> region using a Beckman Model IR-10 double-beam, grating spectrophotometer. Volatile materials were confined in a 100-mm gas cell fitted with KBr windows sealed with rubber O rings at reduced pressure. The instrument was calibrated using polystyrene.

The mass spectra were obtained using a Hitachi-Perkin Elmer RMU-7E mass spectrometer with an ionizing potential of 70 eV, current of 50 μA, ion-source temperature of 150°, accelerating potential of 3600 V, and a pressure between 10<sup>-6</sup> and 5 × 10<sup>-5</sup> Torr as measured by the unit's gauge.

**Reagents.** All reagents were commercial materials. The volatile substances were vacuum distilled just prior to use. Carbon dioxide and COS were purified by distilling the sample through a trap cooled to -95° (toluene slush) to remove impurities of low volatility and then retaining the material which stopped in a trap cooled to -145° (pentane-isopentane slush).<sup>12,13</sup> The same procedure was used to

purify  $(\text{CF}_3)_2\text{S}_2^{14}$  and  $\text{CS}_2^{15}$  using a  $-78^\circ$  bath (Dry Ice-acetone mixture) and a  $-95^\circ$  bath. Sulfur hexafluoride was purified by the same procedure using a  $-95^\circ$  and a  $-130^\circ$  (pentane slush) bath.<sup>16</sup> Carbon tetrafluoride was passed through a trap cooled to  $-130^\circ$  before use.<sup>17</sup> Sulfur was used as sublimed sulfur flowers. The oxygen-free carbon used in this study was "Sterling MT 2700° Graphitized Carbon" prepared by the Cabot Corp.

**Reaction of  $\text{SF}_6$  and  $\text{CS}_2$ .** Sulfur hexafluoride (120 mg, 0.823 mmol) and  $\text{CS}_2$  (136.8 mg, 1.800 mmol) were condensed into a gold tube at  $-196^\circ$ . The tube was sealed and held at  $485^\circ$  (4000 atm) for 18 hr. The tube was opened and the volatile material was separated by passing through traps cooled to  $-95^\circ$  and  $-130^\circ$ . The material that passed was collected at  $-196^\circ$ . The  $-95^\circ$  trap contained  $(\text{CF}_3)_2\text{S}_2^{14,18}$  and  $\text{CS}_2$  (142.6 mg, 1.006 mmol). The  $-130^\circ$  fraction contained  $(\text{CF}_3)_2\text{S}$  (22.4 mg, 0.132 mmol) and the  $-196^\circ$  trap held  $\text{SF}_6$  (24.1 mg, 0.165 mmol).

Additional experiments are summarized in Table I.

**Reaction of  $\text{SF}_6$  and  $\text{CS}_2$ .** Sulfur hexafluoride (142 mg, 0.979 mmol) and  $\text{CS}_2$  (98.8 mg, 1.30 mmol) were condensed into a gold tube at  $-196^\circ$ . The tube was sealed and held at  $540^\circ$  (4000 atm) for 24 hr. The tube was opened and the volatile material was condensed at  $-196^\circ$ . The mixture was passed through a trap cooled to  $-130^\circ$  into a trap at  $-196^\circ$ . The former trap contained a mixture of  $(\text{CF}_3)_2\text{S}_2^{14}$  and  $\text{SF}_4^{19,20}$  (115 mg, 0.830 mmol) and the  $-196^\circ$  trap contained  $\text{CF}_4$  (45 mg, 0.42 mmol). The gold tube contained S (84.3 mg, 2.63 mmol).

**Reaction of  $\text{SF}_6$  and COS.** Sulfur hexafluoride (233 mg, 1.60 mmol) and COS (74.4 mg, 1.24 mmol) were condensed into a gold tube at  $-196^\circ$ . The tube was sealed and held at  $500^\circ$  (4000 atm) for 24 hr. The gold ampoule was opened and the volatile material was condensed at  $-196^\circ$ . The volatile material was passed through a trap at  $-145^\circ$  into a trap at  $-196^\circ$ . The former trap contained a mixture of  $\text{SF}_6$  and  $\text{SOF}_2$  (159 mg, 1.59 mmol) and the latter  $\text{CF}_4$  (109 mg, 1.24 mmol). The gold tube contained S (39.7 mg, 1.24 mmol).

Additional experiments are summarized in Table II.

**Reaction of  $(\text{CF}_3)_2\text{S}_2$  and  $\text{SF}_6$ .** Bis(perfluoromethyl) disulfide (162 mg, 0.802 mmol) and  $\text{SF}_6$  (206 mg, 1.41 mmol) were condensed into a gold tube at  $-196^\circ$ . The tube was sealed and held at  $540^\circ$  (4000 atm) for 24 hr. The gold ampoule was opened and the volatile materials were condensed into a trap at  $-196^\circ$ . The volatile material was separated by passing through a series of cold traps cooled to  $-78^\circ$ ,  $-95^\circ$ ,  $-130^\circ$ ,  $-145^\circ$ , and  $-160^\circ$  into a trap cooled to  $-196^\circ$ . All of the volatile material passed the  $-78^\circ$  trap. The material which stopped in the  $-95^\circ$  trap was a mixture of  $\text{CS}_2$ ,  $(\text{CF}_3)_2\text{S}_2$ ,  $\text{C}_3\text{F}_{12}$ , and  $\text{C}_4\text{F}_{10}$ .<sup>21</sup> The  $-130^\circ$  and  $-145^\circ$  fractions were the same and were combined. This fraction contained  $\text{C}_2\text{F}_6$  and  $\text{C}_3\text{F}_8$ .<sup>21</sup> The  $-160^\circ$  trap contained nothing. The  $-196^\circ$  trap contained  $\text{CF}_4$  (0.20 mmol). The gold tube contained a mixture of carbon and sulfur.

**Reaction of  $\text{SF}_6$  with CO and S.** Sulfur hexafluoride (127 mg, 0.870 mmol) was condensed into a gold tube which had been charged with sulfur (100 mg, 3.13 mmol) at  $-196^\circ$ . Liquid CO was then condensed into the tube at  $-196^\circ$  and the tube sealed. The ampoule was held at  $500^\circ$  (4000 atm) for 24 hr and then opened. The material which condensed at  $-196^\circ$  was found to be a mixture of  $\text{SF}_6$ , COS, and  $\text{COF}_2$  (177 mg, 2.03 mmol). This mixture after treatment with NaOH yielded only  $\text{SF}_6$  (84 mg, 0.58 mmol).

**Reaction of  $\text{SF}_6$  and Carbon.** Sulfur hexafluoride (150 mg, 1.03 mmol) was added to a gold tube which had been charged with carbon (60.1 mg, 5.00 mmol). The tube was sealed and held at  $500^\circ$  (4000 atm) for 24 hr. The ampoule was opened and the volatile materials condensed at  $-196^\circ$ . The mixture was passed through a trap cooled

to  $-160^\circ$  (isopentane slush) into a trap cooled to  $-196^\circ$ . The latter trap contained  $\text{CF}_4$  (42 mg, 0.48 mmol). The former trap contained  $\text{SF}_4$  and a trace of  $\text{SF}_6$  (108 mg, 1.00 mmol).

No reaction was found at  $450^\circ$  (4000 atm) and a 20% conversion took place at  $500^\circ$  (135 atm).

**Decomposition of  $\text{CS}_2$ .** Carbon disulfide (210.0 mg, 2.763 mmol) was condensed into a gold tube at  $-196^\circ$ . The tube was sealed and held at  $500^\circ$  (4000 atm) for 24 hr. The tube was opened and the volatile material was condensed at  $-196^\circ$ . The trap contained only  $\text{CS}_2$  (125.9 mg, 1.656 mmol). The gold tube contained a mixture of carbon and sulfur. No decomposition was found at  $250^\circ$  (4000 atm) and 51% decomposition was observed at  $500^\circ$  (275 atm).

**Reaction of  $\text{SF}_2\text{O}_2$  and  $\text{CS}_2$ .** In a qualitative experiment sulfur fluoride (125.6 mg, 1.231 mmol) and  $\text{CS}_2$  (84.06 mg, 1.106 mmol) were condensed into a gold tube at  $-196^\circ$ . The tube was sealed and held at  $395^\circ$  (4300 atm) for 18 hr. The tube was opened and the volatile material was separated by passing the mixture through a trap cooled to  $-130^\circ$  into a trap at  $-196^\circ$ . The former trap contained unreacted  $\text{CS}_2$  and  $(\text{CF}_3)_2\text{S}_x$  (where  $x = 1, 2, 3$ ; confirmed by mass spectroscopy) and the  $-196^\circ$  trap contained  $\text{SO}_2\text{F}_2$  and COS.

Additional qualitative experiments at  $500^\circ$  (3000 and 4000 atm) gave the same results.

**Acknowledgment.** The authors wish to thank Dr. H. W. Beck for his assistance in obtaining mass spectral data. This research was supported by National Science Foundation Grant GP-19873.

**Registry No.**  $\text{SF}_6$ , 2551-62-4;  $\text{CS}_2$ , 75-15-0; COS, 463-58-1.

## References and Notes

- (1) This report is based on portions of a dissertation submitted by B. W. Callaway to The University of Oklahoma in partial fulfillment of the requirements for the degree of Doctor of Philosophy.
- (2) M. Kh. Karapet'yants and M. L. Karapet'yants, "Thermodynamic Constants of Inorganic and Organic Compounds", J. Schmorak, Translator, Ann Arbor-Humphrey Science Publishers, Ann Arbor, Mich., 1970.
- (3) J. R. Case and F. Nyman, *Nature (London)*, **193**, 473 (1962).
- (4) A. P. Hagen, D. J. Jones, and S. R. Ruttman, *J. Inorg. Nucl. Chem.*, **36**, 1217 (1974).
- (5) R. N. Haszeldine and J. M. Kidd, *J. Chem. Soc.*, 3219 (1953).
- (6) E. H. Man, D. D. Coffman, and E. L. Muettterties, *J. Am. Chem. Soc.*, **81**, 3575 (1959).
- (7) L. E. Trevorrow, J. Fischer, and W. H. Gunther, *Inorg. Chem.*, **2**, 1281 (1963).
- (8) R. J. Harder and W. C. Smith, *J. Am. Chem. Soc.*, **83**, 3422 (1961).
- (9) E. W. Lawless and L. D. Harman, *J. Inorg. Nucl. Chem.*, **31**, 1541 (1969).
- (10) W. R. Hasek, W. C. Smith, and V. A. Engelhardt, *J. Am. Chem. Soc.*, **82**, 543 (1960).
- (11) J. R. Partington and H. H. Neville, *J. Chem. Soc.*, 1230 (1951).
- (12) R. H. Pierson, A. N. Fletcher, and E. St. C. Gantz, *Anal. Chem.*, **28**, 1218 (1956).
- (13) H. J. Collomon, D. C. McKean, and H. W. Thompson, *Proc. R. Soc. London, Ser. A.*, **208**, 341 (1951).
- (14) G. R. A. Brandt, H. J. Emeleus, and R. N. Haszeldine, *J. Chem. Soc.*, 2549 (1952).
- (15) T. Wentink, Jr., *J. Chem. Phys.*, **29**, 188 (1958).
- (16) R. T. Lagemann and E. A. Jones, *J. Chem. Phys.*, **19**, 534 (1951).
- (17) J. Goubeau, W. Bues, and F. W. Kampmann, *Z. Anorg. Allg. Chem.*, **283**, 123 (1956).
- (18) H. A. Carter, C. S.-C. Wang, and J. M. Shreeve, *Spectrochim. Acta, Part A*, **29**, 1479 (1973).
- (19) J. K. O'Loane and M. K. Wilson, *J. Chem. Phys.*, **23**, 1313 (1955).
- (20) R. E. Dodd, L. A. Woodward, and H. L. Roberts, *Trans. Faraday Soc.*, **52**, 1052 (1956).
- (21) American Petroleum Institute Research Project 44, Spectra 196, 401, 444, 445, 729.