


Synthesis, Characterization, And Computational Study of Complexes Containing Pt···H Hydrogen Bonding Interactions

Miguel Baya, Úrsula Belío, and Antonio Martín*

Instituto de Síntesis Química y Catálisis Homogénea (ISQCH), Departamento de Química Inorgánica, Universidad de Zaragoza–CSIC, 50009 Zaragoza, Spain

Supporting Information

ABSTRACT: Complexes $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})\text{L}]$ (bzq = 7,8-benzoquinoline; L = 8-hydroxyquinoline, hqH (**1**); 2-methyl-8-hydroxyquinoline, hqH' (**2**)) have been prepared by replacing the labile acetone ligand in the starting material $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})(\text{Me}_2\text{CO})]$. The ^1H NMR spectra of **1** and **2** show that the signals attributable to the hydroxyl proton of the hqH or hqH' ligands are displaced downfield 2.64 ppm for **1** and 2.74 ppm for **2** with respect to the respective free ligands. Moreover, in both complexes the signals present platinum satellites with $J(\text{Pt},\text{H})$ coupling constant of 67.0 Hz for **1** and 80.6 Hz for **2**. All these features are indicative of the existence of $\text{Pt}\cdots\text{H}-\text{O}$ hydrogen bonds in solution for these complexes. The structures of complexes **1** and **2** have been established by an X-ray diffraction study and allow us to confirm the existence of these interactions in the solid state too. Thus, in both cases the hydroxyl hydrogen atom is pointing toward the metal center, and the measured geometric parameters involving this hydrogen are $\text{Pt}-\text{H} = 2.09(4)$ Å, $\text{O}-\text{H} = 0.94(4)$ Å, $\text{Pt}-\text{H}-\text{O} = 162(4)^\circ$, for **1**, and $\text{Pt}-\text{H} = 2.10(4)$ Å, $\text{O}-\text{H} = 0.91(4)$ Å, $\text{Pt}-\text{H}-\text{O} = 162(4)^\circ$, for **2**, all of which are fully compatible with a hydrogen bond system. Complexes **1** and **2** and the analogues $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{hqH})]^-$ (**A**) and $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{hqH}')^-]$ (**B**), prepared some time ago in our laboratory and also showing $\text{Pt}\cdots\text{H}-\text{O}$ hydrogen bonds, have been the object of theoretical calculations to obtain better insight into the $\text{Pt}\cdots\text{H}$ interactions. Their density functional theory (DFT) calculated structures show excellent agreement with the X-ray determined ones (**1**, **2**, and **B**). Topological analyses of the electron density function ($\rho(\mathbf{r})$) have been performed on the four complexes according to Bader's *Atoms In Molecules* theory. These analyses reveal a bond path that relates the platinum atom and the hydroxyl hydrogen atom, as well as the corresponding bond critical points. The values of the Laplacian $\nabla^2\rho(\mathbf{r})$ and local energy density $H(\mathbf{r})$ indicate that these are closed shell, electrostatic interactions, but with *partial covalence*. The deprotonation of the OH fragment in **1** and **2** with BuLi leads to the formation of the unexpected trinuclear complexes $(\text{NBu}_4)[\text{Li}\{\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})(\text{L})\}_2]$ ($\text{L} = \text{hq}$ (**3**), hq' (**4**)). The X-ray structures of these have shown a change in the coordination of the deprotonated hq and hq' , which are now bonded to the Pt atoms through their O atoms, and which are bridging the Pt and Li metal atoms.



INTRODUCTION

The existence of interactions between metal centers and hydrogen atoms attached to main group elements (especially C, N, and O) has been recognized since the early 1980s.^{1–3} The first of these types of interactions to be studied and understood were the “agostic” interactions.^{1,4} In these, the metal center acts as a Lewis acid, generally receiving electron density from a C–H bond and resulting in a 3-center–2-electron (3c–2e) bond system. The three centers are the metal and the C and H atoms, and the two electrons are those in the C–H bond. An empty orbital of the metal is involved to house the donated electron density. A characteristic of agostic interactions is the upfield displacement in the ^1H NMR spectra of the signal of the hydrogen involved.

In the second type of $\text{M}\cdots\text{H}$ interactions a transition metal atom is acting as a proton acceptor in a formally hydrogen bonding interaction. Although some early reports on IR studies

in solution⁵ mention the possibility of metal centers being involved in hydrogen bridging, these $\text{M}\cdots\text{H}-\text{X}$ systems have been recognized and understood from the early 1990s and now they are well established.^{2,3,6–10} They are substantially similar to “classic” hydrogen bonds, that is, the metal atom is the Lewis base that has a filled orbital with an electron pair that can interact with an electropositive hydrogen atom. Using a molecular orbital method language, the electron pair is donated to create a 3-center–4-electron (3c–4e) system. These hydrogen bonds therefore are favored by electron rich metals such as late transition metals, especially in low oxidation states. Pt(II) complexes have been found to be particularly suited to this interaction because of their planar nature and the electron pair housed in the $5d_{z^2}$ orbital, available to participate in the

Received: August 7, 2013


Published: December 12, 2013

Pt...H–X 3c–4e system of the hydrogen bond. The signal of the proton involved in the M...H–X hydrogen bonding moves downfield in the ^1H NMR spectra, which is a common feature of all hydrogen bond systems.

The terms “anagostic”¹¹ or “pregostic”¹² have been used to refer to M...H–C interactions which clearly do not fit the “agostic” definition. Structurally and spectroscopically, they are more similar to the hydrogen bonding M...H–X systems, but given the low ability of the C atom to act as a proton donor in a C–H fragment, their bonding description is still unclear. Nevertheless, some theoretical studies would seem to indicate that the d_z^2 orbital in d^8 square planar complexes is not involved in the interaction in certain cases.¹¹

While a fair amount of structural solid state studies have been carried out by X-ray or neutron diffraction,^{9,13,14} there is relatively scarce evidence of the existence of M...H hydrogen bridging in solution achieved through simultaneous observation of the downfield on the H signal and M–H coupling.^{6,15–17} The highest value of a $J(\text{Pt},\text{H})$ coupling constant (180 Hz) has been found in complex $[\text{PtBr}\{1\text{-C}_{10}\text{H}_6(\text{NMe}_2)\text{-}8\text{-C},\text{N}\}\{1\text{-C}_{10}\text{H}_6(\text{NHMe}_2)\text{-}8\text{-C},\text{H}\}]^+$.¹⁵ In this example, the hydrogen involved is very acidic because of the ammoniac nature of the N donor fragment. Smaller values of the coupling constants have been reported for amino donor fragments, such as in $[\text{Pt}(\text{C}_6\text{H}_3\text{-}2,6\text{E}_2)(8\text{-acetylaminequinoline})]^+$ ($\text{E} = \text{PPh}_2$, $J(\text{Pt},\text{H}) = 55$ Hz; NMe_2 , $J(\text{Pt},\text{H}) = 33$ Hz).¹⁶ Intermediate values of $J(\text{Pt},\text{H})$ (69 and 88 Hz, respectively) have been found in anionic complexes $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{hqH})]^-$ (**A**) and $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{hqH}')^-]$ (**B**, see Scheme 1),⁶ which contain the 8-hydroxyquinoline (hqH) or

Scheme 1



2-methyl-8-hydroxyquinoline (hqH') ligands, with a O–H donor fragment. Usually, with even less acidic C–H protons such as the ones contained in the 7,8-benzo[*h*]quinoline (bzqH) ligand of complex $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzqH})]^-$,⁶ the value observed for the Pt–H coupling constant is even lower (22 Hz). Nevertheless, higher $J(\text{Pt},\text{H})$ can be observed when the C–H group is strongly oriented toward the Pt atoms by the geometry of the ligand. This is the case of the complexes $[\text{Pt}(\text{Me}_3\text{Si-BAM})\text{Me}_2]$ and $[\text{Pt}(\text{Me}_3\text{Si-BAM})\text{Ph}_2]$ (BAM = bis(7-azaindol-1-yl)methane)¹⁷ for which values of 61.0 and 44.1 Hz, respectively, have been reported. Theoretical studies have also been carried on these complexes containing M...H–X hydrogen bonds.^{2,9,11,18–21} They indicate that most systems show an important electrostatic contribution interaction, as in “classic” hydrogen bonds. Nevertheless, it has been suggested that the importance of a covalent contribution increases as the M...H distance shortens and thus the strength of the interaction increases.^{2,11}

Recently, the first crystallographic evidence by neutron diffraction of intermolecular hydrogen bonding involving a d^8 metal center and a hydrogen atom of a crystallization water

molecule has been reported in the complex $\text{trans-}[\text{PtCl}_2(\text{NH}_3)\text{-}(\text{N-Glycine})]$.⁹ The structure also presents an intramolecular Pt...H–N interaction. Interestingly, theoretical studies of this system conclude that dispersion forces constitute the main component of the intermolecular Pt...H–O contact^{9,18,20} and also support the persistence of this interaction in solution.^{19,20}

In the course of previous research, we prepared anionic *tris* pentafluorophenyl Pt(II) complexes containing Pt...H–X ($\text{X} = \text{O}, \text{C}$) hydrogen bonds.⁶ In this paper we have explored the use of neutral Pt(II) pentafluorophenyl complexes which also contain the 7,8-benzoquinolate chelate planar ligand as precursors for complexes containing Pt...H–O hydrogen bonding with success. The overall charge in the complex and the different steric requirements of the ligands surrounding the metal center might influence the characteristics of the Pt...H interactions, which have been studied both in the solid state (X-ray) and in solution (NMR). Moreover, theoretical calculations have been performed to obtain a greater insight into the nature of the Pt...H interaction. For comparative purposes, studies on **A** and **B**, a couple of similar complexes previously prepared in our laboratory⁶ (see Scheme 1) have also been included in this paper. The study of the reactivity of the Pt...H complexes toward hydrogen abstractors has resulted in unexpected polynuclear complexes which have also been fully characterized.

Synthesis and Characterization of the Complexes $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})\text{L}]$ $\{\text{L} = \text{hqH}, 8\text{-hydroxyquinoline (1)}; \text{L} = \text{hqH}', 2\text{-methyl-}8\text{-hydroxyquinoline (2)}\}$. Complex $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})(\text{Me}_2\text{CO})]$ (bzq = 7,8-benzoquinolate) has proven to be a suitable precursor for the preparation of complexes $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})\text{L}]$ because the acetone group is easily replaced with other L ligands.^{22,23} Thus, the addition of equimolar amounts of 8-hydroxyquinoline (hqH) or 2-methyl-8-hydroxyquinoline (hqH') to dichloromethane solutions of $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})(\text{Me}_2\text{CO})]$ under protective Ar atmosphere and at 273 K allows one to obtain, after 15 min of stirring, the corresponding complexes $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{bzq})\text{L}]$ $\{\text{L} = \text{hqH (1)}, \text{hqH' (2)}\}$, see Scheme 1) as yellow solids which precipitate after partial evaporation of the solvent.

The IR spectra of complexes **1** and **2** confirm the replacement of the acetone in the starting material, since the ν_{CO} vibration band corresponding to this ligand which appears at 1669 cm^{-1} is no longer present, and bands assignable to the hqH and hqH' ligands can now be observed (see Experimental Section).

The ^{19}F NMR spectra of **1** and **2** present the same pattern of five signals, indicating that all the five fluorine atoms of the C_6F_5 ligands are inequivalent. The two *ortho*-F appear at lower field with platinum satellites. At higher field, one signal for the *para*-F and one for each of the *meta*-F can be observed. The inequivalence of the fluorine atoms in analogous positions of the pentafluorophenyl rings indicates the difficulty of this group to rotate around the Pt– C_{ipso} probably because of the bulkiness and rigidity of the neighboring chelating bzq ligand.

The ^1H NMR spectra of these complexes are more interesting. Figures 1 and 2 show these spectra for complexes **1** and **2**, respectively. They show the signals corresponding to the hydroxyquinoline ligands in the aromatic area besides the ones attributed to the bzq ligand with the expected relative intensity. Moreover, in the case of complex **2**, a singlet signal corresponding to the methylic hydrogen atoms appears at 3.40 ppm. However, the most striking feature of these spectra is the presence at low field of a sharp signal with platinum satellites assignable to the hydroxylic proton of the hydroxyquinoline




Figure 1. ^1H NMR spectrum of **1**. Inset: Detail of the signal of the OH hydrogen atom showing the ^{195}Pt satellites.




Figure 2. ^1H NMR spectrum of **2**. Inset: Detail of the signal of the OH hydrogen atom showing the ^{195}Pt satellites.

ligands. This signal appears at 10.92 ppm in **1**, with a coupling constant $J(\text{Pt},\text{H}) = 67.0$ Hz, and at 10.99 ppm in **2**, with a coupling constant $J(\text{Pt},\text{H}) = 80.6$ Hz. The downfield displacement of these signals with respect to the free ligands⁶ (2.64 ppm for **1** and 2.74 ppm for **2**) and, most importantly, the existence of the Pt–H coupling, accounts for the existence of the Pt···H–O hydrogen bond in solution for complexes **1** and

2. Similar Pt–H coupling constants have been reported for the related complexes $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{hqH})]^-$ (**A**) and $[\text{Pt}(\text{C}_6\text{F}_5)_3(\text{hqH}')^-]$ (**B**),⁶ (69 Hz, and 88 Hz, respectively, see Table 1). Nevertheless, the chemical displacement of the signal of the hydrogen involved in the interaction is greater (3.70 ppm and 4.09 ppm respectively).

Table 1. Relevant Structural and Magnetic Parameters Illustrating the Pt···H–O Contacts in Complexes **1**, **2**, **A**, and **B**^a

complex	1	2	A ^b	B ^b
Pt···H (X-ray), Å	2.09(4)	2.10(4)		2.19
Pt···H (DFT calculations, gas phase), Å	2.18	2.14	2.16	2.11
δ ¹ H NMR (CD ₂ Cl ₂), ppm	10.92	10.99	12.22	12.34
δ ¹ H NMR (DFT calculations, gas phase), ppm	10.71	10.95	11.95	12.07
$\Delta\delta$ ¹ H NMR (CD ₂ Cl ₂), ppm	+2.64	+2.74	+3.70	+4.09
$\Delta\delta$ ¹ H NMR (DFT calculations, gas phase), ppm	+3.02	+2.89	+4.26	+4.01
$J_{\text{Pt-H}}$, Hz	67.0	80.6	69	88
$T_{1, \text{min}}$ central signal, s ^c	1.55	1.47		
$T_{1, \text{min}}$ satellites, s ^c	1.45	1.28		
$\Delta T_{1, \text{min}}$	-0.10	-0.19		

^aSee Experimental Section for further details. ^bReference 6. ^c $T_{1, \text{min}}$ found at 193 K.

The close vicinity of the platinum center and the hydroxylic hydrogen atom is also manifest in another magnetic property of the latter, the relaxation rate.²⁴ Thus the $I = 1/2$ platinum isotope in the ¹⁹⁵Pt···H–O isotopomer makes an additional contribution to the relaxation rate of the corresponding hydrogen atom. As a consequence, the $T_{1(\text{min})}$ values measured for the satellite signals (which are due to the ¹⁹⁵Pt···H–O isotopomer) are slightly shorter than those measured for the central signal (which belong to the rest of the platinum isotopomers).²⁵ In CD₂Cl₂ as solvent, the measured $T_{1(\text{min})}$ values for the central singlet are 1.55 (**1**) and 1.45 (**2**) s, whereas those measured for the doublet corresponding to the ¹⁹⁵Pt isotopomers are 1.45 (**1**) and 1.28 (**2**) s. All those magnetic parameters are summarized in Table 1

The structures of complexes **1** and **2** have been established by single crystal X-ray diffraction studies. Figures 3 and 4 show




Figure 3. View of the molecular structure of the complex [Pt(C₆F₅)(bzq)(hqH)] (**1**). Ellipsoids are drawn at their 50% probability level.

views of the structures of **1** and **2** respectively, and Table 2 lists a selection of relevant bond distances and angles for both complexes. As expected, **1** and **2** are square planar complexes in which the pentafluorophenyl ligand is located trans to the nitrogen donor atom of the cyclometalated bzq ligand, as has previously been found in complexes with the formula [Pt(C₆F₅)(bzq)L].²³ In both structures, the bzq planes are




Figure 4. View of the molecular structure of the complex [Pt(C₆F₅)(bzq)(hqH')] (**2**). Ellipsoids are drawn at their 50% probability level.

Table 2. Selected Bond Lengths (Å) and Angles (deg) for [Pt(C₆F₅)(bzq)(hqH)]·CH₂Cl₂ (**1**·CH₂Cl₂) and [Pt(C₆F₅)(bzq)(hqH')]·CHCl₃ (**2**·CHCl₃)

	1 ·CH ₂ Cl ₂	2 ·CHCl ₃
Pt–C(17)	1.995(3)	1.990(2)
Pt–C(1)	2.012(3)	2.011(2)
Pt–N(1)	2.089(2)	2.083(2)
Pt–N(2)	2.144(2)	2.176(2)
Pt–H(1)	2.09(4)	2.10(4)
O–C(27)	1.355(4)	1.357(3)
O–H(1)	0.94(4)	0.91(4)
C(17)–Pt–C(1)	91.76(12)	92.57(9)
C(17)–Pt–N(1)	81.96(11)	81.80(8)
C(1)–Pt–N(1)	173.52(10)	174.24(8)
C(17)–Pt–N(2)	174.89(10)	174.90(8)
C(1)–Pt–N(2)	93.17(10)	90.03(8)
N(1)–Pt–N(2)	93.14(9)	95.67(7)
Pt–H(1)–O	162(4)	162(4)


coplanar to the Pt basal square planes (dihedral angle 3.4(1)° for **1** and 4.4(1)° for **2**), while hqH and C₆F₅ ligands are almost perpendicular to the latter (dihedral angles are 80.7(1)° and 78.5(1)° respectively for **1** and 84.0(1)° and 84.6(1)° for **2**). With these dispositions, the OH fragments of the hqH and hqH' ligands have optimal orientations for the hydrogen atoms to establish interactions with the Pt centers. It is noteworthy that the quality of the X-ray diffraction data collected has allowed, in both structures, to find and refine the position of these hydrogen atoms (H(1)) without restraints, and that from all the possible orientations, the hydrogen atoms are pointing toward the metal centers. Thus, the measured geometric parameters involving H(1) are Pt–H(1) = 2.09(4) Å, O–H(1) = 0.94(4) Å, Pt–H(1)–O 162(4)° for **1** and Pt–H(1) = 2.10(4) Å, O–H(1) = 0.91(4) Å, Pt–H(1)–O 162(4)° for **2**. If the O–H(1) distances are normalized to 0.993,²⁶ then the Pt–H(1) distances are 2.04 Å in **1** and 2.07 Å in **2**. In any case, these parameters are fully consistent with the existence of Pt···H(1)–O hydrogen bond systems in the solid state;^{2,3} in fact, the Pt–H(1) distances found in **1** and **2** (2.09(4) Å, 2.10(4) Å) are the shortest reported for this kind of hydrogen bonding.

Table 3. Comparison of Selected Distances (Å) and Angles (deg) Obtained for **1** (X-ray, DFT), **2** (X-ray, DFT), [Pt(C₆F₅)₃(hqH)][−] **A** (DFT), and **B** (X-ray,⁶ DFT)

	Pt···H	H–O	O–C	Pt···H–O	H–O–C	Pt···H–O–C
1 (X-ray)	2.09(4)	0.94(4)	1.355(4)	162(4)	110.2	−29.1
1 -DFT	2.18	0.983	1.338	153.6	112.7	−45.3
2 (X-ray)	2.10(4)	0.91(4)	1.357(3)	162(4)	109.8	−29.3
2 -DFT	2.15	0.983	1.343	152.4	111.7	−50.1
A -DFT	2.16	0.99	1.329	157.8	113.0	−20.8
B (X-ray) ⁶	2.19	0.84	1.354(6)	160.7	108.4	−32.0
B -DFT	2.11	0.99	1.331	153.9	113.1	−33.7

Slightly longer distances have been found in complexes [PtBr{1-C₁₀H₆(NMe₂)-8-C,N}{1-C₁₀H₆(NHMe₂)-8-C,H}] (2.11(5) Å),¹⁵ **B** (2.19 Å),⁶ and [Pt(C₆H₃-2,6(PPh₂)₂)(8-acetylaminoquinoline)](CF₃SO₃) (2.2(1) Å).¹⁶ Nevertheless, fine comparisons of the Pt–H distances must be performed with caution because of the inherent uncertainty of the location of the hydrogen atoms from single crystal X-ray diffraction studies. As expected, the Pt–H distance found in the structure of *trans*-[PtCl₂(NH₃)(N-Glycine)]·H₂O, determined by neutron diffraction,⁹ is much longer (2.885(3) Å), since it arises from an intermolecular interaction between the Pt center and a crystallization water molecule.

Computational Studies. The molecular structures of the complexes **1**, **2**, **A**, and **B** have been optimized by density functional theory (DFT) methods, at the M06 level of theory (see Experimental Section for further details). A comparison of the most relevant structural parameters of **1**, **1**-DFT, **2**, **2**-DFT, **A**-DFT, **B**, and **B**-DFT is included in Table 3. Views of the optimized structures for **1**-DFT, **2**-DFT, **A**-DFT, and **B**-DFT are included in Figure 5. The geometries of **1**-DFT and **2**-DFT

**Figure 5.** Optimized structures (DFT) for **1**, **2**, **A**, and **B**.

are consistent with the structures found by X-ray diffraction (see Figures S1 and S2, Supporting Information). In agreement with the existence of a Pt···H short contact, the Pt···H(1) distance is 2.18 Å, the H(1)–O bond length is 0.98 Å, and the Pt–H(1)–O is 153.6° for **1**-DFT, while for **2**-DFT the calculated parameters are 2.15, 0.98 Å, and 152.4°, respectively. These Pt–H distances are slightly longer than those determined crystallographically and support 4e-3c type interactions. The optimized geometry of **B** (**B**-DFT) is also consistent with the X-ray determined structure.⁶ It also shows

that the hydrogen atom of the hydroxyl fragment is oriented toward the platinum atom resulting in a very short distance of 2.11 Å (2.19 Å, X-ray), and all the Pt···H–O structural parameters are consistent with a hydrogen bonding 4e-3c type interaction (see Table 3). The optimized geometry of the analogous **A** (**A**-DFT) is very similar, thus supporting the existence of a Pt···H–O hydrogen bonding system.

The coherence observed for these structures has led us to investigate the Pt···H contacts in more detail through DFT methods. Topological analyses of the electron density function ($\rho(\mathbf{r})$) obtained for **1**-DFT, **2**-DFT, **A**-DFT, and **B**-DFT have been performed. According to Bader's *Atoms In Molecules* theory,^{27–31} the critical points (CPs) in the $\rho(\mathbf{r})$ function are the points in space at which the first derivatives of the function vanish (i.e., each individual derivative in the gradient operator is zero). CPs indicate chemically meaningful points and are classified according to their rank and signature. The rank is the number of nonzero curvatures of the electron density $\rho(\mathbf{r})$ at the CP, whereas the signature is the algebraic sum of the signs of the curvatures. For example, CPs of the (3,-3) type are indicative of the nuclear positions, whereas CPs of the (3,-1) type are evidence of chemical bonds. Complementarily, a bond path (BP) is a single line of maximum electron density linking the nuclei of two chemically bonded atoms. A BP is an indicator of chemical bonding of all kinds; weak, strong, closed-shell, and open-shell interactions. The point on the BP with the lowest electron density value (minimum along the path) is the bond critical point (BCP).

The analyses of the electron density functions in **1**-DFT, **2**-DFT, **A**-DFT, and **B**-DFT reveal a BP relating the platinum atom and the hydroxyl hydrogen atom in all four cases, as well as the corresponding BCPs. Observation of the properties of the electron density at the referred CPs sheds light onto the nature and properties of the discussed contacts. Cremer et al. have stated that to provide a thorough description of the CP, electrostatic and also energetic aspects must be considered.³² Thus, a negative value of $\nabla^2\rho(\mathbf{r})$ indicates a covalent (shared electron) interaction, while a positive value is associated with a closed-shell, electrostatic interaction. Complementarily, a negative value of the local energy density function $H(\mathbf{r})$ corresponds to partial covalence, while a positive $H(\mathbf{r})$ indicates a purely closed-shell, electrostatic interaction.^{11,27,28}

The results of our study on the referred BCPs found for **1**-DFT, **2**-DFT, **A**-DFT, and **B**-DFT are shown in Table 4. The electron density $\rho(\mathbf{r})$ at a BCP correlates with the strength of an atomic interaction. For a typical C–C covalent bond the value of $\rho(\mathbf{r})$ is about 1.7 au.³³ For conventional purely organic hydrogen bonding, values of between 0.0123 and 0.0276 au have been reported.³³ Bergès et al.²⁰ have found $\rho(\mathbf{r})$ values of about 0.020 au for the intermolecular Pt···H interactions between d⁸ square planar Pt(II) complexes and the water

Table 4. Topological Characteristics of Critical Point Pt...H–O in Complexes 1-DFT, 2-DFT, A-DFT, and B-DFT^a

complex	1-DFT	2-DFT	A-DFT	B-DFT
$\rho(r)$ (au)	0.034	0.036	0.035	0.039
$\nabla^2\rho(r)$	0.070	0.076	0.070	0.079
Ellipt	0.028	0.034	0.028	0.034
Pt...H (Å)	2.18	2.15	2.16	2.11
BP length (Å)	2.21	2.18	2.20	2.14
Pt–CP (Å)	1.51	1.49	1.51	1.47
CP–H (Å)	0.70	0.69	0.69	0.67
$G(r)$ (au)	0.022	0.024	0.022	0.026
$V(r)$ (au)	–0.026	–0.029	–0.027	–0.032
$H(r)$ (au)	–0.004	–0.005	–0.005	–0.006
$G(r)/\rho(r)$	0.648	0.661	0.630	0.655
$E(\text{HB}, \text{kcal mol}^{-1})$	–8.15	–9.11	–8.34	–9.92

^aBP: Bond path; CP: Critical point.

molecules discussed above. In the complexes studied in this paper, the values of $\rho(r)$ are higher, ranging from 0.034 to 0.039, and also higher than those reported by Oldfield and co-workers¹¹ for d⁸ square planar complexes containing intramolecular M...H–X (X = C, N) interactions (range 0.012–0.025), or by Pérez-Prieto and co-workers³⁴ in intramolecular M...H–C (M = Pd, Ag, Mo) interactions (range 0.019–0.034). Thus, these values of $\rho(r)$ at the BCP seem to indicate a significant Pt...H interaction in the complexes studied here. According to the $E_{\text{cont}} = 1/2V(r_{\text{CP}})$ relationship, the energies of the hydrogen bonds can be estimated to be between –8.1 and –10.0 kcal mol^{–1}.^{35,36}

With respect to the sign of the Laplacian $\nabla^2\rho(r)$, in all four cases their values are positive (see Table 4), thus indicating closed-shell, electrostatic interactions. This same result is also observed in conventional organic hydrogen bonds,³³ in the referred Pt(II)·H–OH interactions^{20,21} or in other studies on M...H–X systems^{11,34}. Nevertheless, and as stated before,^{11,33,34} considering the values and signs of $\nabla^2\rho(r)$ and $H(r)$ together allows a better understanding of these interactions. Thus, in the four examples studied here, all the $H(r)$ values are negative (see Table 4), which means that the Pt...H–O hydrogen bonds have *partial covalence*. Analogous results have been found in other pregeostic or hydrogen bonded M...H–X interactions,^{11,34} in contrast to conventional organic hydrogen bond systems for which positive values of $H(r)$ are always calculated, and thus no covalent component is deduced in the interaction.³³

It has been observed that a more negative value of $H(r)$ is related to the decrease in the distance between the interacting atoms, both in certain specific organic hydrogen bonding systems^{33,37} and in other nonbonded interactions.³⁸ In these specific hydrogen bonding systems the donor or proton acceptors are ylides³⁷ or organic acids,³³ and when known, the X...H distances have been shown to be very short. These “special” hydrogen bonds are sometimes termed “Low Barrier Hydrogen Bonds” (LBHB) and are postulated as transition states in several organic and enzyme catalytic events.^{33,37} The values of $H(r)$ reported for the complexes investigated here are the most negative ones calculated for M...H–X interactions^{11,34} (see Table 4), and certainly, the Pt–H distances calculated or measured by X-ray (complexes 1 and B, see Table 3), are among the shortest reported for this kind of complexes.^{6,15,16}

Some authors have stated that the shorter the Pt...H contact, the more negative the value of $H(r)$ is and the greater the downfield displacement of the interacting hydrogen is in the ¹H

NMR spectra.¹¹ In the case of the four complexes studied here, the calculated or measured values of the Pt–H distances are very similar, ranging from 2.09 Å to 2.19 Å (see Table 1). Nevertheless, the value of the downfield displacement of the signal of the interacting hydrogen in the ¹H NMR with respect to the free ligand is significantly greater for the complexes A and B (3.70 and 4.09 ppm, respectively) than it is for 1 and 2 (2.64 and 2.74 ppm, respectively). Thus, in these cases the Pt–H distance would seem to be not the only factor determining the magnitude of the downfield displacement in the signal of the hydrogen involved in the interaction.

Natural Bond Orbital (NBO) analyses have been performed for 1-DFT, 2-DFT, A-DFT, and B-DFT (see Supporting Information, Table S7). As a result of the anionic nature of the latter pair of complexes, the atomic charges on the platinum centers of 1-DFT and 2-DFT (+0.23 and +0.22) are higher than those in A-DFT and B-DFT (+0.11 and +0.09). This trend is also observed in the Mulliken charges (+0.09 in 1-DFT, +0.06 in 2-DFT, –0.09 in A-DFT, and –0.11 in B-DFT). Oppositely, the calculated charges on the hydroxyl hydrogen and on the oxygen atoms are almost identical in the series of four cases under study (average values are +0.52 for the hydrogen and –0.70 for the oxygen atoms according to NBO, and +0.41 for the hydrogen and –0.56 for the oxygen atoms according to Mulliken). The ¹H chemical shifts of the free quinoline and quinaldine molecules and of the 1-DFT, 2-DFT, A-DFT, and B-DFT complexes have also been calculated by DFT methods and compared to the experimental values (see Table 1). The calculated shifts for the hydroxylic hydrogen atoms in the organic substrates (7.69 and 8.06 ppm respectively) and in the four organometallic complexes (10.71, 10.95, 11.95, and 12.07 ppm respectively) are in a good agreement with the experimental ones, from both a qualitative and a quantitative point of view (see Table 1). Remarkably, the computed shifts reproduce the experimentally observed downfield delta shifts upon formation of the Pt...H interactions, and predict higher deshieldings for the hydroxylic proton of the aromatic ligands when coordinated to the anionic [Pt(C₆F₅)₃][–] fragment (+4.26 for A-DFT and +4.01 for B-DFT), than when coordinated to the neutral [Pt(C₆F₅)(bzq)] (+3.02 for 1-DFT and +2.89 for 2-DFT). The sum of these observations suggests that the strengths of the Pt...H interactions and the extent of the downfield shifts of the hydroxylic hydrogen atoms participating in the platinum–hydrogen contacts are determined not only by the short values of the Pt...H distances but also by the charges of the platinum centers in these sets of compounds. According to this, a more negative charge on the platinum center would favor stronger hydrogen bonds as well as more downfield chemical ¹H shifts in the resulting complexes.

Reactivity of Complexes 1 and 2 toward Bases. The hydrogen bridging M...H interactions have been described as the first step in processes of protonation of the metal and possible migration of the proton to a ligand with elimination of the protonated ligand.² For example, complex [Pt-(C₆F₅)₃(bzqH)][–], which shows evidence of the existence of the Pt...H interaction through Pt–H coupling in its ¹H NMR spectrum (see above), undergoes cyclometalation of the bzq ligand with elimination of C₆F₅H when it is refluxed in 1,2-dichloroethane for 3 h, giving rise to [Pt(C₆F₅)₂(bzq)][–].³⁹ Furthermore, the preparation of the starting material [Pt-(C₆F₅)(bzq)(Me₂CO)] is achieved via this pathway, with coordination of the Hbzq ligand to the platinum center in reflux

of acetone solutions of cis -[Pt(C₆F₅)₂(THF)₂]²³ However, complexes **1** and **2** do not undergo a similar process, with the cyclometalation of the hqH ligand, when their solutions are refluxed for several hours, even in relatively high boiling point solvents such as 1,2-dichloroethane.

Since the “internal” deprotonation, chelation, and corresponding formation of C₆F₅H do not take place, we tested the acidity of the OH hydrogen of the hydroxyquinoline ligands in **1** and **2** toward several “external” deprotonating reagents. Successful abstraction of the proton from the hydroxyl fragment should afford a monoanionic complex with a formally negative oxygen atom that could be used as a building block for preparing compounds of higher nuclearity, for example, by reaction with acidic metals such as Ag(I), Au(I), or Tl(I). Although no deprotonation was achieved using KOH, K(Me₃CO), or 1,8-bis(dimethylamino)naphthalene (proton sponge), the reactions with BuLi led to unexpected results. Thus, the treatment of tetrahydrofuran (THF) solutions of **1** or **2** with equimolar amounts of BuLi for 1 h at 198 K and the subsequent room temperature addition of (NBu₄)ClO₄ afforded, after workup, two yellow solids in good yields (see Experimental Section for details). The ¹H NMR spectra of these yellow solids do indeed show the absence of the downfield signal corresponding to the OH hydrogen atoms of the hydroxyquinoline ligands, with signals corresponding to the other protons of these and the bzq ligands in 1:1 ratio. They also show signals for NBu₄⁺, but with half the intensity expected for a (NBu₄)[Pt(C₆F₅)(bzq)(hq)] or (NBu₄)[Pt(C₆F₅)(bzq)(hq')] stoichiometry. Their ¹⁹F NMR spectra show signals for five fluorine atoms in the region and intensity expected.

The nature of the yellow solids could only be established when their structures were determined by single crystal X-ray diffraction. These studies concluded that the stoichiometry of the prepared complexes was in fact (NBu₄)[Li{Pt(C₆F₅)(bzq)(L)}₂] (L = hq (**3**), hq' (**4**)). For complex **4**, two sets of different crystals have been obtained, corresponding to two pseudopolymorphs,⁴⁰ one triclinic (*P* $\bar{1}$) which incorporates two CH₂Cl₂ solvent molecules in the asymmetric unit, and one monoclinic (*P*₂₁/*n*) with one CH₂Cl₂ molecule. The structural parameters for both pseudopolymorphs are very similar and will be denoted **4a** and **4b**, respectively. Figures 6 and 7 show views of the complexes **3** and **4a**, and Tables 5 and 6 list a selection of




Figure 6. View of the molecular structure of the anion of the complex (NBu₄)[Li{Pt(C₆F₅)(bzq)(hq)}₂] (**3**).




Figure 7. View of the molecular structure of the anion of the complex (NBu₄)[Li{Pt(C₆F₅)(bzq)(hq')}₂] (triclinic pseudopolymorph, **4a**).

their relevant bond distances and angles. A figure, a table of selected bond distances and angles, and crystallographic data for **4b** are included as Supporting Information.

Complexes **3**, **4a**, and **4b** are isostructural, with the obvious difference of the methyl substituent on the 2-methylhydroxyquinolate ligand, and small differences mainly in the conformation of the pentafluorophenyl and hydroxyquinolate ligands that probably can be accounted by the flexibility of the molecules in the crystalline environment.⁴¹ The three complexes are trinuclear, with two “Pt(C₆F₅)(bzq)(L)” subunits bridged by a lithium atom. The most remarkable feature of the structures is the change of coordination mode of the hydroxyquinolate ligands. Thus, the deprotonation of the hydroxyl group and the presence of the lithium atom cause a change in the bond between the platinum center and the hydroxyquinolate ligand, which now is established through a Pt–O bond. This change in the donor atom allows the oxygen atom to act as a bridge between the two metals. Furthermore, in this way, the lithium also coordinates to the now available nitrogen atom and is able to reach the four coordination with a distorted tetrahedral environment. A few lithium hydroxyquinolate complexes have been previously described in the literature, which are tetrametallic or hexametallic with cyclic structures in which the Li atoms are bridged by three oxygen atoms of the hq ligand and only the fourth coordination position is occupied by a N atom.^{42,43}

The two Pt square planes are nearly coplanar, in a disposition that is probably optimal to reduce the steric repulsions of the bulky benzoquinolate chelate ligands. Furthermore, the two bzq planes are separated about 3.4 Å, a distance that could indicate the existence of $\pi \cdots \pi$ interactions of the planar aromatic rings in a similar fashion to that previously reported for other complexes containing the bzq ligand.^{23,44–46}

CONCLUSION

The combination of the 8-hydroxyquinoline type ligands and square planar Pt(II) complexes has shown to be a good way to design complexes which contain Pt \cdots H–O hydrogen bonds. Thus, **1** and **2** are examples of complexes exhibiting this kind of interaction both in solution and in the solid state. Their ¹H NMR spectra show the two expected features that prove the interactions, namely, (a) downfield displacements of the signals of the involved hydrogen atoms and, even more conclusive, (b)

Table 5. Selected Bond Lengths (Å) and Angles (deg) for (NBu₄)[Li{Pt(C₆F₅)(bzq)(hq)}₂·1.875CH₂Cl₂ (3·1.875CH₂Cl₂)

Pt(1)–C(17)	1.978(4)	Pt(1)–C(1)	2.003(4)	Pt(1)–N(1)	2.076(3)
Pt(1)–O(1)	2.110(3)	Pt(2)–C(45)	1.981(4)	Pt(2)–C(29)	2.012(4)
Pt(2)–N(3)	2.071(3)	Pt(2)–O(2)	2.122(3)	Li–O(2)	1.886(7)
Li–O(1)	1.910(7)	Li–N(4)	2.030(7)	Li–N(2)	2.057(7)
C(17)–Pt(1)–C(1)	97.23(16)	C(17)–Pt(1)–N(1)	81.88(14)		
C(1)–Pt(1)–N(1)	173.30(14)	C(17)–Pt(1)–O(1)	174.75(13)		
C(1)–Pt(1)–O(1)	87.99(14)	N(1)–Pt(1)–O(1)	93.01(11)		
C(45)–Pt(2)–C(29)	95.78(16)	C(45)–Pt(2)–N(3)	82.00(15)		
C(29)–Pt(2)–N(3)	174.15(15)	C(45)–Pt(2)–O(2)	175.70(13)		
C(29)–Pt(2)–O(2)	87.70(13)	N(3)–Pt(2)–O(2)	94.29(12)		
O(2)–Li–O(1)	113.4(4)	O(2)–Li–N(4)	85.5(3)		
O(1)–Li–N(4)	129.1(4)	O(2)–Li–N(2)	130.8(4)		
O(1)–Li–N(2)	84.4(3)	N(4)–Li–N(2)	119.4(4)		

Table 6. Selected Bond Lengths (Å) and Angles (deg) for (NBu₄)[Li{Pt(C₆F₅)(bzq)(hq')}₂·2CH₂Cl₂ (4a·2CH₂Cl₂)

Pt(1)–C(17)	1.973(3)	Pt(1)–C(1)	2.003(3)	Pt(1)–N(1)	2.078(3)
Pt(1)–O(1)	2.111(2)	Pt(2)–C(46)	1.976(3)	Pt(2)–C(30)	2.002(4)
Pt(2)–N(3)	2.076(3)	Pt(2)–O(2)	2.100(2)	Li–O(2)	1.888(6)
Li–O(1)	1.903(6)	Li–N(2)	2.028(6)	Li–N(4)	2.058(6)
C(17)–Pt(1)–C(1)	96.04(13)	C(17)–Pt(1)–N(1)	81.98(12)		
C(1)–Pt(1)–N(1)	174.03(12)	C(17)–Pt(1)–O(1)	175.51(11)		
C(1)–Pt(1)–O(1)	88.09(11)	N(1)–Pt(1)–O(1)	94.09(10)		
C(46)–Pt(2)–C(30)	95.92(14)	C(46)–Pt(2)–N(3)	81.75(13)		
C(30)–Pt(2)–N(3)	175.35(12)	C(46)–Pt(2)–O(2)	176.02(11)		
C(30)–Pt(2)–O(2)	87.66(11)	N(3)–Pt(2)–O(2)	94.55(10)		
O(2)–Li–O(1)	115.0(3)	O(2)–Li–N(2)	128.2(3)		
O(1)–Li–N(2)	85.1(2)	O(2)–Li–N(4)	84.8(2)		
O(1)–Li–N(4)	127.2(3)	N(2)–Li–N(4)	121.7(3)		

the existence of Pt–H couplings with values of J of appreciable magnitude, indeed some of the largest reported so far. Moreover, the X-ray structures of **1** and **2** have revealed that the structural parameters of the fragment Pt–H–O are typical of a hydrogen bonding system. In particular, the Pt–H distances are very short, 2.09(4) and 2.10(4) Å, and indicate a quite strong interaction. The *Atoms in Molecules* study on complexes **1** and **2**, and also of the related complexes **A** and **B** (see Scheme 1), confirms the existence of interactions between the metal centers and the OH hydrogen atoms of electrostatic nature but with a *partial covalence*. This description is derived from the values of the Laplacian $\nabla^2\rho(\mathbf{r})$ and the local energy density function $H(\mathbf{r})$ calculated for the Pt···H–O systems. In particular, besides their sign, the magnitude of the value of $H(\mathbf{r})$ has also been directly related with the distance Pt–H, which correlates well with the observations for **1**, **2**, **A**, and **B**, and with the magnitude of the downfield displacement of the ¹H NMR signal of the hydrogen.¹¹ Nevertheless, in the cases studied in this paper the magnitude of this NMR displacement seems to be also related to the difference of charge between the Pt and the H, as suggested by NBO analyses of the complexes.


Several reagents have been tested as deprotonating agents for the hydroxyl OH in **1** and **2**, and only BuLi has proved to work properly. Nevertheless, the process evolves toward the formation of unexpected trinuclear Pt₂Li complexes, in which the coordination mode of the deprotonated 8-hydroxyquinolate type ligands changes. The hq and hq' ligands are now O-coordinated to the Pt atom, and O,N-chelate to the Li atom, in such a way that the Li is tetracoordinated and bridges two platinum subunits.

EXPERIMENTAL SECTION

General Comments. Literature methods were used to prepare the starting material [Pt(C₆F₅)(bzq)(Me₂CO)].²² Elemental analyses were carried out with a Perkin-Elmer 2400 CHNS analyzer. IR spectra were recorded on a Perkin-Elmer Spectrum 100 FT-IR spectrometer (ATR in the range 250–4000 cm⁻¹). Mass spectrometry was performed with the Microflex matrix-assisted laser desorption/ionization-time-of-flight (MALDI-TOF) Bruker or an Autoflex III MALDI-TOF Bruker instruments. NMR spectra in solution were recorded at 298 K on Bruker Avance 400 spectrometer with SiMe₄ and CFCl₃ as external references for ¹H, ¹³C, and ¹⁹F. The signal attributions and coupling constant assessment was made on the basis of a multinuclear NMR analysis of each compound including ¹H COSY, ¹H-¹³C HMQC, ¹H-¹³C HMBC, and APT.

Preparation of [Pt(C₆F₅)(bzq)(8-hydroxyquinoline)] (1**).** To a solution of [Pt(bzq)(C₆F₅)(Me₂CO)] (0.150 g, 0.251 mmol) in CH₂Cl₂ (20 mL) at 273 K and under Ar atmosphere, 0.251 mmol (0.036 g) of 8-hydroxyquinoline were added. After 15 min of stirring the solution was concentrated to about 2 mL. The yellow precipitate which appeared was filtered off, washed with *n*-hexane (10 mL), and air-dried. Yield 0.148 g (0.216 mmol), 86% yield. IR ν = 2946 (w, ν_{OH}), 1576 (vw), 1497 (m), 1450 (m), 1438 (m), 1208 (w), 1059 (m), 952 (s), 801 (m, C₆F₅, X-sensitive vibr.)⁴⁷ cm⁻¹. ¹H NMR (400.132 MHz, CD₂Cl₂, 298 K): δ = 10.92 (1H, s, J(H,Pt) = 67.0 Hz, H; hqH'–OH), 9.62 (1H, d, ³J(H2,H3) = 5.1 Hz, ³J(H2,Pt) = 23.4 Hz, H2, see Scheme 2 for the hydrogen and carbon numbering scheme), 8.44 (1H, dd, ³J(H4,H3) = 8.4 Hz, ⁴J(H4,H2) = 1.6 Hz, H4), 8.37 (1H, dd, ³J(H17,H16) = 8.1 Hz, ⁴J(H17,H15) = 1.3 Hz, H17), 7.85 (1H, d, ³J(H6,H5) = 8.8 Hz, H6), 7.65 (1H, d, ³J(H6,H5) = 8.8 Hz, H5), 7.64 (1H, dd, ³J(H7,H8) = 7.9 Hz, ⁴J(H7,H9) = 0.8 Hz, H7), 7.62 (1H, t, ³J(H19,H20) = ³J(H19,H18) = 7.7 Hz, H19), 7.60–7.52 (3H, m, overlapped signals of H15, H18 and H3), 7.39 (1H, dd, ³J(H8,H9) = 7.9 Hz, ³J(H8,H7) = 7.9 Hz, H8), 7.33 (1H, dd, ³J(H20,H19) = 7.7 Hz, ⁴J(H20,H18) = 1.5 Hz, H20), 7.29 (1H, dd, ³J(H16,H17) = 8.1 Hz, ³J(H16,H15) = 5.2 Hz, H16), 7.00 (1H, d,

Scheme 2



$^3J(\text{H}_9, \text{H}_8) = 7.8 \text{ Hz}$, $^3J(\text{H}_9, \text{Pt}) = 62.5 \text{ Hz}$, H9 ppm. ^{19}F NMR (376.479 MHz, CD_2Cl_2 , 298 K): $\delta = -117.9$ (*o*-F, m, $^3J(\text{F}, \text{Pt}) = 515 \text{ Hz}$), -121.2 (*o*-F, m, $^3J(\text{F}, \text{Pt}) = 444 \text{ Hz}$), -163.3 (*p*-F, t), -164.5 (*m*-F, br m), -165.0 (*m*-F, br m) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100.624 MHz, CD_2Cl_2 , 298 K): $\delta = 155.4$ (s, C12), 153.2 (s, C2), 152.2 (s, C21), 146.2 (s, C15), 142.7 (s, C11), 140.3 (s, C4), 138.6 (s, C17), 136.4 (s, C22), 135.9 (s, C10), 134.7 (s, $^2J(\text{C}, \text{Pt}) = 114 \text{ Hz}$, C9), 134.2 (s, C14), 132.3 (s, C23), 130.2 (s, C5), 129.8 (s, C8), 129.4 (s, C19), 123.5 (s, C6), 123.2 (s, C7), 122.2 (s, C16), 122.0 (s, C18), 120.3 (s, C3), 117.3 (s, C20) ppm. Mass spectra MALDI+ DCTB: $m/z = 517.0$ $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_9\text{H}_6\text{NO})]$, 684.0 $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_9\text{H}_6\text{NOH})\text{H}]^+$. Elemental analysis calcd (%) for $\text{C}_{28}\text{H}_{15}\text{F}_5\text{N}_2\text{O}$: C 49.06, H 2.21, N 4.09; found: C 48.72, H 2.25, N 4.02.

Preparation of $[\text{Pt}(\text{C}_6\text{F}_5)(\text{bzq})(2\text{-methyl-8-hydroxyquinoline})]$ (2). To a solution of $[\text{Pt}(\text{bzq})(\text{C}_6\text{F}_5)(\text{Me}_2\text{CO})]$ (0.150 g, 0.251 mmol) in CH_2Cl_2 (20 mL) at 273 K and under Ar atmosphere, 0.251 mmol (0.040 g) of 2-methyl-8-hydroxyquinoline were added. After 15 min of stirring the solution was concentrated to about 2 mL. The yellow precipitate which appeared was filtered off, washed with *n*-hexane (10 mL), and air-dried. Yield 0.154 g (0.220 mmol), 88% yield. IR $\nu = 2938$ (w, ν_{OH}), 1569 (w), 1504 (m), 1450 (m), 1438 (m), 1258 (m), 1060 (m), 955 (s), 797 (m, C_6F_5 , X-sensitive vibr.). ^{19}F NMR (400.132 MHz, CD_2Cl_2 , 298 K): $\delta = 10.99$ (1H, s, $J(\text{H}, \text{Pt}) = 80.6 \text{ Hz}$, H; hqH'-OH), 8.42 (1H, dd, $^3J(\text{H}_4, \text{H}_3) = 8.1 \text{ Hz}$, $^4J(\text{H}_4, \text{H}_2) = 1.2 \text{ Hz}$, H4, see Scheme 2 for the hydrogen and carbon numbering scheme), 8.27 (1H, d, $^3J(\text{H}_{17}, \text{H}_{16}) = 8.5 \text{ Hz}$, H17), 8.10 (1H, dd, $^3J(\text{H}_2, \text{H}_3) = 5.3 \text{ Hz}$, $^4J(\text{H}_2, \text{H}_4) = 1.2 \text{ Hz}$, $^3J(\text{H}_2, \text{Pt}) = 19.4 \text{ Hz}$, H2), 7.88 (1H, d, $^3J(\text{H}_6, \text{H}_5) = 8.8 \text{ Hz}$, H6), 7.67 (1H, d, $^3J(\text{H}_5, \text{H}_6) = 8.8 \text{ Hz}$, H5), 7.66 (1H, dd, $^3J(\text{H}_7, \text{H}_8) = 7.9 \text{ Hz}$, $^4J(\text{H}_7, \text{H}_9) = 0.7 \text{ Hz}$, H7), 7.52 (1H, t, $^3J(\text{H}_{19}, \text{H}_{20}) = ^3J(\text{H}_{19}, \text{H}_{18}) = 7.8 \text{ Hz}$, H19), 7.44–7.38 (3H, m, overlapped signals of H18, H8 and H3), 7.42 (1H, d, $^3J(\text{H}_{16}, \text{H}_{17}) = 8.5 \text{ Hz}$, H16), 7.28 (1H, dd, $^3J(\text{H}_{20}, \text{H}_{19}) = 7.8 \text{ Hz}$, $^4J(\text{H}_{20}, \text{H}_{18}) = 1.6 \text{ Hz}$, H20), 7.02 (1H, d, $^3J(\text{H}_9, \text{H}_8) = 7.2 \text{ Hz}$, $^3J(\text{H}, \text{Pt}) = 62.0 \text{ Hz}$, H9), 3.40 (3H, s, H; hqH'-CH₃) ppm. ^{19}F NMR (376.479 MHz, CD_2Cl_2 , 298 K): $\delta = -117.8$ (*o*-F, m, $^3J(\text{F}, \text{Pt}) = 435 \text{ Hz}$), -118.5 (*o*-F, $^3J(\text{F}, \text{Pt}) = 468 \text{ Hz}$), -163.5 (*p*-F, t), -164.9 (*m*-F, br m), -165.1 (*m*-F, br m) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100.624 MHz, CD_2Cl_2 , 298 K): $\delta = 163.5$ (s, C15), 155.6 (s, C12), 152.8 (s, C21), 146.8 (s, C2), 142.3 (s, C11), 140.8 (s, C17), 138.7 (s, C4), 137.2 (s, C22), 136.5 (s, C10), 134.6 (s, $^2J(\text{C}, \text{Pt}) = 110 \text{ Hz}$, C9), 134.4 (s, C14), 130.7 (s, C23), 130.4 (s, C6), 130.1 (s, C8), 128.4 (s, C19), 127.7 (s, C13), 124.5 (s, C16), 123.7 (s, C5), 123.2 (s, C7), 122.6 (s, C3), 120.5 (s, C18), 118.4 (s, C20) ppm. Mass spectra MALDI+ DCTB: $m/z = 531.0$ $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_{10}\text{H}_8\text{NOH})\text{H}]^+$, 699.0 $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_{10}\text{H}_8\text{NOH})\text{H}]^+$. Elemental analysis calcd (%) for $\text{C}_{29}\text{H}_{17}\text{F}_5\text{N}_2\text{O}$: C 49.79, H 2.45, N 4.01; found: C 49.87, H 2.14, N 4.10.

Preparation of $(\text{NBu}_4)[\text{Li}(\text{Pt}(\text{C}_6\text{F}_5)(\text{bzq})(8\text{-hydroxyquinoline})_2)]$ (3). To a solution of 1 (0.343 g, 0.500 mmol) in tetrahydrofuran (60 mL) at 195 K and under Ar atmosphere, BuLi (2.5 M solution in hexane; 0.220 mL, 0.550 mmol) was added. After 60 min of stirring the solution was allowed to reach room temperature, and the solution was hydrolyzed for 10 min to remove the excess of BuLi. The solution was evaporated to dryness, and the yellow solid was treated with *i*PrOH (10 mL) and NBu_4ClO_4 (0.085 g, 0.250 mmol) was added. The resultant yellow suspension was filtered off, washed with *n*-hexane (10 mL), and air-dried. Yield 0.281 g (0.174

mmol), 68% yield. IR $\nu = 2964$ (vw), 1567 (vw), 1493 (m), 1450 (m), 1436 (m), 1276 (vw), 1055 (m), 951 (s), 880 (w), 796 (m, C_6F_5 , X-sensitive vibr.), $^{47} 409$ (w) cm^{-1} . ^1H NMR (400.132 MHz, CD_2Cl_2 , 298 K): $\delta = 9.10$ (2H, dd, $^3J(\text{H}_2, \text{H}_3) = 5.1 \text{ Hz}$, $^4J(\text{H}_2, \text{H}_4) = 0.9$, H2, see Scheme 2 for the hydrogen and carbon numbering scheme), 8.14 (2H, dd, $^3J(\text{H}_{17}, \text{H}_{16}) = 8.4 \text{ Hz}$, $^4J(\text{H}_{17}, \text{H}_{15}) = 1.4 \text{ Hz}$, H17), 7.90 (2H, dd, $^3J(\text{H}_{15}, \text{H}_{16}) = 4.1 \text{ Hz}$, $^4J(\text{H}_{15}, \text{H}_{17}) = 1.4 \text{ Hz}$, H15), 7.74 (2H, d, $^3J(\text{H}_{20}, \text{H}_{19}) = 7.8 \text{ Hz}$, H20), 7.44 (2H, d, $^3J(\text{H}_6, \text{H}_5) = 8.7 \text{ Hz}$, H6), 7.38 (2H, d, $^3J(\text{H}_7, \text{H}_8) = 7.6 \text{ Hz}$, H7), 7.22 (2H, t, $^3J(\text{H}_{19}, \text{H}_{20}) = ^3J(\text{H}_{19}, \text{H}_{18}) = 8.0 \text{ Hz}$, H19), 7.20 (2H, dd, $^3J(\text{H}_{16}, \text{H}_{17}) = 8.4 \text{ Hz}$, $^3J(\text{H}_{16}, \text{H}_{15}) = 4.1 \text{ Hz}$, H16), 7.11 (2H, t, $^3J(\text{H}_8, \text{H}_9) = ^3J(\text{H}_8, \text{H}_7) = 7.5 \text{ Hz}$, H8), 6.97 (2H, dd, $^3J(\text{H}_{18}, \text{H}_{19}) = 8.0 \text{ Hz}$, $^4J(\text{H}_{18}, \text{H}_{20}) = 0.8 \text{ Hz}$, H18), 6.86 (4H, d, overlapped signals of H5 and H9), 6.67 (2H, d, $^3J(\text{H}_4, \text{H}_3) = 8.0 \text{ Hz}$, H4), 6.17 (2H, dd, $^3J(\text{H}_3, \text{H}_4) = 8.0 \text{ Hz}$, $^3J(\text{H}_3, \text{H}_2) = 5.1 \text{ Hz}$, H3), 2.70 (16H, m, $\alpha\text{-CH}_2\text{-NBu}_4^+$), 1.28 (16H, m, $\beta\text{-CH}_2\text{-NBu}_4^+$), 1.13 (16H, m, $\gamma\text{-CH}_2\text{-NBu}_4^+$), 0.84 (24H, t, $\text{CH}_3\text{-NBu}_4^+$) ppm. ^{19}F NMR (376.479 MHz, CD_2Cl_2 , 298 K): $\delta = -117.5$ (2*o*-F, m, $^3J(\text{F}, \text{Pt}) = 597 \text{ Hz}$), -166.6 (*m*-F, br m), -167.0 (*m*-F, br m), -167.3 (*p*-F, t) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100.624 MHz, CD_2Cl_2 , 298 K): $\delta = 166.2$ (s, C21), 153.6 (s, C12), 148.7 (s, C2), 146.0 (s, C15), 145.2 (s, C22), 142.8 (s, C11), 137.6 (s, C10), 136.7 (s, C17), 134.0 (s, C4), 133.4 (s, C14), 133.3 (s, $^2J(\text{C}, \text{Pt}) = 139 \text{ Hz}$, C9), 130.2 (s, C23), 129.0 (s, C8), 128.6 (s, C19), 127.8 (s, C6), 125.0 (s, C13), 123.8 (s, C5), 122.3 (s, C3), 121.0 (s, C16), 119.7 (s, C7), 114.9 (s, C20), 111.4 (s, C18), 59.0 (s, $\alpha\text{-CH}_2\text{-NBu}_4^+$), 24.1 (s, $\beta\text{-CH}_2\text{-NBu}_4^+$), 20.0 (s, $\gamma\text{-CH}_2\text{-NBu}_4^+$), 13.8 (s, $\text{CH}_3\text{-NBu}_4^+$). Mass spectra MALDI⁻ DCTB: $m/z = 557.0$ $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{OH})]$, 684.0 $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_9\text{H}_6\text{NOH})\text{H}]^+$, 1375.0 $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_9\text{H}_6\text{NO})\text{Li}(\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_9\text{H}_6\text{NO}))]^-$. Elemental analysis calcd (%) for $\text{C}_{72}\text{H}_{64}\text{F}_{10}\text{LiN}_5\text{O}_2\text{Pt}_2$: C 53.43, H 3.99, N 4.33; found: C 53.19, H 3.88, N 3.92.

Preparation of $(\text{NBu}_4)[\text{Li}(\text{Pt}(\text{C}_6\text{F}_5)(\text{bzq})(2\text{-methyl-8-hydroxyquinoline})_2)]$ (4). To a solution of 2 (0.350 g, 0.500 mmol) in tetrahydrofuran (60 mL) at 195 K and under Ar atmosphere, BuLi (2.5 M solution in hexane; 0.220 mL, 0.550 mmol) was added. After 60 min of stirring the solution was allowed to reach room temperature, and the solution was hydrolyzed for 10 min to remove the excess of BuLi. The solution was evaporated to dryness, and the yellow solid was treated with *i*PrOH (10 mL) and NBu_4ClO_4 (0.085 g, 0.250 mmol) was added. The resultant yellow suspension was filtered off, washed with *n*-hexane (10 mL), and air-dried. Yield 0.284 g (0.173 mmol), 69% yield. IR $\nu = 2963$ (vw), 1562 (vw), 1496 (m), 1450 (m), 1435 (m), 1274 (vw), 1057 (m), 952 (s), 880 (w), 796 (m, C_6F_5 , X-sensitive vibr.), $^{47} 356$ (w) cm^{-1} . ^1H NMR (400.132 MHz, CD_2Cl_2 , 298 K): $\delta = 9.10$ (2H, d, $^3J(\text{H}_2, \text{H}_3) = 5.1 \text{ Hz}$, H2, see Scheme 2 for the hydrogen and carbon numbering scheme), 8.03 (2H, d, $^3J(\text{H}_{17}, \text{H}_{16}) = 8.4 \text{ Hz}$, H17), 7.74 (2H, d, $^3J(\text{H}_{20}, \text{H}_{19}) = 7.8 \text{ Hz}$, H20), 7.42 (2H, d, $^3J(\text{H}_6, \text{H}_5) = 8.7 \text{ Hz}$, H6), 7.38 (2H, d, $^3J(\text{H}_7, \text{H}_8) = 7.8 \text{ Hz}$, H7), 7.15 (2H, t, $^3J(\text{H}_{19}, \text{H}_{20}) = ^3J(\text{H}_{19}, \text{H}_{18}) = 7.8 \text{ Hz}$, H19), 7.11 (2H, t, $^3J(\text{H}_8, \text{H}_9) = ^3J(\text{H}_8, \text{H}_7) = 7.8 \text{ Hz}$, H8), 7.09 (2H, d, $^3J(\text{H}_{16}, \text{H}_{17}) = 8.4 \text{ Hz}$, H16), 6.90 (2H, d, $^3J(\text{H}_{18}, \text{H}_{19}) = 7.8 \text{ Hz}$, H18), 6.84 (2H, d, $^3J(\text{H}_5, \text{H}_6) = 8.7 \text{ Hz}$, H5), 6.82 (2H, d, $^3J(\text{H}_9, \text{H}_8) = 7.8 \text{ Hz}$, H9), 6.65 (2H, d, $^3J(\text{H}_4, \text{H}_3) = 8.0 \text{ Hz}$, H4), 6.21 (2H, dd, $^3J(\text{H}_3, \text{H}_4) = 8.0 \text{ Hz}$, $^4J(\text{H}_3, \text{H}_2) = 5.1 \text{ Hz}$, H3), 2.71 (16H, m, $\alpha\text{-CH}_2\text{-NBu}_4^+$), 1.29 (16H, m, $\beta\text{-CH}_2\text{-NBu}_4^+$), 1.14 (16H, m, $\gamma\text{-CH}_2\text{-NBu}_4^+$), 0.85 (24H, t, $\text{CH}_3\text{-NBu}_4^+$) ppm. ^{19}F NMR (376.479 MHz, CD_2Cl_2 , 298 K): $\delta = -117.2$ (*o*-F, m, $^3J(\text{F}, \text{Pt}) = 557 \text{ Hz}$), -117.7 (*o*-F, m, $^3J(\text{F}, \text{Pt}) = 549 \text{ Hz}$), -166.8 (*m*-F, br m), -167.1 (*m*-F, br m), -167.6 (*p*-F, t) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100.624 MHz, CD_2Cl_2 , 298 K): $\delta = 165.6$ (s, C21), 155.5 (s, C15), 153.5 (s, C12), 148.7 (s, C2), 144.3 (s, C22), 142.8 (s, C11), 138.0 (s, C10), 137.1 (s, C17), 134.0 (s, C4), 133.4 (s, C14), 133.1 (s, $^2J(\text{C}, \text{Pt}) = 101 \text{ Hz}$, C9), 129.0 (s, C8), 128.3 (s, C23), 127.7 (s, C6), 127.6 (s, C19), 125.0 (s, C13), 124.0 (s, C5), 122.4 (s, C3), 121.7 (s, C16), 119.6 (s, C7), 115.1 (s, C20), 111.4 (s, C18), 58.9 (s, $\alpha\text{-CH}_2\text{-NBu}_4^+$), 24.1 (s, $\beta\text{-CH}_2\text{-NBu}_4^+$), 23.5 (s, $\gamma\text{-CH}_2\text{-NBu}_4^+$), 20.0 (s, $\text{CH}_3\text{-NBu}_4^+$). Mass spectra MALDI⁻ DCTB: $m/z = 698.0$ $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_{10}\text{H}_8\text{NOH})\text{H}]^+$, 1403.0 $[\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_{10}\text{H}_8\text{NO})\text{Li}(\text{Pt}(\text{C}_{13}\text{H}_8\text{N})(\text{C}_6\text{F}_5)(\text{C}_{10}\text{H}_8\text{NO}))]^-$.

Table 7. Crystal Data and Structure Refinement for Complexes **1**·CH₂Cl₂, **2**·CHCl₃, **3**·1.875CH₂Cl₂, and **4a**·2CH₂Cl₂

	1 ·CH ₂ Cl ₂	2 ·CHCl ₃	3 ·1.875CH ₂ Cl ₂	4a ·2CH ₂ Cl ₂
formula	C ₂₈ H ₁₅ F ₃ N ₂ O ₂ Pt ·CH ₂ Cl ₂	C ₂₉ H ₁₇ F ₃ N ₂ O ₂ Pt ·CHCl ₃	C ₇₂ H ₆₄ F ₁₀ LiN ₅ O ₂ Pt ₂ ·1.875CH ₂ Cl ₂	C ₇₄ H ₆₈ F ₁₀ LiN ₅ O ₂ Pt ₂ ·2CH ₂ Cl ₂
<i>M_t</i>	770.44	818.90	1777.64	1816.31
crystal system	monoclinic	monoclinic	triclinic	triclinic
space group	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> 2 ₁ / <i>n</i>	<i>P</i> $\bar{1}$	<i>P</i> $\bar{1}$
<i>a</i> /Å	10.3728(2)	10.9721(2)	14.3558(2)	13.8434(1)
<i>b</i> /Å	16.5660(3)	15.5286(2)	14.5503(2)	14.2761(2)
<i>c</i> /Å	15.0527(2)	16.6883(2)	17.2806(3)	18.7328(2)
α /deg	90	90	78.609(1)	79.406(1)
β /deg	96.075(2)	107.556(2)	81.501(1)	86.245(1)
γ /deg	90	90	80.284(1)	81.744(1)
<i>V</i> /Å ³	2572.1(1)	2710.9(1)	3463.2(1)	3598.4(1)
<i>Z</i>	4	4	2	2
<i>D_c</i> /g cm ⁻³	1.990	2.006	1.705	1.676
<i>T</i> /K	100(1)	100(1)	100(1)	100(1)
μ /mm ⁻¹	5.728	5.536	4.257	4.108
<i>F</i> (000)	1480	1576	1750	1792
2 θ range/deg	8.4–57.7	8.8–57.9	8.3–57.8	8.4–57.8
collected reflections	28470	30600	76190	77177
unique reflections	6142	6571	16370	17183
<i>R</i> _{int}	0.0249	0.0226	0.0337	0.0340
<i>R</i> ₁ , <i>wR</i> ₂ ^a (<i>I</i> > 2 σ (<i>I</i>))	0.0217, 0.0515	0.0174, 0.0424	0.0290, 0.0796	0.0276, 0.0745
<i>R</i> ₁ , <i>wR</i> ₂ ^a (all data)	0.0246, 0.0523	0.0191, 0.0432	0.0402, 0.0813	0.0396, 0.0769
GOF (<i>F</i> ²) ^b	1.048	1.059	1.040	1.024

$$^a R_1 = \frac{\sum(|F_o| - |F_c|)}{\sum|F_o|}, wR_2 = \left[\frac{\sum w(F_o^2 - F_c^2)^2}{\sum w(F_o^2)^2} \right]^{1/2}, ^b \text{Goodness-of-fit} = \left[\frac{\sum w(F_o^2 - F_c^2)^2}{(n_{\text{obs}} - n_{\text{param}})} \right]^{1/2}.$$

Elemental analysis calcd (%) for C₇₄H₆₈F₁₀LiN₅O₂Pt₂: C 53.98, H 4.16, N 4.25; found: C 53.66, H 4.33, N 3.99.

X-ray Structure Determinations. Crystal data and other details of the structure analyses are presented in Table 7. Suitable crystals for X-ray diffraction studies were obtained by slow diffusion of *n*-hexane into concentrated solutions of the complexes in 3 mL of CH₂Cl₂ (**1**, **3**, and **4**) or CHCl₃ (**2**). Crystals were mounted at the end of quartz fibers. X-ray intensity data were collected on an Oxford Diffraction Xcalibur diffractometer. The diffraction frames were integrated and corrected for absorption using the CrysAlis RED program.⁴⁸ The structures were solved by Patterson and Fourier methods and refined by full-matrix least-squares on *F*² with SHELXL-97.⁴⁹ All non-hydrogen atoms were assigned anisotropic displacement parameters and refined without positional constraints, except as noted below. For **1**·CH₂Cl₂, **3**·1.875CH₂Cl₂, and **4a**·2CH₂Cl₂, all hydrogen atoms were constrained to idealized geometries and assigned isotropic displacement parameters equal to 1.2 times the *U*_{iso} values of their attached parent atoms (1.5 times for the methyl hydrogen atoms), with the exception of the position of the hydrogen attached to the OH group of the hydroxyquinoline ligand (H(1)) in complex **1**·CH₂Cl₂, which was found in the electron density maps and allowed to refine with no positional or thermal restraints. For **2**·CHCl₃, the positions of all hydrogen atoms were found in the electron density maps and allowed to refine with no positional or thermal restraints. In the structure of **3**·1.875CH₂Cl₂, the dichloromethane solvent molecules were very diffuse, and restraints in their geometry and thermal parameters were used. In the structure of **4a**·2CH₂Cl₂, the γ -CH₂ and CH₃ groups of two of the butyl chains of the cation are disordered over two sets of positions refined with occupancy 0.7/0.3 and 0.6/0.4. Restraints were used in the geometry parameters involving these atoms. Full-matrix least-squares refinement of these models against *F*² converged to final residual indices given in Table 7.

CCDC-950300 (**1**), CCDC-950301 (**2**), CCDC-950302 (**3**), CCDC-950303 (**4a**), and CCDC-950304 (**4b**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Computational Details. Quantum mechanical calculations were performed with the Gaussian09 package⁵⁰ at the DFT/M06 level of theory.⁵¹ SDD basis set and its corresponding effective core potentials (ECPs) were used to describe the platinum atom.⁵² An additional set of *f*-type functions was also added.⁵³ Carbon, fluorine, hydrogen, nitrogen, and oxygen atoms were described with a 6-31G* basis set⁵⁴ except for the hydrogen atoms close to the metal (hydroxyl and methyl hydrogen atoms), which were described with a 6-31G** basis set.⁵⁵ The structures of the platinum complexes and hydroxyquinoline ligands were fully optimized with these basis sets and with no symmetry restrictions. All minima were subsequently characterized by analytically computing the Hessian matrix. Atomic coordinates (*x*, *y*, *z*) for the optimized structures are collected in the Supporting Information, Tables S3–S6. Topological analyses of the electron density distribution functions $\rho(\mathbf{r})$ were performed by using the AIMAll program package⁵⁶ based on the extended wave function obtained by M06 calculations. The AIM extended wave function format allows QTAIM analyses of molecular systems containing heavy atoms described with ECPs. Atomic charges were calculated by using the NBO analysis option as incorporated in Gaussian 09.⁵⁷ NMR chemical shifts were calculated in the previously optimized structures, but using the 6-311++g(d) basis set for all the light atoms in the molecules.^{58,59}

■ ASSOCIATED CONTENT

📄 Supporting Information

Further details of the structure determinations of **1**·CH₂Cl₂, **2**·CHCl₃, **3**·1.875CH₂Cl₂, **4a**·2CH₂Cl₂, and **4b**·CH₂Cl₂ in cif format. Tables of selected bond lengths and angles, crystal data and structure refinement, and a view of the molecular structure for **4b**. Tables of atomic coordinates for the optimized structures of **1**-DFT, **2**-DFT, **A**-DFT, and **B**-DFT. Table of calculated NBO and Mulliken atomic charges for complexes **1**-DFT, **2**-DFT, **A**-DFT, and **B**-DFT. Overlay drawings of the X-ray diffraction and DFT calculated structures of **1** and **2**. This

material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E mail: tello@unizar.es.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by the Spanish MICINN (DGPTC/FEDER) (Project CTQ2008-06669-C02-01/BQU) and MINECO (CTQ2012-35251), the Gobierno de Aragón (Grupo Consolidado E21: Química Inorgánica y de los Compuestos Organometálicos), and the Universidad de Zaragoza (JIUZ2012-CIE-02). The authors thank the Instituto de Biocomputación y Física de Sistemas Complejos (BIFI) and the Centro de Supercomputación de Galicia (CESGA) for generous allocation of computational resources.

DEDICATION

Dedicated to Prof. Antonio Laguna on the occasion of his 65th birthday.

REFERENCES

- Brookhart, M.; Green, M. L. H.; Parkin, G. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104*, 6908–6914.
- Brammer, L. *Dalton Trans.* **2003**, 3145–3157.
- Martín, A. J. *Chem. Educ.* **1999**, *76*, 578–583.
- Brookhart, M.; Green, M. L. H. *J. Organomet. Chem.* **1983**, *250*, 395–408.
- Epstein, L. M.; Shubina, E. S. *Coord. Chem. Rev.* **2002**, *231*, 165–181.
- Casas, J. M.; Falvello, L. R.; Forniés, J.; Martín, A.; Welch, A. J. *Inorg. Chem.* **1996**, *35*, 6009–6014.
- Casas, J. M.; Forniés, J.; Martín, A. *J. Chem. Soc., Dalton Trans.* **1997**, 1559–1563.
- Falvello, L. R. *Angew. Chem., Int. Ed.* **2010**, *49*, 10045–10047.
- Rizzato, S.; Bergès, J.; Mason, S. A.; Albinati, A.; Kozelka, J. *Angew. Chem., Int. Ed.* **2010**, *49*, 7440–7443.
- Chatterjee, S.; Krause, J. A.; Oliver, A. G.; Connick, W. B. *Inorg. Chem.* **2010**, *49*, 9798–9808.
- Zhang, Y.; Lewis, J. C.; Bergman, R. G.; Ellman, J. A.; Oldfield, E. *Organometallics* **2006**, *25*, 3515–3519.
- Bortolin, M.; Bucher, U. E.; Rügger, H.; Venanzi, I. M.; Albinati, A.; Lianza, F.; Trofimenko, S. *Organometallics* **1992**, *11*, 2514–2521.
- Brammer, L.; Charnock, J. M.; Goggin, P. L.; Goodfellow, R. J.; Orpen, A. G.; Koetzle, T. F. *J. Chem. Soc., Dalton Trans.* **1991**, 1789–1798.
- Braga, D.; Grepioni, F.; Tedesco, E.; Biradha, K.; Desiraju, G. R. *Organometallics* **1997**, *16*, 1846–1856.
- Wehmanoooyevaar, I. C. M.; Grove, D. M.; Kooijman, H.; Vandersluis, P.; Spek, A. L.; Vankoten, G. *J. Am. Chem. Soc.* **1992**, *114*, 9916–9924.
- Albinati, A.; Lianza, F.; Pregosin, P. S.; Muller, B. *Inorg. Chem.* **1994**, *33*, 2522–2526.
- Zhao, S. B.; Wang, R. Y.; Wang, S. *Organometallics* **2009**, *28*, 2572–2582.
- Kozelka, J.; Bergès, J.; Attias, R.; Fraita, J. *Angew. Chem., Int. Ed.* **2000**, *39*, 198–201.
- Vidossich, P.; Ortuño, M. Á.; Ujaque, G.; Lledós, A. *ChemPhysChem* **2011**, *12*, 1666–1668.
- Bergès, J.; Fourré, I.; Pilmé, J.; Kozelka, J. *Inorg. Chem.* **2013**, *52*, 1217–1227.
- Li, Y.; Zhang, G.; Chen, D. *Mol. Phys.* **2012**, *110*, 179–184.
- Berenguer, J. R.; Lalinde, E.; Moreno, M. T.; Sánchez, S.; Torroba, J. *Inorg. Chem.* **2012**, *51*, 11665–11679.
- Martín, A.; Belío, U.; Fuertes, S.; Sicilia, V. *Eur. J. Inorg. Chem.* **2013**, 2231–2247.
- Bakhmutov, V. I. *Practical Nuclear Magnetic Resonance Relaxation for Chemists*; John Wiley & Sons: Chichester, U.K., 2004.
- Crabtree, R. H.; Segmuller, B. E.; Uriarte, R. J. *Inorg. Chem.* **1985**, *24*, 1949–1950.
- Mercury CSD 3.1.1; Cambridge Crystallographic Data Centre: Cambridge, U.K., 2013.
- Bader, R. F. W. *Atoms in Molecules—a Quantum Theory*; Oxford University Press: Oxford, U.K., 1990.
- Bader, R. F. W. *J. Phys. Chem. A* **1998**, *102*, 7314–7323.
- Popelier, P. *Atoms in Molecules: An Introduction*; Pearson Education: Harlow, U.K., 2000.
- Gillespie, R. J.; Popelier, P. L. A. *Chemical Bonding and Molecular Geometry*; Oxford University Press: New York, 2001.
- The Quantum Theory of Atoms in Molecules*; Wiley-VCH: Weinheim, Germany, 2007.
- Cremer, D.; Kraka, E. *Angew. Chem., Int. Ed.* **1984**, *23*, 627–628.
- Arnold, W. D.; Oldfield, E. *J. Am. Chem. Soc.* **2000**, *122*, 12835–12841.
- Ciclosi, M.; Lloret, J.; Estevan, F.; Sanau, M.; Perez-Prieto, J. *Dalton Trans.* **2009**, 5077–5082.
- Espinosa, E.; Molins, E.; Lecomte, C. *Chem. Phys. Lett.* **1998**, *285*, 170–173.
- Espinosa, E.; Alkorta, I.; Rozas, I.; Elguero, J.; Molins, E. *Chem. Phys. Lett.* **2001**, *336*, 457–461.
- Rozas, I.; Alkorta, I.; Elguero, J. *J. Am. Chem. Soc.* **2000**, *122*, 11154–11161.
- Iwaoka, M.; Komatsu, H.; Katsuda, T.; Tomoda, S. *J. Am. Chem. Soc.* **2004**, *126*, 5309–5317.
- Forniés, J.; Ibáñez, S.; Martín, A.; Gil, B.; Lalinde, E.; Moreno, M. T. *Organometallics* **2004**, *23*, 3963–3975.
- Braga, D.; Grepioni, F. *Chem. Soc. Rev.* **2000**, *29*, 229–238.
- Martín, A.; Orpen, A. G. *J. Am. Chem. Soc.* **1996**, *118*, 1464–1470.
- Begley, W. J.; Rajeswaran, M. *Acta Crystallogr., Sect. E* **2006**, *62*, 1200–1202.
- Rajeswaran, M.; Begley, W. J.; Olson, L. P.; Huo, S. *Polyhedron* **2007**, *26*, 3653–3660.
- Forniés, J.; Ibáñez, S.; Martín, A.; Sanz, M.; Berenguer, J. R.; Lalinde, E.; Torroba, J. *Organometallics* **2006**, *25*, 4331–4340.
- Berenguer, J. R.; Díez, A.; Fernández, J.; Forniés, J.; García, A.; Gil, B.; Lalinde, E.; Moreno, M. T. *Inorg. Chem.* **2008**, *47*, 7703–7716.
- Forniés, J.; Gómez, J.; Lalinde, E.; Moreno, M. T. *Inorg. Chem.* **2001**, *40*, 5415–5419.
- Maslow, E. J. *Vibrational Spectra of Organometallic Compounds*; Wiley: New York, 1977.
- CysAlis RED, CCD camera data reduction program; Oxford Diffraction: Oxford, U.K., 2004.
- Sheldrick, G. M. *Acta Crystallogr.* **2008**, *A64*, 112–122.
- Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; J. Montgomery, A. J.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, O.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09, Revision A.02*; Gaussian, Inc.: Wallingford, CT, 2009.
- Zhao, Y.; Truhlar, D. G. *Theor. Chem. Acc.* **2008**, *120*, 215–241.

- (52) Andrae, D.; Haussermann, U.; Dolg, M.; Stoll, H.; Preuss, H. *Theor. Chim. Acta* **1990**, *77*, 123–141.
- (53) Ehlers, A. W.; Böhme, M.; Dapprich, S.; Gobbi, A.; Höllwarth, A.; Jonas, V.; Köhler, K. F.; Stegmann, R.; Veldkamp, A.; Frenking, G. *Chem. Phys. Lett.* **1993**, *208*, 111–114.
- (54) Hariharan, P. C.; Pople, J. A. *Theor. Chim. Acta* **1973**, *28*, 213–222.
- (55) Francl, M. M.; Pietro, W. J.; Hehre, W. J.; Binkley, J. S.; Gordon, M. S.; DeFrees, D. J.; Pople, J. A. *J. Chem. Phys.* **1982**, *77*, 3654–3665.
- (56) Todd, A.; Keith, T. K. *AIMAll*, version 13.05.06; Gristmill Software: Overland Park, KS, 2013.
- (57) Glendening, E. D.; Reed, A. E.; Carpenter, J. E.; Weinhold, F. *NBO*, Version 3.1.
- (58) Feller, D. *Comput. Chem.* **1996**, *17*, 1571–1586.
- (59) Schuchardt, K. L.; Didier, B. T.; Elsethagen, T.; Sun, L.; Gurumoorthi, V.; Chase, J.; Li, J.; Windus, T. L. *J. Chem. Inf. Model.* **2007**, *47*, 1045–1052.