

# Kinetic and Mechanistic Studies on the Reaction of the Vitamin B<sub>12</sub> Complex Aquacobalamin with the HNO Donor Angeli's Salt: Angeli's Salt and HNO React with Aquacobalamin


Harishchandra Subedi,<sup>†</sup> Hanaa A. Hassanin,<sup>†,‡</sup> and Nicola E. Brasch<sup>\*,†,§</sup>

<sup>†</sup>Department of Chemistry and Biochemistry and <sup>§</sup>School of Biomedical Sciences, Kent State University, Kent, Ohio 44242, United States

<sup>‡</sup>Department of Chemistry, Ain Shams University, Abbassia Square, Cairo, 11566, Egypt

## Supporting Information

**ABSTRACT:** We report the first studies on the reaction between an HNO donor compound and vitamin B<sub>12</sub> complexes. Kinetic and mechanistic studies have been carried out on the reaction between the vitamin B<sub>12</sub> derivative aquacobalamin (H<sub>2</sub>OCbl<sup>+</sup>/HOcbl; pK<sub>a</sub> = 7.8) and the HNO donor Angeli's salt. Studies were carried out with aquacobalamin in excess, since nitrite also reacts with aquacobalamin to form nitrocobalamin (NO<sub>2</sub>Cbl). At pH <9.90 aquacobalamin reacts directly with the monoprotonated form of Angeli's salt, HN<sub>2</sub>O<sub>3</sub><sup>-</sup>, to form nitroxylcobalamin (NO<sup>-</sup>-Cbl(III)); NOcbl and nitrite. At pH >10.80 the reaction instead switches predominantly to a mechanism in which spontaneous decomposition of Angeli's salt to give HNO and nitrite becomes the rate-determining step, followed by the rapid reaction between aquacobalamin and HNO/NO<sup>-</sup> to again give NOcbl. Both reactions proceed with a 1:1 stoichiometry and formation of nitrite is confirmed using the Griess assay.




## INTRODUCTION


There is currently much interest in the chemical and biochemical reactivity of the reactive nitrogen species nitrosyl hydride (nitroxyl, HNO).<sup>1–8</sup> HNO is generated in vivo by nitric oxide synthases from L-arginine in the absence of the tetrahydrobiopterin cofactor or by the oxidation of *N*-hydroxy-L-arginine.<sup>9–11</sup> Reduction of NO by enzymes such as ferrocyanochrome *c*<sup>12</sup> and superoxide dismutase<sup>13</sup> leads to HNO generation. Spontaneous decomposition of *S*-nitrosothiols or their reactions with thiols may also generate HNO.<sup>10,11,14</sup> Like NO, elevated HNO levels are associated with damage to cellular components leading to nitrosative stress.<sup>15</sup> Thus far HNO formation in biological systems has not been unequivocally demonstrated, and multiple groups are currently focused on designing molecules which efficiently trap HNO.<sup>16–21</sup>

An interesting aspect of HNO-associated chemistry is the spin state change which occurs upon deprotonation of HNO in alkaline solution (the pK<sub>a</sub> for deprotonation of <sup>1</sup>HNO to give <sup>3</sup>NO<sup>-</sup> (+H<sup>+</sup>) is ~11.4).<sup>10,22–24</sup> Furthermore, HNO has a unique chemical and biological reactivity distinct from NO.<sup>9,25–27</sup> For example, HNO reacts directly with thiols while NO does not.<sup>9</sup> HNO inhibits aldehyde dehydrogenase, and as such may be useful in treating alcoholism.<sup>28</sup> Like NO, HNO is also a vasorelaxant<sup>29</sup> and shows potential in treating cardiovascular disease and preventing ischemia/reperfusion injury and congestive heart failure.<sup>9,25,30</sup> However, one major challenge in elucidating the chemical and biochemical reactivity of HNO stems from its rapid, spontaneous dimerization and

subsequent decomposition to nitrous oxide and water in aqueous solution, eq 1.<sup>31</sup>




Owing to its inherent instability, HNO donor compounds are therefore required to generate HNO in situ, and indeed the design of molecules that release HNO within a useful time frame is an active area of research.<sup>32–34</sup> Currently, sodium trioxodinitrate or Angeli's salt (Na<sub>2</sub>N<sub>2</sub>O<sub>3</sub>, AS), first synthesized by Angeli in 1903,<sup>35</sup> is the most widely used HNO donor in chemical and biochemical studies of HNO reactivity.<sup>36</sup> AS (HN<sub>2</sub>O<sub>3</sub><sup>-</sup>, pK<sub>a</sub>(HN<sub>2</sub>O<sub>3</sub><sup>-</sup>) = 9.70; pK<sub>a</sub>(H<sub>2</sub>N<sub>2</sub>O<sub>3</sub>) = 2.51<sup>37</sup>) decomposes rapidly to give HNO and NO<sub>2</sub><sup>-</sup> at pH 4–8, eq 2, whereas at pH <4, AS is primarily a NO donor.<sup>36,38</sup>



Vitamin B<sub>12</sub> derivatives (also known as cobalamins; Cbls) are a class of redox active cobalt containing complexes belonging to the corrinoid family which are synthesized by a number of microorganisms.<sup>39</sup> Cobalamins are essential coenzymes in all mammalian cells. The two B<sub>12</sub>-dependent enzyme reactions in humans require either adenosylcobalamin (AdoCbl, X = 5'-deoxy-5'-adenosyl (Ado), Figure 1) or methylcobalamin

Received: October 16, 2013

Published: January 17, 2014



**Figure 1.** Structure of vitamin B<sub>12</sub> derivatives (cobalamins): X = CN<sup>-</sup>, CH<sub>3</sub>, Ado, H<sub>2</sub>O, OH<sup>-</sup>, NO<sup>-</sup>, NO<sub>2</sub><sup>-</sup>, etc.

(MeCbl, X = CH<sub>3</sub>, Figure 1).<sup>40</sup> Cbl deficiency in humans is associated with hyperhomocysteinemia and/or methylmalonic acidemia leading to megaloblastic anemia and/or neurological disorders.<sup>41</sup> Cyanocobalamin (CNCbl, X = CN<sup>-</sup>, Figure 1) is the most common form of vitamin B<sub>12</sub> used in vitamin supplements.<sup>42</sup>

Multiple studies have recently been published on the chemistry of HNO donor molecules with porphyrin complexes.<sup>1,21,43–46</sup> Thus far no studies have been reported on the reactivity of HNO or HNO donor molecules with structurally similar vitamin B<sub>12</sub> complexes, although Cbls are well-known to modulate NO-associated events in biological systems.<sup>47–51</sup> We have carried out kinetic and mechanistic studies on the reaction of the cob(III)alamin aquacobalamin/hydroxycobalamin (H<sub>2</sub>Ocbl<sup>+</sup>/HOcbl; X = H<sub>2</sub>O/OH<sup>-</sup>, Figure 1) with the HNO donor Angeli's salt. Our studies show that H<sub>2</sub>Ocbl<sup>+</sup>/HOcbl reacts either directly with AS and/or with its decomposition product, HNO, depending upon the pH conditions used.

## EXPERIMENTAL SECTION

**Reagents.** Hydroxycobalamin hydrochloride (HOcbl·HCl, 98% stated purity by the manufacturer) was purchased from Fluka. The percentage of water in HOcbl·HCl ( $n\text{H}_2\text{O}$ ) (10–15% water, batch dependent) was determined by converting HOcbl·HCl into dicyanocobalamin, (CN)<sub>2</sub>Cbl<sup>-</sup> (0.10 M KCN, pH 11.0,  $\epsilon_{368\text{ nm}} = 3.04 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ).<sup>52,53</sup> Angeli's salt (AS) was purchased from Cayman Chemical and used without further purification. NaBH<sub>4</sub> ( $\geq 98\%$ ), Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> ( $\geq 85\%$ ), NaNO<sub>2</sub> (99.6%), NH<sub>2</sub>OH·HCl ( $\geq 97\%$ ), KCN ( $\geq 99.1\%$ ), 8-hydroxyquinoline ( $\geq 99\%$ ), diethylenetriaminepentaacetic acid (DTPA;  $\geq 98\%$ ), D<sub>2</sub>O (99.8 atom % D), acetone, triflic acid, NaOH, biological buffers (MES, TES, TAPS, and CHES), and phosphate buffers (NaH<sub>2</sub>PO<sub>4</sub> and Na<sub>2</sub>HPO<sub>4</sub>) were obtained from either Fisher Scientific or Acros Organics. TSP (3-(trimethylsilyl)propionic 2,2,3,3-*d*<sub>4</sub> acid, sodium salt) and the Griess reagent were obtained from Sigma Aldrich. Nessler's reagent was purchased from RICCA Chemical. Water was purified using a Barnstead Nanopure Diamond water purification system.

**General Methods.** All solutions were prepared using standard biological buffers and phosphate buffers (0.05–0.30 M), and a constant ionic strength was maintained using sodium triflate (NaCF<sub>3</sub>SO<sub>3</sub>;  $I = 1.0 \text{ M}$ ). Anaerobic solutions were prepared by bubbling argon through the solutions for ~24 h. Stock solutions were stored in the MBRAUN Labmaster 130 (1250/78) glovebox filled with argon, equipped with O<sub>2</sub> and H<sub>2</sub>O sensors and a freezer at -24 °C. Temperature-sensitive solutions were stored in the freezer. All pH

measurements were carried out at room temperature using an Orion Model 710A pH meter equipped with Mettler-Toledo Inlab 423 or 421 electrodes. An electrode was filled with 3 M KCl/saturated AgCl solution (pH 7). The electrodes were calibrated with standard buffer solutions at pH 4.00, 7.00, 10.00, and 12.45. Air-free UV–vis spectrometric measurements were carried out in Schlenk cuvettes (cuvettes fitted with a J-Young or an equivalent stopcock) on a Cary 5000 spectrophotometer equipped with a thermostatted (25.0 ± 0.1 °C) cell changer operating with WinUV Bio software (version 3.00). Freshly prepared solutions were used for kinetic measurements. For rapid reactions, kinetic data were collected under strictly anaerobic conditions at 25.0 ± 0.2 °C using an Applied Photophysics SX20 stopped-flow instrument equipped with a photodiode array detector, operating with Pro-Data SX (version 2.1.4) and Pro-Data Viewer (version 4.1.10) software, using either a 2 or 10 mm path length cell. The instrument was continuously purged with nitrogen gas during data collection and pretreated with anaerobic sodium dithionite (at least for 1 h) to remove oxygen and thoroughly washed with plenty of anaerobic water. Hamilton gastight syringes filled with the anaerobic reactant solutions in the glovebox were used to introduce the reactant solutions into the reservoir syringes of the stopped-flow instrument. Data were fitted using the program Microcal Origin (version 8.0). Data simulations were carried out using Pro-Kineticist software (Applied Photophysics). <sup>1</sup>H NMR spectra were recorded on a Bruker 400 MHz spectrometer equipped with a 5 mm probe at 23 ± 1 °C. TSP was used as internal reference. For <sup>1</sup>H NMR experiments under anaerobic conditions, airtight J-Young NMR tubes (Wilmad, 535-JY-7) were used. Reaction mixtures were equilibrated for 15 min prior to measurements.


**Determination of Cobalamin (Cbl) Concentrations.** The Cbl concentrations were determined by converting Cbl's to dicyanocobalamin, (CN)<sub>2</sub>Cbl<sup>-</sup>. Cobalamins were allowed to react with KCN (0.10 M, pH 11.50) to produce (CN)<sub>2</sub>Cbl<sup>-</sup>. The concentration of the final product was determined using UV–vis spectrometry ( $\epsilon_{368\text{ nm}} = 3.04 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ).<sup>52,53</sup>

**Study of the Spontaneous Decomposition of AS.** Rate constants for the spontaneous decomposition of AS as a function of pH under anaerobic conditions were measured using UV–vis spectrophotometry by following the decay in the Angeli's salt absorbance at 245 nm. The reaction was initiated by adding an aliquot (0.050 mL, 6.00 mM) of a stock solution of AS in 10.0 mM NaOH to a cuvette containing buffer (2.95 mL) which had been thermostatted in the cell holder of the Cary 5000 spectrophotometer. CAPS, CHES, and TAPS buffers (0.30 M) were used, and the total ionic strength was maintained at 1.0 M (NaCF<sub>3</sub>SO<sub>3</sub>). The absorbance at 245 nm versus time data were fitted to a first-order rate equation.

**Griess assay for the Quantification of Nitrite.** The amount of nitrite in the product solutions was determined using the Griess assay<sup>54</sup> under strictly anaerobic conditions. In a typical experiment, an aliquot of Griess reagent (1.50 mL) was added to an equal volume of buffer (0.1 M, 1.50 mL) containing varying concentrations of nitrite (0, 20.0, 40.0, 60.0, 80.0, and 100.0 μM) and NOcbl (1.00 × 10<sup>-4</sup> M) and the absorbance at 586 nm was determined. The resulting calibration curve is shown in Figure S13 in the Supporting Information (note that the calibration curves were found to be pH independent). An aliquot of the product of the reaction between HOcbl (1.0 × 10<sup>-4</sup> M) and AS (1.0 equiv) was subjected to the identical procedure. The concentration of nitrite was estimated from the calibration plot.

**Determination of Percentage of Nitrite Impurity in Commercially Available AS.** The percentage nitrite impurities in commercially available AS was determined using the Griess assay<sup>54</sup> under strictly anaerobic conditions. A typical procedure is as follows: Griess reagent (1.50 mL) was added to a solution of AS (1.17 and 1.43 mM; carried out in duplicate) in 0.10 M CAPS buffer, pH 10.80, and the absorbance was measured at 586 nm. The concentration of nitrite was obtained from the calibration curve (as described elsewhere). The percentage of nitrite impurity was found to be 4.8 ± 0.1%.

**Sample Preparation for Kinetic Measurements on the Reaction of Cbl(III) (H<sub>2</sub>Ocbl<sup>+</sup>/HOcbl) with AS.** All samples were prepared under strictly anaerobic conditions inside the glovebox. Stock



**Figure 2.** (a) UV-vis spectra for the reaction between HOCbl ( $5.00 \times 10^{-5}$  M) and excess AS ( $2.50 \times 10^{-3}$  M) at pH 9.80 (25.0 °C, 0.30 M CHES buffer,  $I = 1.0$  M ( $\text{NaCF}_3\text{SO}_3$ )). Isosbestic points occur at 341, 370, and 498 nm. **Inset:** First and last spectra. (b) Fit of absorbance data at 356 nm versus time to a first-order rate equation, giving  $k_{\text{obs}} = (2.85 \pm 0.02) \times 10^{-3} \text{ s}^{-1}$ .

Cbl(III) solutions were prepared by dissolving solid HOCbl·HCl in the appropriate buffer. Stock AS solutions were prepared by dissolving solid AS in NaOH (10.0 mM). Stock solutions were stored in the freezer (−24 °C) inside the glovebox and used within 24 h. Typically a small aliquot (~0.10 mL) of the AS solution was added to a solution of Cbl(III) (~2.90 mL) in buffer.


## RESULTS AND DISCUSSION

**Studies on the Reaction between  $\text{H}_2\text{OCbl}^+/\text{HOCbl}$  and Angeli's salt (AS) at pH  $\leq 9.90$ .** Upon the addition of AS to a solution of aquacobalamin/hydroxycobalamin ( $\text{H}_2\text{OCbl}^+/\text{HOCbl}$ ;  $\text{p}K_{\text{a}} = 7.8^{55}$ ) under strictly anaerobic conditions, significant changes in the UV-vis spectrum were observed and the reaction solution changed from red to orange. Figure 2a shows typical UV-vis spectra for the reaction between HOCbl ( $5.00 \times 10^{-5}$  M) and excess AS ( $2.50 \times 10^{-3}$  M) as a function of time under anaerobic conditions at pH 9.80 (25.0 °C, 0.30 M CHES buffer,  $I = 1.0$  M,  $\text{NaCF}_3\text{SO}_3$ ). The inset to Figure 2a shows a comparison between the initial spectrum and final spectrum, and it is clear that the product is nitroxy-cobalamin,  $\text{NO}^- \text{-Cbl(III)}$  ( $\lambda_{\text{max}} = 256, 278$  (shoulder), 289, 315, and 478 nm), with sharp isosbestic points observed at 341, 370, and 498 nm, in agreement with literature values for the HOCbl to NOCbl conversion.<sup>56</sup> Figure 2b gives the best fit of the absorbance data at 356 nm versus time to a first-order rate equation, giving an observed rate constant ( $k_{\text{obs}} = (2.85 \pm 0.02) \times 10^{-3} \text{ s}^{-1}$  ( $t_{1/2} = 4.1$  min)).

In order to be able to directly compare rate constants for the reaction between AS and  $\text{H}_2\text{OCbl}^+/\text{HOCbl}$  with rate constants for spontaneous AS decomposition, observed rate constants for AS decomposition ( $k_{\text{L}}$ ) were determined under the identical experimental conditions (25.0 °C,  $I = 1.0$  M ( $\text{NaCF}_3\text{SO}_3$ )) in the pH 7.00–11.40 range by following the decay of AS at 245 nm. The resulting plot of  $k_{\text{L}}$  versus pH is shown in Figure S1 in the Supporting Information, giving a first-order rate constant for  $\text{HN}_2\text{O}_3^-$  decomposition,  $k_{\text{AS}} = (5.16 \pm 0.23) \times 10^{-4} \text{ s}^{-1}$  and  $\text{p}K_{\text{a}}(\text{HN}_2\text{O}_3^-) = 9.48 \pm 0.16$ . These values are in very good agreement with literature values ( $\text{p}K_{\text{a}}(\text{HN}_2\text{O}_3^-) = 9.70^{37}$  and  $k_{\text{AS}} = 6.7 \times 10^{-4} \text{ s}^{-1}$  (25 °C,  $I = 0.25$  M)<sup>57,58</sup>).


The rate constant for the spontaneous acid-catalyzed decomposition of AS at pH 9.80 ( $k_{\text{L}} = (1.34 \pm 0.01) \times 10^{-4} \text{ s}^{-1}$ ; Supporting Information, Figure S1) was found to be more than 1 order of magnitude (~20 times) slower than the

reaction of AS ( $2.50 \times 10^{-3}$  M) with HOCbl ( $5.00 \times 10^{-5}$  M) at pH 9.80 ( $k_{\text{obs}} = (2.85 \pm 0.02) \times 10^{-3} \text{ s}^{-1}$ ). Therefore, decomposition of AS is not important under these reaction conditions and a direct reaction occurs between  $\text{H}_2\text{OCbl}^+/\text{HOCbl}$  and AS. The dependence of  $k_{\text{obs}}$  on AS concentration (2.50–10.0 mM) at pH 9.80 was determined, and the data are summarized in Figure 3. Fitting the data to a straight line passing through the origin gives a slope of  $1.18 \pm 0.01 \text{ M}^{-1} \text{ s}^{-1}$ .



**Figure 3.** Plot of  $k_{\text{obs}}$  versus AS concentration at pH 9.80 (25.0 °C, 0.30 M CHES buffer,  $I = 1.0$  M ( $\text{NaCF}_3\text{SO}_3$ )). The data have been fitted to a straight line passing through the origin, giving a second-order rate constant,  $k_{\text{app}} = 1.18 \pm 0.01 \text{ M}^{-1} \text{ s}^{-1}$ .

It occurred to us that nitrite could interfere with the determination of the rate constant for the reaction of interest, since  $\text{H}_2\text{OCbl}^+$  reacts rapidly with nitrite to form nitro-cobalamin,  $\text{NO}_2\text{Cbl}$ .<sup>59</sup> There are several possible sources of nitrite. Nitrite is produced in the reaction between  $\text{H}_2\text{OCbl}^+/\text{HOCbl}$  and AS (see below) and is also a product of AS decomposition (eq 2). Furthermore, a control experiment showed that commercial AS can contain ~5% nitrite impurity (Experimental Section). Importantly, the UV-vis spectra of  $\text{H}_2\text{OCbl}^+$  and  $\text{NO}_2\text{Cbl}$  are practically identical,<sup>60</sup> hence if  $\text{H}_2\text{OCbl}^+$  is partially converted to  $\text{NO}_2\text{Cbl}$  by reacting with nitrite as the reaction proceeds prior to being converted to NOCbl, little or no change in the isosbestic points would be observed. Values of  $K = 2.2 \times 10^5 \text{ M}^{-1}$  and  $k_1 = 99.82 \text{ M}^{-1} \text{ s}^{-1}$  at 25 °C have been reported for the formation of  $\text{NO}_2\text{Cbl}$  from the reaction of  $\text{H}_2\text{OCbl}^+$  with nitrite, eq 3.<sup>59,61,62</sup>




Control experiments showed that the rate of the reaction between  $\text{H}_2\text{OCbl}^+/\text{HOCl}$  and nitrite is still rapid even in alkaline solution despite the reaction proceeding through  $\text{H}_2\text{OCbl}^+$  ( $\text{p}K_a(\text{H}_2\text{OCbl}^+) = 7.8^{55}$ ). Specifically, Figures S2a and S2b in the Supporting Information show that  $k_{\text{app}} = 5.95 \pm 0.04 \text{ M}^{-1} \text{ s}^{-1}$  and  $k_{\text{app}} = 41.5 \pm 0.2 \text{ M}^{-1} \text{ s}^{-1}$  at pH 9.80 and 8.80, respectively, where  $k_{\text{app}}$  is the apparent rate constant for the reaction between  $\text{H}_2\text{OCbl}^+/\text{HOCl}$  and nitrite (25.0 °C, 0.30 M CAPS or TAPS buffer,  $I = 1.0 \text{ M}$  ( $\text{NaCF}_3\text{SO}_3$ )). Note that the hydroxo ligand of  $\text{HOCl}$  is inert to substitution.<sup>63</sup> It therefore is possible that  $\text{NO}_2\text{Cbl}$  is formed as an intermediate in the reaction. To determine whether forming  $\text{NO}_2\text{Cbl}$  would have an effect on the observed rate constant, the rate of the reaction between the authentic  $\text{NO}_2\text{Cbl}$  and AS was determined at pH 9.80 and 9.09 (25.0 °C, 0.30 M CHES buffer,  $I = 1.0 \text{ M}$  ( $\text{NaCF}_3\text{SO}_3$ )). Figure S3 in the Supporting Information shows the plots of absorbance at 356 nm versus time for the reaction of  $\text{HOCl}$  or  $\text{NO}_2\text{Cbl}$  ( $5.00 \times 10^{-5} \text{ M}$ ) with excess AS ( $5.00 \times 10^{-3} \text{ M}$ ) at pH 9.80 (25.0 °C, 0.30 M CHES buffer,  $I = 1.0 \text{ M}$  ( $\text{NaCF}_3\text{SO}_3$ )). The data fit well to first-order rate equations giving  $k_{\text{obs}} = (8.29 \pm 0.01) \times 10^{-3}$  and  $(5.79 \pm 0.04) \times 10^{-3} \text{ s}^{-1}$  for the reaction of AS with  $\text{HOCl}$  and  $\text{NO}_2\text{Cbl}$ , respectively. Substituting  $\text{HOCl}$  by  $\text{NO}_2\text{Cbl}$  still results in the complete formation of  $\text{NOCl}$ , but leads to a slight decrease in the rate of the reaction. It is likely that  $\text{NO}_2\text{Cbl}$  does not itself react directly with AS, but that  $\text{NO}_2\text{Cbl}$  is in equilibrium with  $\text{H}_2\text{OCbl}^+$  and the latter species reacts with AS to form  $\text{NOCl}$ . However, experiments were not carried out to probe this mechanism further.

It was, therefore, likely that nitrite interferes with the reaction of interest. Given that  $\text{NO}_2^-$  is the non-cobalamin product of the reaction between  $\text{H}_2\text{OCbl}^+/\text{HOCl}$  and AS (see below), the maximum concentration of nitrite produced during the reaction is  $\sim[\text{AS}]$ . In order to minimize the amount of  $\text{NO}_2^-$  produced during the reaction of  $\text{H}_2\text{OCbl}^+/\text{HOCl}$  with AS, the kinetics of the reaction between  $\text{H}_2\text{OCbl}^+$  and AS were instead investigated with excess  $\text{H}_2\text{OCbl}^+$  (at least 7 times compared to the concentration of AS), rather than with AS in excess. Importantly, control experiments demonstrated that the addition of 1.0 mol equiv  $\text{NO}_2^-$  ( $5.00 \times 10^{-5} \text{ M}$ ) decreases the rate constant by only  $\sim 10\%$  for the reaction between AS ( $5.00 \times 10^{-5} \text{ M}$ ) and excess  $\text{H}_2\text{OCbl}^+$  ( $5.00 \times 10^{-4} \text{ M}$ ) at pH  $4.15 \pm 0.03$  ( $k_{\text{obs}} = (4.91 \pm 0.03) \times 10^{-2}$  and  $(4.40 \pm 0.02) \times 10^{-2} \text{ s}^{-1}$  in the absence and presence of  $5.00 \times 10^{-5} \text{ M}$   $\text{NaNO}_2$ , respectively), whereas there was no difference observed at pH  $7.20 \pm 0.03$  ( $k_{\text{obs}} = 6.78 \pm 0.03$  and  $6.86 \times 10^{-2} \text{ s}^{-1}$ ) in the absence and presence of  $5.00 \times 10^{-5} \text{ M}$   $\text{NaNO}_2$ , respectively). Hence since only  $\sim 0.14$  mol equiv  $\text{NO}_2^-$  (=7 times excess  $\text{H}_2\text{OCbl}^+/\text{HOCl}$ ) is produced in our experiments,  $\text{NO}_2^-$  production will not significantly alter the observed rate of the reaction. Control experiments also demonstrated that the rate constant is unaffected by the presence of the metal chelator DTPA ( $k_{\text{obs}} = (4.91 \pm 0.03) \times 10^{-2}$  and  $(4.94 \pm 0.05) \times 10^{-2} \text{ s}^{-1}$  in the absence and presence of 50.0  $\mu\text{M}$  DTPA, respectively, at pH  $4.15 \pm 0.03$ ), showing that free metal ions are not involved in the reaction.

It was established that spontaneous decomposition of AS is not significant on the time scale of the experiment. Typical UV-vis spectra as a function of time for the reaction of AS


( $6.50 \times 10^{-5} \text{ M}$ ) with excess  $\text{H}_2\text{OCbl}^+/\text{HOCl}$  ( $3.25 \times 10^{-4} \text{ M}$ ) at pH 5.00 (25.0 °C, 0.30 M acetate buffer,  $I = 1.0 \text{ M}$  ( $\text{NaCF}_3\text{SO}_3$ )) under anaerobic conditions are shown in Figure S4, Supporting Information. The isosbestic points were the same as those observed for the conversion of  $\text{H}_2\text{OCbl}^+/\text{HOCl}$  to  $\text{NOCl}$  at pH 7.40 within the limitations of the small absorbance changes due to an incomplete reaction.<sup>56</sup> Kinetic data were collected at a range of  $\text{H}_2\text{OCbl}^+$  concentrations in order to determine the apparent second-order rate constant for the reaction at pH 5.00, maintaining a  $\text{H}_2\text{OCbl}^+:\text{AS}$  ratio of  $\sim 7:1$ . The data are summarized in Figure 4, which shows a plot



**Figure 4.** Plot of  $k_{\text{obs}}$  versus concentration of  $\text{H}_2\text{OCbl}^+$  at pH  $5.00 \pm 0.03$  (25.0 °C, 0.30 M acetate buffer,  $I = 1.0 \text{ M}$  ( $\text{NaCF}_3\text{SO}_3$ )). The data have been fitted to a straight line passing through the origin, giving a slope ( $k_{\text{app}} = 129.4 \pm 1.4 \text{ M}^{-1} \text{ s}^{-1}$ ).

of  $k_{\text{obs}}$  versus the concentration of  $\text{H}_2\text{OCbl}^+$  at pH  $5.00 \pm 0.03$ . The data fit to a straight line passing through the origin, giving a slope (second-order rate constant,  $k_{\text{app}} = 129.4 \pm 1.4 \text{ M}^{-1} \text{ s}^{-1}$ ).

Values of  $k_{\text{app}}$  were similarly determined in the pH range 4.15–9.90 (individual plots given in Figures S5–S11 in the Supporting Information), and the results are summarized in Figure 5. Plots of  $k_{\text{obs}}$  versus  $[\text{H}_2\text{OCbl}^+/\text{HOCl}]$  at pH 9.05, 9.40, and 9.90 had a small but significant y-intercept. This arises because under these pH conditions decomposition of AS is no




**Figure 5.** Plot of  $k_{\text{app}}$  versus pH for the reaction between  $\text{H}_2\text{OCbl}^+/\text{HOCl}$  and AS. The best fit of the data to eq 5 fixing  $\text{p}K_a(\text{H}_2\text{OCbl}^+) = 7.8$  and  $\text{p}K_a(\text{HN}_2\text{O}_3^-) = 9.48$  gives  $k_{\text{Cbl(III)}} = 122.6 \pm 5.3 \text{ M}^{-1} \text{ s}^{-1}$ .

longer insignificant compared to the reaction of interest. In this case the data were fitted to eq 4.

$$k_{\text{obs}} = k_{\text{app}}[\text{H}_2\text{OCbl}^+/\text{HOCbl}] + k_t \quad (4)$$

The stoichiometry of the reaction between  $\text{H}_2\text{OCbl}^+$  and AS was determined using UV–vis spectroscopy. UV–vis spectra were taken of equilibrated solutions of  $\text{H}_2\text{OCbl}^+$  ( $5.00 \times 10^{-5}$  M) and AS (0, 0.25–2.5 equiv) at pH 6.00. The isosbestic points obtained (Figure 6) are in agreement with the literature




**Figure 6.** UV–vis spectra obtained from titration for the reaction of  $\text{H}_2\text{OCbl}^+$  ( $5.00 \times 10^{-5}$  M) with AS (0, 0.25–2.5 mol equiv) at pH 6.00 under anaerobic conditions. Isosbestic points occur at 337, 365, and 490 nm. **Inset:** Plot of absorbance at 351 nm versus mol equiv of AS at pH 6.00 (0.30 M MES buffer).

values for the conversion of  $\text{H}_2\text{OCbl}^+/\text{HOCbl}$  to  $\text{NOCbl}$  at pH 7.4.<sup>56</sup> The plot of absorbance at 351 nm versus mol equiv of AS (inset to Figure 6) gives a stoichiometry of 1:1  $\text{H}_2\text{OCbl}^+:\text{AS}$  at pH 6.00. The stoichiometry and the Cbl product(s) were also determined at pD 9.86 using  $^1\text{H}$  NMR spectroscopy.  $\text{HOCbl}$  is completely converted to  $\text{NOCbl}$  upon reacting  $\text{HOCbl}$  with 1.1 mol equiv AS (pD 9.86, 24 °C, 0.30 M CHES buffer, Supporting Information, Figure S12).

The non-cobalamin product was also identified. Nitrite was shown to be the non-cobalamin product using the Griess assay.<sup>54</sup> A calibration curve of absorbance versus nitrite concentration was generated (see the Experimental Section for details). From the Griess test, it was found that 0.79 equiv  $\text{NO}_2^-$  was produced in the reaction between  $\text{HOCbl}$  ( $1.00 \times 10^{-4}$  M) and AS (1.0 mol equiv) at pH 9.80, using a calibration curve of absorbance versus nitrite concentration (Figure S13, Supporting Information). Interestingly, others have reported significantly less  $\text{NO}_2^-$  formation than expected from AS decomposition based on the stoichiometry of the reaction.<sup>64–66</sup> This has been attributed to N–N bond homolysis (resulting in formation of  $2 \cdot\text{NO}$  and  $\text{H}_2\text{O}$ ) in addition to N–N heterolysis of  $\text{HN}_2\text{O}_3^-$  ( $\text{HNO}$  and  $\text{NO}_2^-$  formed).<sup>38</sup> We have noticed in our studies with AS that a slight excess of AS ( $\sim 1.1$  mol equiv) is always required for reactions to proceed to completion.

From Figure 5 it is clear that the reaction of  $\text{HOCbl}/\text{H}_2\text{OCbl}^+$  with AS becomes faster with decreasing pH, becoming pH independent at  $\text{pH} \leq 6$ . Scheme 1 gives the proposed reaction pathways involved for  $\text{pH} \leq 9.90$ . The

### Scheme 1. Proposed Mechanism for the Reaction of $\text{H}_2\text{OCbl}^+/\text{HOCbl}$ with AS at $\text{pH} \leq 9.90$




equilibrium between  $\text{H}_2\text{OCbl}^+$ , nitrite, and  $\text{NO}_2\text{Cbl}$  is shown for completeness; however, the effect of this equilibrium under the conditions of the kinetic experiments is insignificant. It is likely that rapid substitution of the aqua ligand of  $\text{H}_2\text{OCbl}^+$  by AS occurs prior to AS undergoing N–N bond cleavage. It is well established that  $\beta$ -axial ligand substitution reactions for cobalamins occur via a dissociative interchange mechanism.<sup>67</sup> Furthermore, only  $\text{H}_2\text{OCbl}^+$ , not  $\text{HOCbl}$ , undergoes these reactions, since  $\text{HOCbl}$  is inert to  $\beta$ -axial ligand substitution,<sup>58</sup> hence in Scheme 1 only  $\text{H}_2\text{OCbl}^+$  ( $\text{p}K_a = 7.8^{55}$ ), not  $\text{HOCbl}$ , reacts with AS. The corresponding rate equation is given in eq 5.

$$k_{\text{app}} = \frac{kK[\text{H}^+]^2}{([\text{H}^+] + K_a(\text{H}_2\text{OCbl}^+)) + ([\text{H}^+] + K_a(\text{HN}_2\text{O}_3^-))} \quad (5)$$

The best fit of the data in Figure 5 to eq 5 with  $\text{p}K_a(\text{H}_2\text{OCbl}^+) = 7.8^{55}$  and  $\text{p}K_a(\text{HN}_2\text{O}_3^-) = 9.48$  gives  $kK = 123 \pm 5 \text{ M}^{-1} \text{ s}^{-1}$ . This value is of a comparable magnitude to values reported for the reactions of  $\text{H}_2\text{OCbl}^+$  with a wide variety of ligands.<sup>62,68</sup> By fitting the data to this eq 5, we assume that the reaction between  $\text{H}_2\text{OCbl}^+$  and  $\text{N}_2\text{O}_3^{2-}$  is not important under the pH conditions of our study. Indeed, if the data are instead fitted to a model in which both  $\text{HN}_2\text{O}_3^-$  and  $\text{N}_2\text{O}_3^{2-}$  react with  $\text{H}_2\text{OCbl}^+$ ,  $kK$  is the same within experimental error as expected ( $kK = 121 \pm 5 \text{ M}^{-1} \text{ s}^{-1}$ ; see Supporting Information, Figure S14 and Scheme S1), since practically no Cbl remains in the reactive  $\text{H}_2\text{OCbl}^+$  form at pH conditions where the concentration of  $\text{N}_2\text{O}_3^{2-}$  is significant ( $\text{pH} > 9$ ).

Two mechanisms have been proposed for the reaction of AS with porphyrins and heme proteins. AS reacts directly with  $\text{Mn}^{\text{III}}$  porphyrins ( $\text{Mn}^{\text{III}}\text{TPEPy}^{69}$ ) and  $\text{Fe}^{\text{III}}$  porphyrins ( $\text{Fe}^{\text{III}}\text{TPEPy}^{70}$ ) to form an AS-bound intermediate (rate-determining step, RDS) followed by rapid decomposition of intermediate to form the final nitrosyl products,  $\text{Mn}^{\text{II}}\text{TPEPy}(\text{NO})$  and  $\text{Fe}^{\text{II}}\text{TPEPy}(\text{NO})$ , respectively. Alternatively, the reaction may occur via a mechanism in which  $\text{HNO}$ , not AS, reacts. This mechanism has been proposed for the reaction of AS with  $\text{Mn}^{\text{III}}\text{TPPS}$ ,<sup>69</sup> microperoxidase-11 ( $\text{Fe}^{\text{III}}\text{MP11}$ ),<sup>70</sup>  $\text{Fe}^{\text{III}}(\text{TPPS})$ ,<sup>71</sup> methemoglobin (methHb) and metmyoglobin (metMb),<sup>72,73</sup> and horseradish peroxidase,<sup>74</sup> to form the corresponding  $\text{Mn}^{\text{II}}(\text{NO})$  and  $\text{Fe}^{\text{II}}(\text{NO})$  complexes. Indeed, evidence for this latter mechanism occurring in our system was obtained at pH 10.80 and 11.40 (see below).

**Studies on the Reaction between  $\text{HOCbl}$  and AS at High pH Conditions (pH 10.80, 11.40).** Figure 7a gives a



**Figure 7.** (a) Plot of absorbance at 356 nm versus time for reaction of HOcbl ( $1.00 \times 10^{-4}$  M) with excess AS ( $1.00 \times 10^{-2}$  M) at pH 10.80 (25.0 °C, 0.30 M CAPS buffer,  $I = 1.0$  M ( $\text{NaCF}_3\text{SO}_3$ )). The linear dependence suggests that the reaction is independent of HOcbl concentration. (b) Plot of absorbance at 356 nm versus time for reaction of HOcbl ( $1.00 \times 10^{-4}$  M) with 1.0 equiv AS at pH 10.80 (25.0 °C, 0.30 M CAPS buffer,  $I = 1.0$  M ( $\text{NaCF}_3\text{SO}_3$ )). The best fit of the data to a first-order reaction gave  $k_{\text{obs}} = (7.45 \pm 0.01) \times 10^{-5} \text{ s}^{-1}$ .

plot of absorbance versus time for the reaction of HOcbl ( $1.00 \times 10^{-4}$  M) with excess AS ( $1.00 \times 10^{-2}$  M) at pH 10.80 (25.0 °C, 0.30 M CAPS buffer,  $I = 1.0$  M ( $\text{NaCF}_3\text{SO}_3$ )). An almost linear dependence is observed, with HOcbl reacting with AS to again give NOcbl. A similar result was obtained at  $5.00 \times 10^{-4}$  M HOcbl. The linear dependence suggests that the reaction is independent of HOcbl concentration (i.e., zero order in [HOcbl]), and that the decomposition of AS is instead the rate-determining step at these pH conditions. Therefore, the same experiment was repeated using 1.0 mol equiv AS at pH 10.80 (0.30 M CAPS buffer,  $I = 1.0$  M,  $\text{NaCF}_3\text{SO}_3$ ). Under these conditions the absorbance at 356 nm versus time data fit well to a first-order rate equation, with an observed rate constant ( $k_{\text{obs}}$ ) of  $(7.45 \pm 0.01) \times 10^{-5} \text{ s}^{-1}$  (see Figure 7b). A similar rate constant was observed with 0.5 equiv AS at the same conditions ( $(8.00 \pm 0.05) \times 10^{-5} \text{ s}^{-1}$ ). Given that the reaction is stoichiometric (see below),  $k_{\text{obs}}$  would be expected to be independent of HOcbl for [HOcbl]:[AS]  $\geq 1.0$  (also see data simulations in the Supporting Information). The reaction of excess HOcbl ( $1.00 \times 10^{-4}$  M) with AS ( $1.40 \times 10^{-5}$  M) at pH 10.80 (25.0 °C, 0.30 M CAPS buffer,  $I = 1.0$  M,  $\text{NaCF}_3\text{SO}_3$ ) gives  $k_{\text{obs}} = (5.32 \pm 0.06) \times 10^{-5} \text{ s}^{-1}$ ; see Figure S15 in the Supporting Information. The observed rate constant for the spontaneous decomposition of AS at the same pH is  $(3.17 \pm 0.01) \times 10^{-5} \text{ s}^{-1}$ . The observed rate constants for the reaction between HOcbl and AS are therefore up to 2.5 times larger than one would expect based solely on a pathway involving rate-determining AS decomposition. Possible explanations for this discrepancy are that there is a small contribution to the observed reaction from the direct reaction (Scheme 1) or that weak association of HOcbl with AS in aqueous solution promotes AS decomposition.

The reaction between HOcbl and AS was also studied at pH 11.40 (0.40 M phosphate buffer), using 0.50 and 1.0 mol equiv AS. The best fit of the absorbance versus time data to a first-order equation gives  $k_{\text{obs}} = (1.10 \pm 0.01) \times 10^{-5}$  and  $(1.15 \pm 0.01) \times 10^{-5} \text{ s}^{-1}$ , respectively. At pH 11.40 the observed rate constant for the spontaneous decomposition of AS is  $(4.70 \pm 0.02) \times 10^{-6} \text{ s}^{-1}$ ; hence once again the rate of the reaction is slightly faster than one would expect based purely on AS decomposition.


Since the reaction of AS with HOcbl at higher pH conditions occurs via a different mechanism, the stoichiometry of the reaction of HOcbl with 0.55, 1.1, and 2.2 mol equiv AS was investigated by  $^1\text{H}$  NMR spectroscopy at pD 10.86 (0.50

M CAPS). A minimum of 1.1 mol equiv AS is required for the reaction to proceed to completion (Figure S16, Supporting Information), with NOcbl once again formed as the Cbl product. Reacting 0.50 mol equiv AS yielded approximately ~50% NOcbl and ~50% unreacted starting material (HOcbl). An attempt was also made to determine the stoichiometry of the reaction at pD 11.40. However, since the reaction at this pH condition is very slow and Cbls are not stable in alkaline solution,<sup>75,76</sup> considerable cobalamin decomposition to Co(II) corrinoid species was observed at this pH value.

Finally, the amount of  $\text{NO}_2^-$  produced at pH 10.80 was determined using the Griess assay<sup>54</sup> under strictly anaerobic conditions. From the calibration curve of absorbance versus nitrite concentration, the resulting absorbance corresponded to 0.81 and 0.78 mol equiv nitrite produced (see Experimental Section). Hence nitrite is the non-Cbl product.

The proposed major reaction pathway for the reaction between  $\text{H}_2\text{OCbl}^+/\text{HOcbl}$  and AS at pH  $\geq 10.80$  is summarized in Scheme 2. Decomposition of monoprotonated

#### Scheme 2. Proposed Major Mechanism for the Reaction of $\text{H}_2\text{OCbl}^+/\text{HOcbl}$ with AS at pH $\geq 10.80$



form of AS ( $\text{HN}_2\text{O}_3^-$ ;  $\text{p}K_a$  9.48) to give  $\text{HNO}/\text{NO}^-$  is the rate-limiting step of the reaction followed by fast substitution of aqua ligand of  $\text{H}_2\text{OCbl}^+$  by  $\text{HNO}/\text{NO}^-$  to form NOcbl. Since decomposition of  $\text{HN}_2\text{O}_3^-$  is rate-limiting, our experimental data do not allow us to determine whether  $^3\text{NO}^-$  in addition to  $^1\text{HNO}$  ( $\text{p}K_a(^1\text{HNO}) \sim 11.4$ ;<sup>23</sup>  $^1\text{HNO} \rightleftharpoons ^3\text{NO}^- + \text{H}^+$ ) reacts with  $\text{H}_2\text{OCbl}^+$  and/or HOcbl to form NOcbl. Furthermore, no information is obtained on the mechanism of this reaction.

## CONCLUSIONS

Kinetic and mechanistic studies have been carried out on the reaction of aquacobalamin with the HNO donor Angeli's salt using UV-vis and  $^1\text{H}$  NMR spectroscopies. At lower pH

conditions ( $\leq 9.90$ ),  $\text{H}_2\text{OCbl}^+$  reacts directly with AS with a 1:1 stoichiometry, giving  $\text{NOCl}$  and  $\text{NO}_2^-$ . A direct transfer of a nitroxyl group to the cobalt(III) center of  $\text{H}_2\text{OCbl}^+$  from the ligand was also observed for the reaction of  $\text{R}_2\text{N-NONOates}$  with  $\text{H}_2\text{OCbl}^+$ .<sup>77</sup> Under strongly alkaline conditions (pH  $\geq 10.80$ ) the rate-determining step instead involves AS decomposition to give  $\text{HNO/NO}^-$  and  $\text{NO}_2^-$  (RDS), with  $\text{HNO/NO}^-$  subsequently rapidly reacting with  $\text{H}_2\text{OCbl}^+$  to give  $\text{NOCl}$ .

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

Additional kinetic data,  $^1\text{H}$  NMR spectra, calibration plot for nitrite detection, and data simulations; Figures S1–S16 and Scheme S1. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [nbrasch@kent.edu](mailto:nbrasch@kent.edu).

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This material is based upon work supported by the U.S. National Science Foundation under CHE-0848397.

## ■ REFERENCES

- (1) Doctorovich, F.; Bikiel, D.; Pellegrino, J.; Suárez, S. A.; Larsen, A.; Martí, M. A. *Coord. Chem. Rev.* **2011**, *255*, 2764.
- (2) Basudhar, D.; Bharadwaj, G.; Cheng, R. Y.; Jain, S.; Shi, S.; Heinecke, J. L.; Holland, R. J.; Ridnour, L. A.; Caceres, V. M.; Spadari-Bratfisch, R. C.; Paolucci, N.; Velazquez-Martinez, C. A.; Wink, D. A.; Miranda, K. M. *J. Med. Chem.* **2013**, *56*, 7804.
- (3) Aizawa, K.; Nakagawa, H.; Matsuo, K.; Kawai, K.; Ieda, N.; Suzuki, T.; Miyata, N. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 2340.
- (4) Nakagawa, H. *J. Inorg. Biochem.* **2013**, *118*, 187.
- (5) Zhang, Y. *J. Inorg. Biochem.* **2013**, *118*, 191.
- (6) Fukuto, J. M.; Cisneros, C. J.; Kinkade, R. L. *J. Inorg. Biochem.* **2013**, *118*, 201.
- (7) Nakagawa, H. *Nitric Oxide* **2011**, *25*, 195.
- (8) Sirsalmath, K.; Suárez, S. A.; Bikiel, D. E.; Doctorovich, F. *J. Inorg. Biochem.* **2013**, *118*, 134.
- (9) Irvine, J. C.; Ritchie, R. H.; Favaloro, J. L.; Andrews, K. L.; Widdop, R. E.; Kemp-Harper, B. K. *Trends Pharmacol. Sci.* **2008**, *29*, 601.
- (10) Fukuto, J. M.; Dutton, A. S.; Houk, K. N. *ChemBioChem* **2005**, *6*, 612.
- (11) Miranda, K. M.; Espey, M. G.; Yamada, K.; Krishna, M.; Ludwick, N.; Kim, S.; Jourdeuil, D.; Grisham, M. B.; Feelisch, M.; Fukuto, J. M.; Wink, D. A. *J. Biol. Chem.* **2001**, *276*, 1720.
- (12) Sharpe, M. A.; Cooper, C. E. *Biochem. J.* **1998**, *332*, 9.
- (13) Murphy, M. E.; Sies, H. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 10860.
- (14) Filipovic, M. R.; Miljkovic, J. L.; Nauser, T.; Royzen, M.; Klos, K.; Shubina, T.; Koppenol, W. H.; Lippard, S. J.; Ivanović-Burmazović, I. *J. Am. Chem. Soc.* **2012**, *134*, 12016.
- (15) Naughton, P.; Foresti, R.; Bains, S. K.; Hoque, M.; Green, C. J.; Motterlini, R. *J. Biol. Chem.* **2002**, *277*, 40666.
- (16) Kawai, K.; Ieda, N.; Aizawa, K.; Suzuki, T.; Miyata, N.; Nakagawa, H. *J. Am. Chem. Soc.* **2013**, *135*, 12690.
- (17) Johnson, G. M.; Chozinski, T. J.; Salmon, D. J.; Moghaddam, A. D.; Chen, H. C.; Miranda, K. M. *Free Radical Biol. Med.* **2013**, *63*, 476.
- (18) Reisz, J. A.; Zink, C. N.; King, S. B. *J. Am. Chem. Soc.* **2011**, *133*, 11675.
- (19) Cline, M. R.; Toscano, J. P. *J. Phys. Org. Chem.* **2011**, *24*, 993.

- (20) Zhou, Y.; Liu, K.; Li, J.-Y.; Fang, Y.; Zhao, T.-C.; Yao, C. *Org. Lett.* **2011**, *13*, 1290.
- (21) Suárez, S. A.; Bikiel, D. E.; Wetzler, D. E.; Martí, M. A.; Doctorovich, F. *Anal. Chem.* **2013**, *85*, 10262.
- (22) Bartberger, M. D.; Liu, W.; Ford, E.; Miranda, K. M.; Switzer, C.; Fukuto, J. M.; Farmer, P. J.; Wink, D. A.; Houk, K. N. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 10958.
- (23) Shafirovich, V.; Lymar, S. V. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 7340.
- (24) Reisz, J. A.; Bechtold, E.; King, S. B. *Dalton Trans.* **2010**, *39*, 5203.
- (25) Miranda, K. M.; Paolucci, N.; Katori, T.; Thomas, D. D.; Ford, E.; Bartberger, M. D.; Espey, M. G.; Kass, D. A.; Feelisch, M.; Fukuto, J. M.; Wink, D. A. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 9196.
- (26) Fukuto, J. M.; Bianco, C. L.; Chavez, T. A. *Free Radical Biol. Med.* **2009**, *47*, 1318.
- (27) Jackson, M. I.; Han, T. H.; Serbulea, L.; Dutton, A.; Ford, E.; Miranda, K. M.; Houk, K. N.; Wink, D. A.; Fukuto, J. M. *Free Radical Biol. Med.* **2009**, *47*, 1130.
- (28) Shoeman, D. W.; Shirota, F. N.; DeMaster, E. G.; Nagasawa, H. *T. Alcohol* **2000**, *20*, 55.
- (29) Fukuto, J. M.; Bartberger, M. D.; Dutton, A. S.; Paolucci, N.; Wink, D. A.; Houk, K. N. *Chem. Res. Toxicol.* **2005**, *18*, 790.
- (30) Pagliaro, P.; Mancardi, D.; Rastaldo, R.; Penna, C.; Gattullo, D.; Miranda, K. M.; Feelisch, M.; Wink, D. A.; Kass, D. A.; Paolucci, N. *Free Radical Biol. Med.* **2003**, *34*, 33.
- (31) Paolucci, N.; Jackson, M. I.; Lopez, B. E.; Miranda, K.; Tocchetti, C. G.; Wink, D. A.; Hobbs, A. J.; Fukuto, J. M. *Pharmacol. Ther.* **2007**, *113*, 442.
- (32) Guthrie, D. A.; Kim, N. Y.; Siegler, M. A.; Moore, C. D.; Toscano, J. P. *J. Am. Chem. Soc.* **2012**, *134*, 1962.
- (33) Huang, Z.; Kaur, J.; Bhardwaj, A.; Alsaleh, N.; Reisz, J. A.; DuMond, J. F.; King, S. B.; Seubert, J. M.; Zhang, Y.; Knaus, E. E. *J. Med. Chem.* **2012**, *55*, 10262.
- (34) Sutton, A. D.; Williamson, M.; Weismiller, H.; Toscano, J. P. *Org. Lett.* **2011**, *14*, 472.
- (35) Angeli, A.; Angelico, F. *Gazz. Chim. Ital.* **1903**, *33*, 245.
- (36) Fukuto, J. M.; Jackson, M. I.; Kaludercic, N.; Paolucci, N. In *Methods in Enzymology*; Enrique, C., Lester, P., Eds.; Academic Press: Waltham, MA, 2008; Vol. 440, p 411.
- (37) Sturrock, P. E.; Ray, J. D.; McDowell, J.; Hunt, H. R. *Inorg. Chem.* **1963**, *2*, 649.
- (38) Dutton, A. S.; Fukuto, J. M.; Houk, K. N. *J. Am. Chem. Soc.* **2004**, *126*, 3795.
- (39) Keck, B.; Munder, M.; Renz, P. *Arch. Microbiol.* **1998**, *171*, 66.
- (40) Banerjee, R. *Chemistry and Biochemistry of B<sub>12</sub>*; Wiley: New York, 1999.
- (41) Carmel, R.; Jacobsen, D. W. *Homocysteine in Health and Disease*; Cambridge University Press: Cambridge U.K., 2001.
- (42) Koyyalamudi, S. R.; Jeong, S.-C.; Cho, K. Y.; Pang, G. J. *Agric. Food Chem.* **2009**, *57*, 6327.
- (43) Goodrich, L. E.; Roy, S.; Alp, E. E.; Zhao, J.; Hu, M. Y.; Lehnert, N. *Inorg. Chem.* **2013**, *52*, 7766.
- (44) Doctorovich, F.; Bikiel, D. E.; Pellegrino, J.; Suarez, S. A.; Martí, M. A. *Adv. Inorg. Chem.* **2012**, *64*, 97.
- (45) Doctorovich, F.; Bikiel, D.; Pellegrino, J.; Suárez, S. A.; Martí, M. A. *J. Porphyr. Phthalocya.* **2010**, *14*, 1012.
- (46) Suárez, S. A.; Fonticelli, M. H.; Rubert, A. A.; de la Llave, E.; Scherlis, D.; Salvarezza, R. C.; Martí, M. A.; Doctorovich, F. *Inorg. Chem.* **2010**, *49*, 6955.
- (47) Brouwer, M.; Chamulitrat, W.; Ferruzzi, G.; Sauls, D. L.; Weinberg, J. B. *Blood* **1996**, *88*, 1857.
- (48) Schubert, R.; Krien, U.; Wulfsen, I.; Schiemann, D.; Lehmann, G.; Ulfög, N.; Veh, R. W.; Schwarz, J. R.; Gago, H. *Hypertension* **2004**, *43*, 891.
- (49) Jiang, F.; Li, C. G.; Rand, M. J. *Eur. J. Pharmacol.* **1997**, *340*, 181.
- (50) Weil, M.; Abeles, R.; Nachmany, A.; Gold, V.; Michael, E. *Cell Death Differ.* **2004**, *11*, 361.

- (51) Sampaio, A. L.; Dalli, J.; Brancaleone, V.; D'Acquisto, F.; Perretti, M.; Wheatley, C. *Mediators Inflammation* **2013**, 2013, 741804.
- (52) Barker, H. A.; Smyth, R. D.; Weissbach, H.; Toohey, J. I.; Ladd, J. N.; Volcani, B. E. *J. Biol. Chem.* **1960**, 235, 480.
- (53) Cavallini, D.; Scandurra, R. In *Methods in Enzymology*; Donald, B. M., Lemuel, D. W., Eds.; Academic Press: Waltham, MA, 1971; Vol. 18, Part C, p 3.
- (54) Green, L. C.; Wagner, D. A.; Glogowski, J.; Skipper, P. L.; Wishnok, J. S.; Tannenbaum, S. R. *Anal. Biochem.* **1982**, 126, 131.
- (55) Suarez-Moreira, E.; Yun, J.; Birch, C. S.; Williams, J. H. H.; McCaddon, A.; Brasch, N. E. *J. Am. Chem. Soc.* **2009**, 131, 15078.
- (56) Zheng, D.; Birke, R. L. *J. Am. Chem. Soc.* **2001**, 123, 4637.
- (57) Bonner, F. T.; Ravid, B. *Inorg. Chem.* **1975**, 14, 558.
- (58) Huges, M. N.; Wimbledon, P. E. *J. Chem. Soc., Dalton Trans.* **1976**, 703.
- (59) Firth, R. A.; Hill, H. A. O.; Pratt, J. M.; Thorp, R. G.; Williams, R. J. P. *J. Chem. Soc., A* **1969**, 381.
- (60) Suarez-Moreira, E.; Hannibal, L.; Smith, C. A.; Chavez, R. A.; Jacobsen, D. W.; Brasch, N. E. *Dalton Trans.* **2006**, 5269.
- (61) Knapton, L.; Marques, H. M. *Dalton Trans.* **2005**, 889.
- (62) Marques, H. M.; Knapton, L. *J. Chem. Soc., Dalton Trans.* **1997**, 3827.
- (63) Xia, L.; Cregan, A. G.; Berben, L. A.; Brasch, N. E. *Inorg. Chem.* **2004**, 43, 6848.
- (64) Reif, A.; Zecca, L.; Riederer, P.; Feelisch, M.; Schmidt, H. H. H. *W. Free Radical Biol. Med.* **2001**, 30, 803.
- (65) Cline, M. R.; Tu, C.; Silverman, D. N.; Toscano, J. P. *Free Radical Biol. Med.* **2011**, 50, 1274.
- (66) Zeller, A.; Wenzl, M. V.; Beretta, M.; Stessel, H.; Russwurm, M.; Koesling, D.; Schmidt, K.; Mayer, B. *Mol. Pharmacol.* **2009**, 76, 1115.
- (67) Hamza, M. S. A.; Dücker-Benfer, C.; van Eldik, R. *Inorg. Chem.* **2000**, 39, 3777.
- (68) Meier, M.; van Eldik, R. *Inorg. Chem.* **1993**, 32, 2635.
- (69) Martí, M. A.; Bari, S. E.; Estrin, D. A.; Doctorovich, F. *J. Am. Chem. Soc.* **2005**, 127, 4680.
- (70) Suárez, S. A.; Martí, M. A.; De Biase, P. M.; Estrin, D. A.; Bari, S. E.; Doctorovich, F. *Polyhedron* **2007**, 26, 4673.
- (71) Bari, S. E.; Martí, M. A.; Amorebieta, V. T.; Estrin, D. A.; Doctorovich, F. *J. Am. Chem. Soc.* **2003**, 125, 15272.
- (72) Bazylinski, D. A.; Hollocher, T. C. *J. Am. Chem. Soc.* **1985**, 107, 7982.
- (73) Doyle, M. P.; Mahapatro, S. N.; Broene, R. D.; Guy, J. K. *J. Am. Chem. Soc.* **1988**, 110, 593.
- (74) Huang, J.; Sommers, E. M.; Kim-Shapiro, D. B.; King, S. B. *J. Am. Chem. Soc.* **2002**, 124, 3473.
- (75) Yamada, R.; Kato, T.; Shimizu, S.; Fukui, S. *Biochim. Biophys. Acta* **1966**, 117, 13.
- (76) Lee, L.-P.; Schrauzer, G. N. *J. Am. Chem. Soc.* **1968**, 90, 5274.
- (77) Hassanin, H. A.; Hannibal, L.; Jacobsen, D. W.; El-Shahat, M. F.; Hamza, M. S. A.; Brasch, N. E. *Angew. Chem., Int. Ed.* **2009**, 48, 8909.