

room-temperature magnetic moment and the exchange integral for values of $\mu_{\text{eff}}(\text{RT})$ in the range 0.3–1.6 μ_{B} and $-2J$ in the range 190–1000 cm^{-1} . However, if one includes a larger selection of spin-coupled complexes involving a variety of ligand bridges and covering an extended range of $-2J$ and $\mu_{\text{eff}}(\text{RT})$ it becomes apparent that the relationship is not linear. Figure 1 contains $-2J$ vs. $\mu_{\text{eff}}(\text{RT})$ data for a varied selection of binuclear copper(II) complexes involving unrelated ligands, which have been synthesized and studied in a variety of different laboratories, where variable-temperature magnetic measurements have been carried out on a variety of different susceptometers with room-temperature magnetic moments quoted at different room temperatures.

Substitution of the Van Vleck equation (eq 1) into the expression (eq 3) (TIP, temperature-independent paramagnetism,

$$\mu_{\text{eff}} = 2.828(\chi_{\text{M}}T)^{1/2} \quad (3)$$

not included in the corrected molar susceptibility term) for the effective magnetic moment gives (eq 4) an expression, in which

$$-2J = kT \ln \left(\frac{3.0003g^2}{\mu_{\text{eff}}^2} - 3 \right) \quad (4)$$

the exchange integral ($-2J$) and the effective magnetic moment at temperature T are variables ($N\beta^2/3k = 0.12505$). A nonlinear regression analysis of the 36 data (Figure 1) using this two-parameter expression (eq 4) and using a weighting scheme in between a constant and proportional standard deviation¹⁶ gave the theoretical line (Figure 1) and a two-parameter fit of $g = 2.198 \pm 0.008$ and $T = 295.6 \pm 4.5$ K.

The very reasonable fit of the data with a sensible mean g value and mean room-temperature indicates, quite clearly, that realistic estimates of exchange integrals can be obtained for antiferromagnetically coupled binuclear copper(II) complexes by the use of this graphical fit or by substituting known values of g , T , and χ_{M} into the Van Vleck equation itself. The errors from this estimation are not likely to be unreasonable, and, in fact, for very strongly coupled systems, where variable temperature susceptibility measurements may involve large errors because of the weak paramagnetism involved, especially at low temperatures, measurements of $-2J$ values are likely to provide very reasonable results.

The upsurge of interest in spin-coupled copper compounds, which are relevant as models for spin-coupled binuclear copper protein active sites, has led to the synthesis and study of a large number of binuclear copper complexes, many of which have not been investigated by variable-temperature susceptibility studies because of the lack of suitable equipment. However, since most chemistry departments have room-temperature susceptibility equipment, which can be purchased or constructed at a modest cost, the results of room-temperature magnetic moment measurements can be used to provide reasonable estimates of exchange integrals. Of the simple techniques suitable for such measurements the Faraday method, using a sensitive microbalance with resolution at the microgram level, would be favored, especially for weakly paramagnetic, strongly coupled complexes.

Acknowledgment. We thank the Natural Sciences and Engineering Research Council of Canada for financial support for this study.

- (8) Dooley, D. M.; Scott, R. A.; Ellinghaus, J.; Solomon, E. I.; Gray, H. B. *Proc. Natl. Acad. Sci. U.S.A.* **1978**, *75*, 3019.
- (9) Morgenstern-Badarau, I.; Cocco, D.; Desideri, A.; Rotilio, G.; Jardonov, J.; Dupre, N. *J. Am. Chem. Soc.* **1986**, *108*, 300.
- (10) Thompson, L. K.; Hanson, A. W.; Ramaswamy, B. S. *Inorg. Chem.* **1984**, *23*, 2459.
- (11) Thompson, L. K.; Hartstock, F. W.; Robichaud, P.; Hanson, A. W. *Can. J. Chem.* **1984**, *62*, 2755.
- (12) Mandal, S. K.; Woon, T. C.; Thompson, L. K.; Newlands, M. J.; Gabe, E. J. *Aust. J. Chem.*, in press.
- (13) Thompson, L. K.; Woon, T. C.; Murphy, D. B.; Gabe, E. J.; Lee, F. L.; Le Page, Y. *Inorg. Chem.* **1985**, *24*, 4719.
- (14) Robichaud, P.; Thompson, L. K. *Inorg. Chim. Acta* **1984**, *85*, 137.
- (15) Crawford, V. H.; Richardson, H. W.; Wasson, J. R.; Hodgson, D. J.; Hatfield, W. E. *Inorg. Chem.* **1976**, *15*, 2107.
- (16) Duggleby, R. G. *Anal. Biochem.* **1981**, *110*, 9.

Contribution No. 7342 from the Department of Chemistry, A. A. Noyes Laboratory of Chemical Physics, California Institute of Technology, Pasadena, California 91125

Crystal Structure of Bis(*p*-toluidine)bis(acetonitrile)tetraiodotetracopper: Correction of Space Group

William P. Schaefer

Received December 23, 1985

The crystal structure of the tetrameric copper cluster bis(*p*-toluidine)bis(acetonitrile)tetraiodotetracopper has recently been described.¹ It was reported in the triclinic space group $P\bar{1}$ with two formula weights in the unit cell. In fact it should be formulated in the monoclinic space group $C2/c$, with $Z = 4$.

The reported cell dimensions are $a = 14.274$ (5) Å, $b = 7.587$ (2) Å, $c = 14.239$ (5) Å, $\alpha = 88.61$ (2)°, $\beta = 110.07$ (2)°, and $\gamma = 88.70$ (2)°. The vectors $\mathbf{c} + \mathbf{a}$, $\mathbf{c} - \mathbf{a}$ and $-\mathbf{b}$ describe an effectively monoclinic cell with $a = 16.340$ Å, $b = 23.366$ Å, $c = 7.587$ Å, $\alpha = 90.05$, $\beta = 92.35$ °, and $\gamma = 90.15$ °. The atomic coordinates, transformed according to the relations $x_{\text{m}} = 1/2x_{\text{t}}$, $+1/2z_{\text{t}} + 1/4$, $y_{\text{m}} = -1/2x_{\text{t}} + 1/2z_{\text{t}} - 1/4$, and $z_{\text{m}} = -y_{\text{t}}$, yield values that are compatible with the space group $C2/c$ within the reported esd's (Table I, supplementary material). Since the coordinate shifts necessary to achieve the higher symmetry are no greater than the esd's, the change in space group does not change the description of the structure, other than to introduce an exact crystallographic twofold axis in place of the approximate one. Further confirmation of the correctness of this description is obtained from the structure factors, which show the appropriate absences (triclinic hkh , $k = 2n + 1$)³ for space group $C2/c$.

Acknowledgment. I thank Richard E. Marsh for his interest and stimulation. This work was supported in part by the Exxon Educational Foundation.

Supplementary Material Available: Table I, positional parameters for $\text{Cu}_4\text{I}_4\text{C}_{18}\text{H}_{24}\text{N}_4$ in $C2/c$ (1 page). Ordering information is given on any current masthead page.

- (1) Rath, N. P.; Holt, E. M.; Tanimura, K. *Inorg. Chem.* **1985**, *24*, 3934–3938.
- (2) The translational components of $1/4$ are needed to put the origin at the conventional center of symmetry.
- (3) Of the 250 reflections of this type listed in the supplementary data, 119 had k odd and 118 of these were marked with an asterisk, presumably indicating unobserved. The sole exception, 14,1,14, had $F_o = 208$ where some unobserved reflections had $F_o = 164$ *. Of the 131 such reflections with k even, 59 were not marked and were presumably considered observed.

Contribution from the Chemistry Departments,
State University of New York at Buffalo,
Buffalo, New York 14214,
and Gettysburg College, Gettysburg, Pennsylvania 17325

Synthesis and Characterization of a Ruthenium(II) Clathrochelate Complex

J. G. Muller,^{1a} J. J. Grzybowski,^{*1b} and K. J. Takeuchi^{*1a}

Received November 14, 1985

Metal clathrochelates consist of a metal ion encapsulated by a bicyclic macrocyclic ligand.² These complexes often possess greater thermodynamic and kinetic stability than that found in analogous open-chain complexes, because of their greatly reduced

- (1) (a) State University of New York at Buffalo. (b) Gettysburg College.
- (2) Busch, D. H. *Rec. Chem. Prog.* **1964**, *25*, 107.