

# Notes

## Hydrolysis of $\text{Cr}_2\text{O}_7^{2-}$ . Nucleophilic Catalysis by TRIS and Rate Laws for the Formation and Decay of a (TRIS) $\text{CrO}_3$ Intermediate


Nicola E. Brasch, David A. Buckingham,\* and Charles R. Clark

Department of Chemistry, University of Otago, PO Box 56, Dunedin, New Zealand

Received November 30, 1993

### Introduction


As part of a  $^{17}\text{O}$  study of oxygen exchange into  $\text{HCrO}_4^-$  and  $\text{Cr}_2\text{O}_7^{2-}$  in aqueous solution, we have had cause to re-examine the buffer-catalyzed hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$ , eq 1. The classic work in



this area by Perlmutter-Hayman and associates in the 1960's<sup>1,2</sup> established both general acid (HOAc and chloroacetic acid) and base (some 36 oxygen and nitrogen bases were examined) pathways in addition to  $\text{H}^+$ - and  $\text{OH}^-$ -catalyzed and spontaneous reactions, eq 2, but although Perlmutter-Hayman suggested a


$$k_{\text{obs}} = k_0 + k_{\text{H}}[\text{H}^+] + k_{\text{OH}}[\text{OH}^-] + k_{\text{AH}}[\text{AH}] + k_{\text{B}}[\text{B}] \quad (2)$$

direct nucleophilic role for B, eq 3, it was never proven. In other



reports large negative entropies<sup>3</sup> and volumes<sup>4</sup> of activation have been used to support a nucleophilic role for  $\text{NH}_3$  and 2,6-lutidine, and the reverse reaction of  $\text{HCrO}_4^-$  with oxyanions to give  $\text{BCrO}_3$  products is well established.<sup>5–8</sup> This suggests that such intermediates are thermodynamically quite stable. However, only for  $\text{B} = \text{S}_2\text{O}_3^{2-}$  has a mechanism involving nucleophilic catalysis been established.<sup>8</sup>

In our experiments we have found the reaction of  $\text{Cr}_2\text{O}_7^{2-}$  with TRIS to be spectrophotometrically biphasic, thus requiring the formation of an intermediate. TRIS is traditionally regarded as a nonnucleophilic buffer so that this observation is of particular interest. Furthermore no reduction of Cr(VI) was evident in the



**Figure 1.** Variation of absorbance with time for the hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$  in the presence of TRIS buffer ( $[\text{TRIS}]_{\text{T}} = 0.050 \text{ M}$ ,  $[\text{Cr}]_{\text{T}} = 4 \times 10^{-3} \text{ M}$ ,  $\lambda = 450 \text{ nm}$ ,  $\text{pH} = 7.84$ ,  $25^\circ\text{C}$ ). The data have been fitted to two consecutive first-order reactions with rate constants  $k_{\text{fast}} = 0.917 \text{ s}^{-1}$  and  $k_{\text{slow}} = 0.080 \text{ s}^{-1}$ , and the computed fit is superimposed on the experimental data. (Data were collected over two different time intervals: 0–6 and 6–73 s.)

time scale of the hydrolysis. This report details the rate laws for the formation and decay of this intermediate.

### Experimental Section

Rate data were collected using a Durrum D110 stopped-flow spectrophotometer interfaced to an OLIS 3820 data collection system and a Northstar Horizon computer. The OLIS nonlinear least-squares program Versatile Data Fit was used to minimize observed absorbance changes according to the expression  $A_t - A_\infty = Me^{-k_{\text{fast}}t} + Ne^{-k_{\text{slow}}t}$  where  $k_{\text{fast}}$  and  $k_{\text{slow}}$  represent the first and second observed reactions and M and N are functions of the rate constants and the extinction coefficients of the reactants, intermediate, and products.<sup>9</sup> Rapid scan spectra were obtained using a Harrick rapid scan monochromator in association with the Durrum.

Equal volumes of a slightly acidified ( $\text{pH} 4.5$ )  $\text{Na}_2\text{CrO}_4$  solution ( $(4.16\text{--}8.33) \times 10^{-3} \text{ M}$ ,  $I = 1.0 \text{ M}$ ,  $\text{NaClO}_4$ ) and TRIS buffer ( $0.100\text{--}0.500 \text{ M}$ ,  $I = 1.0 \text{ M}$ ,  $\text{NaClO}_4$ ) were mixed in the Durrum spectrophotometer and the resulting absorbance decreases followed at  $450 \text{ nm}$  ( $25.0^\circ\text{C}$ ). Under these conditions the reaction mixtures initially contained some 32–43% of  $[\text{Cr}]_{\text{T}}$  as  $\text{Cr}_2\text{O}_7^{2-}$  with the remainder being  $\text{CrO}_4^{2-}$  ( $K_{\text{D}} = [\text{Cr}_2\text{O}_7^{2-}]/[\text{HCrO}_4^-]^2 = 80 \text{ M}^{-1}$ ,  $\text{p}K_{\text{a}}(\text{HCrO}_4^-) = 5.64$ ).<sup>10</sup> Rapid-scan spectra had previously shown that  $450 \text{ nm}$  gave a significant absorbance change for the second reaction as well as a clear differentiation from the first. The  $\text{pH}$  ( $25^\circ\text{C}$ ) of the recovered effluent was measured using a Radiometer PHM 62  $\text{pH}$  meter equipped with G2020B/K4040 electrodes.

Acidity constants of buffer species ( $\text{B}/\text{BH}^+$ ) are given as concentration constants,  $K_{\text{a}} = [\text{B}][\text{H}^+]/[\text{BH}^+]$ , with  $[\text{H}^+]$  determined from  $\text{pH}$  measurements,  $\text{pH} = -\log a_{\text{H}^+}$ ,  $[\text{H}^+] = a_{\text{H}^+}/\gamma_{\pm}$ , and  $\gamma_{\pm} = 0.767$  ( $25.0^\circ\text{C}$ ,  $I = 1.0 \text{ M}$ ,  $\text{NaClO}_4$ ).

### Results

Figure 1 gives a typical absorbance–time trace for the hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$  in the presence of TRIS buffer. Two rate processes are evident, both involving absorbance decreases. It was not possible to find a wavelength which gave absorbance changes in opposite senses, but the difference in time scales makes analysis of the rates quite straightforward. Table 1 lists rate constants  $k_{\text{fast}}$  and  $k_{\text{slow}}$  at different TRIS concentrations. Under the  $\text{pH}$  conditions 7.55–8.96, and at the low Cr(VI) concentrations used

- (1) Perlmutter-Hayman, B.; Wolff, M. A. *J. Phys. Chem.* **1967**, *71*, 1416.
- (2) Baharad, R.; Perlmutter-Hayman, B.; Wolff, A. *J. Phys. Chem.* **1969**, *73*, 4391.
- (3) Moore, P.; Kettle, S. F. A.; Wilkins, R. G. *Inorg. Chem.* **1966**, *5*, 221.
- (4) Moore, P.; Ducommun, Y.; Nichols, P. J.; Merbach, A. E. *Helv. Chim. Acta* **1983**, *66*, 2445.
- (5) Frennesson, S.-A.; Beattie, J. K.; Haight, G. P. *J. Am. Chem. Soc.* **1968**, *90*, 6018.
- (6) Grace, M. R.; Tregloan, P. A. *Inorg. Chem.* **1992**, *31*, 4524.
- (7) Okumura, A.; Daido, Y. *Inorg. Chim. Acta* **1988**, *144*, 63.
- (8) Muirhead, K. A.; Haight, G. P.; Beattie, J. K. *J. Am. Chem. Soc.* **1972**, *94*, 3006.


(9) Jackson, W. G.; Harrowfield, J. M. B.; Vowles, P. D. *Int. J. Chem. Kinet.* **1977**, *9*, 535.

(10) Brasch, N. E. Ph.D. Thesis, University of Otago, 1993.

**Table 1.** Rate Data for the Hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$  in the Presence of TRIS Buffer (25 °C,  $I = 1.0 \text{ M}$ ,  $\text{NaClO}_4$ )

[TRIS] <sub>T</sub> (M)	pH	$10^8[\text{H}^+]$ (M)	[TRIS] <sup>a</sup> (M)	$k_{\text{fast}}(\text{obs})$ (s <sup>-1</sup> )	$k_{\text{slow}}(\text{obs})$ (s <sup>-1</sup> )	$k_{\text{slow}}(\text{calc})^b$ (s <sup>-1</sup> )	$k_{\text{slow}}(\text{calc})^c$ (s <sup>-1</sup> )
0.250	7.55	3.67	0.0440	2.50	0.200	0.221	0.228
0.100	7.68	2.7324	0.0224	1.27	0.112	0.106	0.114
0.050	7.84	1.88	0.0147	0.917	0.080	0.063	0.074
0.100	7.86	1.80	0.0304	1.64	0.142	0.128	0.137
0.100	7.90	1.64	0.0324	1.83	0.157	0.133	0.142
0.250	8.02	1.25	0.0967	5.70	0.350	0.358	0.357
0.100	8.20	0.823	0.0487	2.70	0.175	0.150	0.160
0.200	8.22	0.786	0.100	5.70	0.310	0.300	0.300
0.400	8.23	0.768	0.202	11.4	0.580	0.600	0.579
0.200	8.24	0.750	0.102	5.87	0.325	0.299	0.299
0.100	8.60	0.327	0.0706	3.70	0.133	0.124	0.139
0.100	8.64	0.299	0.0725	3.80	0.130	0.119	0.134
0.250	8.80	0.207	0.198	10.00	0.220	0.244	0.246
0.100	8.96	0.143	0.0846	4.77	0.100	0.077	0.099

<sup>a</sup>  $[\text{TRIS}] = K_a^c[\text{TRIS}]_T/([\text{H}^+] + K_a^c)$ ;  $K_a^c = 6.03 \times 10^{-9} \text{ M}$ . <sup>b</sup> Best fit of data to eq 5, with  $k_2 = 6.2 \text{ M}^{-1} \text{ s}^{-1}$  and  $K_a^c = 8.3 \times 10^{-9} \text{ M}$  for (TRIS)CrO<sub>3</sub>.  
<sup>c</sup> Best fit of data to eq 6, with  $k_2 = 6.4 \text{ M}^{-1} \text{ s}^{-1}$ ,  $k_2' = 3.5 \times 10^{-10} \text{ M}^{-1} \text{ s}^{-1}$ , and  $K_a^c = 1.0 \times 10^{-8} \text{ M}$  for (TRIS)CrO<sub>3</sub>.

**Figure 2.** Plot of  $k_{\text{fast}}$  vs [TRIS] for reaction of  $\text{Cr}_2\text{O}_7^{2-}$  at 25 °C and  $I = 1.0 \text{ M}$  ( $\text{NaClO}_4$ ).

(( $2.08\text{--}4.16 \times 10^{-3} \text{ M}$ ), only hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$  was important; reverse dimerization was negligible.<sup>11</sup> The final absorbance was shown to be identical to that of  $\text{CrO}_4^{2-}$  at the same pH in the absence of buffer, and the separate absorbance changes ( $\Delta A_{\text{fast}}$ ,  $\Delta A_{\text{slow}}$ ) were both shown to be independent of TRIS concentration. The latter fact means that the first reaction is not measurably reversible, and the former means that we are dealing only with (catalyzed) hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$ ; i.e., no reduction of Cr(VI) to Cr(III) is involved in the time scale of the experiment. No reaction was observed when Cr(VI) and TRIS solutions initially at the same pH (i.e.  $\sim 8.0$ ) were mixed; thus we are dealing with the reaction of  $\text{Cr}_2\text{O}_7^{2-}$ , and not of  $\text{HCrO}_4^-/\text{CrO}_4^{2-}$ , with TRIS.

### Discussion

Figure 2 gives a plot of  $k_{\text{fast}}$  vs [TRIS], where TRIS represents the basic form of the buffer. The linear correspondence establishes the rate law


$$k_{\text{fast}} = k_1[\text{TRIS}] \quad (4)$$

and the near-zero intercept confirms that spontaneous hydrolysis ( $k_0 = 0.03 \text{ s}^{-1}$ ) is insignificant. Also the fact that the data agree with eq 4 irrespective of pH confirms that  $\text{OH}^-$  catalysis (eq 2) is unimportant.<sup>12</sup> The rapidity of the first reaction compared to the second (at least a 10-fold difference in rate is involved) almost certainly means that the first observed reaction (Figure 1) corresponds to the first mechanistic process,<sup>9</sup> and hence we assign it to the addition of TRIS to  $\text{Cr}_2\text{O}_7^{2-}$ ;  $k_1 = 55 \text{ M}^{-1} \text{ s}^{-1}$  at 25 °C

(11) Linge, H. G.; Jones, A. L. *Aust. J. Chem.* **1968**, *21*, 1445.


(12) Hydrolysis rate constants  $k_0 = 0.031 \text{ s}^{-1}$ ,  $k_{\text{H}} = 4.0 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ , and  $k_{\text{OH}} = 1.01 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$  at 25.0 °C and  $I = 1.0 \text{ M}$  ( $\text{NaClO}_4$ ) have been confirmed by other work in this laboratory.

### Scheme 1



### Scheme 2

#### EITHER




and  $I = 1.0 \text{ M}$  ( $\text{NaClO}_4$ ).<sup>13</sup> Scheme 1 pictures this as a concerted addition–elimination process at Cr(VI) since we view five-coordinate Cr(VI) as a transition state rather than as a transient intermediate (proton transfer from O to N may also be synchronous). Perhaps the most unusual feature of this interpretation is that of TRIS functioning as an alcohol nucleophile rather than as an amine nucleophile, with the intermediate species being  $^+\text{H}_3\text{NC}(\text{CH}_2\text{OH})_2\text{CH}_2\text{OCrO}_3^-$  (designated (TRIS)CrO<sub>3</sub>). This feature may be related to the oxidative properties of Cr(VI) toward alcohols in acidic solution<sup>14,15</sup> and suggests that direct attack by O nucleophiles at Cr(VI) is more facile than direct attack by amines.

The second reaction shows a more complex dependence on TRIS concentration and on pH. Consider the  $k_{\text{slow}}$  data at  $[\text{TRIS}]_T = 0.100 \text{ M}$ , Table 1. As the pH increases from 7.68 to 8.20,  $k_{\text{slow}}$  increases, but as the pH increases further (8.60, 8.64, 8.96), it now decreases. However at a fixed pH (e.g. 8.2)  $k_{\text{slow}}$  shows a linear increasing [TRIS]. This data fits reasonably well the rate law

(13) A contribution to the rate constant  $k_1$  (eq 4) from TRIS buffer catalyzed addition of solvent leading directly to hydrolyzed product (i.e. general base catalysis) cannot be ruled out by these results.

(14) Westheimer, F. H. *Chem. Rev.* **1949**, *45*, 419.

(15) Wiberg, K. B. In *Oxidation in Organic Chemistry*; Academic Press: New York, 1965; Chapter 2, p 69.



**Figure 3.** Spectra derived from rapid-scan measurements (440–540 nm, path length 2.0 cm) following ratio mixing (1:14.4) of  $5.555 \times 10^{-2}$  M Cr(VI) (pH 3.5) with 0.50 M TRIS buffer ( $I = 1.0$  M ( $\text{NaClO}_4$ ),  $[\text{Cr(VI)}]_{\text{T}} = 3.607 \times 10^{-3}$  M,  $25^\circ\text{C}$ , pH 8.8). Spectra correspond to the following: (A)  $1.803 \times 10^{-3}$  M  $\text{Cr}_2\text{O}_7^{2-}$ , calculated on the basis that the initial reaction mixture contains  $1.291 \times 10^{-3}$  M  $\text{Cr}_2\text{O}_7^{2-}$  and  $1.024 \times 10^{-3}$  M  $\text{CrO}_4^{2-}$  ( $K_{\text{D}} = 80$  M $^{-1}$ ; see text); (B)  $3.607 \times 10^{-3}$  M (TRIS)- $\text{CrO}_3$ , calculated on the basis that this species is fully formed 0.3 s ( $10t_{1/2}$ ) after mixing and that the initial reaction has no general base component (cf. ref 13); (C)  $3.607 \times 10^{-3}$  M  $\text{CrO}_4^{2-}$ .

$$k_{\text{slow}} = \frac{k_2[\text{TRIS}][\text{H}^+]}{K_{\text{a}}^{\text{c}} + [\text{H}^+]} \quad (5)$$

with  $k_2 = 6.2$  M $^{-1}$  s $^{-1}$  and  $K_{\text{a}}^{\text{c}} = 8.3 \times 10^{-9}$  M, where  $K_{\text{a}}^{\text{c}}$  represents the acid dissociation constant of the (TRIS) $\text{CrO}_3$  intermediate. Agreement is shown by the calculated rate constants given in the penultimate column of the table. A somewhat better fit (last column) is found by including a [TRIS]-independent pathway  $k_2'$ , viz.

$$k_{\text{slow}} = \frac{k_2[\text{TRIS}][\text{H}^+] + k_2'}{K_{\text{a}}^{\text{c}} + [\text{H}^+]} \quad (6)$$

Such a rate law is interpreted in terms of Scheme 2. The rate constant  $k_2$  represents the general TRIS-base-catalyzed hydrolysis of (TRIS) $\text{CrO}_3$  and  $k_2'$  the solvolytic OH $^{-}$  hydrolysis of the same

**Table 2.** Second-Order Rate Constants for the Catalyzed Hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$  ( $25^\circ\text{C}$ ,  $I = 1.0$ ,  $\text{NaClO}_4$ )

catalyst <sup>a</sup>	$\text{p}K_{\text{a}}^{\text{c,buffer}}^{\text{b}}$	$k_{\text{B}}$ (M $^{-1}$ s $^{-1}$ )
CAPS	10.46	76
CHES	9.48	47
DABCO	9.18	$4.35 \times 10^3$
TAPS	8.34	3.0
TRIS	8.12	55
TES	7.45	1.2
BES	7.15	0.53
BISTRIS	6.74	$2.2 \times 10^{-2}$
MES	6.25	0.82

<sup>a</sup> CAPS, 3-(cyclohexylamino)-1-propanesulfonic acid; CHES, 2-(cyclohexylamino)ethanesulfonic acid; DABCO, 1,4-diazaobicyclo[2.2.2]octane; TAPS, 3-[tris(hydroxymethyl)methylamino]-1-propanesulfonic acid; TRIS, tris(hydroxymethyl)aminomethane; TES, 2-[tris(hydroxymethyl)methylamino]-1-ethanesulfonic acid; BES, *N,N*-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid; BISTRIS, bis(2-hydroxyethyl)amino-tris(hydroxymethyl)methane; MES, 4-morpholineethanesulfonic acid. <sup>b</sup> From measurements on half-neutralized solutions (0.20 M).

species.<sup>16</sup> This interpretation gives  $k_2 = 6.4$  M $^{-1}$  s $^{-1}$ ,  $k_2(\text{OH}^-) = k_2'/K_{\text{w}} = 2.2 \times 10^4$  M $^{-1}$  s $^{-1}$ , and  $K_{\text{a}}^{\text{c}} = 1.0 \times 10^{-8}$  M. It suggests that TRIS is not as efficient a general base ( $k_2 = 6.4$  M $^{-1}$  s $^{-1}$ ) as it is as a nucleophile ( $k_1 = 55$  M $^{-1}$  s $^{-1}$ ), an observation supported by the fact that the OH $^{-}$ -catalyzed hydrolysis of (TRIS) $\text{CrO}_3$  is somewhat faster ( $2.2 \times 10^4$  M $^{-1}$  s $^{-1}$ ) than the OH $^{-}$ -catalyzed hydrolysis of  $\text{Cr}_2\text{O}_7^{2-}$  ( $1.01 \times 10^3$  M $^{-1}$  s $^{-1}$ ). The acidity of the intermediate (TRIS) $\text{CrO}_3$  species ( $\text{p}K_{\text{a}}^{\text{c}} = 8.0$ ) is similar to that of TRIS buffer itself ( $\text{p}K_{\text{a}}^{\text{c}} = 8.12$ ), which is in keeping with the amine function not being involved in the first reaction. Spectra (440–540 nm) of  $\text{Cr}_2\text{O}_7^{2-}$ ,  $\text{CrO}_4^{2-}$ , and the intermediate (TRIS)- $\text{CrO}_3$  species are given in Figure 3.

Apart from triethanolamine, which also gave clear indications of biphasic kinetics at 450 nm,<sup>17</sup> we were unable to find two-step spectrophotometric processes for other "nonnucleophilic" buffers, and the reactions followed a single exponential. Table 2 lists  $k_{\text{B}}$  values (eq 2). Of the buffers studied BISTRIS, TES, TAPS, and BES all contain—OH functionalities (BISTRIS has five), and we fully expected to see similar formation of (B) $\text{CrO}_3$  intermediates. It may be that in these cases decay of the intermediate is faster than its formation ( $k_{\text{B}} = k_1$ , eq 3). However, the bis tertiary amine DABCO is an especially effective catalyst ( $k_{\text{B}} = 4.35 \times 10^3$  M $^{-1}$  s $^{-1}$ ) and is less likely to act as a nucleophile, so that an entirely general base catalyzed role<sup>13</sup> is possible for these bases.

**Acknowledgment.** The authors thank Prof. Gil Haight for his helpful comments.

(16) Alternatively this could be interpreted as solvent ( $\text{H}_2\text{O}$ ) assisted hydrolysis of the [(TRIS-H) $\text{CrO}_3\text{H}]^{-}$  conjugate base.

(17) Biphasic kinetics were clearly evident at 450 nm, but the optical density change for the second reaction was very small making its analytical separation from the first uncertain.