

# Additions and Corrections

1989, Volume 28

**Christopher King, Ju-Chun Wang, Md. Nazrul I. Khan, and John P. Fackler, Jr.\*:** Luminescence and Metal–Metal Interactions in Binuclear Gold(I) Compounds.

Page 2147. In Table II, the  $\lambda_{\max}$  value of 571 nm for  $[\text{Au}(\text{dppm})]_2(\text{BH}_3\text{CN})_2$  should be 490 nm.

1992, Volume 31

**Runyu Han and Gerard Parkin\*:** [Tris(3-*tert*-butylpyrazolyl)-hydroborato]beryllium Hydride: Synthesis, Structure, and Reactivity of a Terminal Beryllium Hydride Complex.

Pages 983–988. The X-ray structure of a complex with terminal beryllium–hydride bonds was first reported for the dimer  $[(\text{Me}_2\text{NCH}_2\text{CH}_2\text{NMe})\text{BeH}]_2$ .<sup>1</sup> We thank Dr. Norman Bell for providing us with this information.

- (1) (a) Bell, N. A.; Coates, G. E.; Schneider, M. L.; Shearer, H. M. M. *J. Chem. Soc., Chem. Commun.* **1983**, 828–829. (b) Bell, N. A.; Coates, G. E.; Schneider, M. L.; Shearer, H. M. M. *Acta Crystallogr.* **1984**, *C40*, 608–610.

1993, Volume 32

**Krishan Kumar,\* C. Allen Chang, and M. F. Tweedle:** Equilibrium and Kinetic Studies of Lanthanide Complexes of Macrocyclic Polyamino Carboxylates.

Page 591. Equations 8 and 9 should read as follows:

$$\begin{aligned} -d[\text{LnL}]_T/dt &= k_{\text{obsd}}([\text{LnL}] + [\text{LnL}(\text{H})] + [\text{LnL}(\text{H}_2)]) \\ &= [k_d + \{k_2K_1[\text{H}^+]/(1 + K_1[\text{H}^+])\}] \times \\ &\quad [\text{LnL}(\text{H})] \quad (8) \end{aligned}$$

$$\begin{aligned} k'_{\text{obsd}} &= k_{\text{obsd}}\{(1/K_H[\text{H}^+]) + 1 + (K_1[\text{H}^+])\} \\ &= k_d + \{k_2K_1[\text{H}^+]/(1 + K_1[\text{H}^+])\} \quad (9) \end{aligned}$$

Page 592. In Table III, the values of  $k_d$  ( $\text{s}^{-1}$ ),  $K_1$  ( $\text{M}^{-1}$ ), and  $k_2$  ( $\text{s}^{-1}$ ), respectively, should read as follows:  $(4.0 \pm 1.5) \times 10^{-4}$ ,  $1.3 \pm 0.1$ ,  $(1.93 \pm 0.11) \times 10^{-2}$  for Gd(DO3A);  $b$ ,  $0.35 \pm 0.03$ ,  $(1.66 \pm 0.09) \times 10^{-3}$  for Gd(HP-DO3A).

**Tamotsu Sugimori, Kimio Shibakawa, Hideki Masuda, Akira Odani, and Osamu Yamauchi\*:** Ternary Metal(II) Complexes with Tyrosine-Containing Dipeptides. Structures of Copper(II) and Palladium(II) Complexes Involving L-Tyrosylglycine and Stabilization of Copper(II) Complexes Due to Intramolecular Aromatic Ring Stacking.

Page 4956. The caption for Figure 3 should read as follows: Species distributions as a function of pH in the 1:1:1 Cu(II)–bpy–L-tyr–L-phe system (1 mM). Species: a, Cu(bpy); b, Cu(bpy)(L); c, Cu(bpy)(LH<sub>-1</sub>); d, Cu(bpy)<sub>2</sub>; e, Cu(LH<sub>-1</sub>); f, Cu(LH<sub>-2</sub>); g, Cu(bpy)(LH<sub>-2</sub>); h, Cu(LH<sub>-2</sub>)(OH); i, Cu<sub>2</sub>(LH<sub>-1</sub>)<sub>2</sub>.

Page 4957. The right-hand side of eq 6

$$1 + \frac{1}{K_{\text{st}} + 1}$$

should read as follows:

$$1 - \frac{1}{K_{\text{st}} + 1}$$

**Hiroki Oshio,\* Etsuo Ino, Iwao Mogi, and Tasuko Ito\*:** A Weak Antiferromagnetic Interaction between Mn<sup>2+</sup> Centers through a TCNQ Column: Crystal Structures and Magnetic Properties of  $[\text{Mn}^{\text{II}}(\text{tpa})(\text{TCNQ})(\text{CH}_3\text{OH})](\text{TCNQ})_2\text{CH}_3\text{CN}$ ,  $[\text{Mn}^{\text{II}}(\text{tpa})(\mu\text{-O}_2\text{CCH}_3)_2(\text{TCNQ})_2\text{CH}_3\text{CN}$ , and  $[\text{Mn}^{\text{II}}(\text{tpa})(\text{NCS})_2]\cdot\text{CH}_3\text{CN}$  (tpa = Tris(2-pyridylmethyl)amine).

Pages 5697–5703. In this paper, TCNQ molecules (**A**, **B**, **C**, **D**) in  $[\text{Mn}^{\text{II}}(\text{tpa})(\text{TCNQ})(\text{CH}_3\text{OH})](\text{TCNQ})_2\text{CH}_3\text{CN}$  have been assigned as being anionic. The IR of **2** showed three bands (2152, 2179, 2184  $\text{cm}^{-1}$ ) characteristic of  $[\text{TCNQ}]^-$  and one band (2216  $\text{cm}^{-1}$ ) characteristic of  $[\text{TCNQ}]^0$ . These IR data lead us to conclude that **A** and **B** are monoanionic and that **C** and **D** are neutral. All other conclusions remain unaffected by this change. We thank Prof. Joel S. Miller (University of Utah) for suggesting these IR measurements.

1994, Volume 33

**Younbong Park and John D. Corbett\*:** Pr<sub>12</sub>I<sub>17</sub>Fe<sub>2</sub>: A Novel Hypostoichiometric Compound with Only Isolated Clusters.

Page 1706. In Table 2, some of the positional parameters and  $B(\text{eq})$  values were omitted or misplaced. The corrections are as follows:

atom	x	y	z	$B(\text{eq}), \text{\AA}^2$
14			0.8390(1)	2.58(4)
17		0.82610(8)	0.2541(1)	1.75(3)
18	0.37527(8)	0.2102(1)	0.3298(1)	2.55(4)