

Additions and Corrections

1994, Volume 33

Jianwei Ho, Tricia L. Breen, Andrzej Ozarowski, and Douglas W. Stephan*: Early Metal Mediated P–P Bond Formation in $\text{Cp}_2\text{M}(\text{PR})_2$ and $\text{Cp}_2\text{M}(\text{PR})_3$ Complexes.

Pages 865–870. Full details of the crystallographic study of $\text{Cp}_2\text{Zr}(\text{PPh})_3$ (**1**) and $\text{Cp}_2\text{Zr}(\text{PCy})_3$ (**2**), respectively, were previously reported: Hey-Hawkins, E.; et al. *Z. Naturforsch.* **1988**, 43B, 1271; *Z. Anorg. Allg. Chem.* **1992**, 615, 35.

IC951921R

Thomas F. Fässler* and Markus Hunziker: Ge_9^{3-} and Pb_9^{3-} : Two Novel, Naked, Homopolyatomic Zintl Ions with Paramagnetic Properties.

Page 5380. It has been brought to our attention that an isolated Ge_9^{3-} anion had been previously found in (2,2,2-crypt-K⁺)₃[P(C₆H₅)₃Ge₉³⁻]. The paramagnetic property and the crystal structure were reported in: Belin, C.; Mercier, H.; Angilella, V. *New J. Chem.* **1991**, 15, 931. Angilella, V.; Belin, C. *J. Chem. Soc., Faraday Trans.* **1991**, 87, 203. The structural similarities of the two compounds indicate that the distortions of the Ge_9^{3-} anions may not be due to electrostatic interactions or crystal packing, as anticipated before.

IC951923B