

2003, Volume 42

Patty K.-L. Fu, Patricia M. Bradley, and Claudia Turro*:
Stabilization of Duplex DNA Structure and Suppression of
Transcription in Vitro by Bis(quinone diimine) Complexes of Rh(III)
and Ru(II)

Page 880. The left side of eq 1 should read $(\epsilon_b - \epsilon_f)/$
 $(\epsilon_a - \epsilon_f)$.

IC0400179

10.1021/ic0400179

Published on Web 02/26/2004

2004, Volume 43

**Hong Zhao, Zhi-Rong Qu, Qiong Ye, Xi-Sen Wang, Jing
Zhang,* Ren-Gen Xiong,* and Xiao-Zeng You:** An Unusual
Mixed-Valence Cu(I)–Cu(II) 3-D Framework

Pages 1813–1815. The formula for compound **1** is
represented as $[\text{Cu(II)}(2\text{-PYA})_2]_3 \cdot [\text{Cu(I)}(2\text{-PYA})]_2 \cdot (\text{H}_2\text{O})_2$;
however, the crystallography indicates that the ratio of Cu(II)
to Cu(I) is 4:1. For charge balance, one of the water
molecules is required to be a hydroxide. The formula should
therefore be represented as $\text{Cu(II)}_4\text{Cu(I)}(2\text{-PYA})_8(\text{OH})(\text{H}_2\text{O})$.
We acknowledge Dr. B. Abrahams' contribution to discus-
sions relating to the formula.

IC040024X

10.1021/ic040024x

Published on Web 02/26/2004