

Silylation, Sulfidation, and Benzene-1,2-dithiolate Complexation Reactions of Oxo- and Oxosulfidomolybdates(VI) and -Tungstates(VI)

Jun-Jieh Wang and R. H. Holm*

Department of Chemistry and Chemical Biology, Harvard University,
Cambridge, Massachusetts 02138

Received June 30, 2007

The synthesis and structures of two types of molecules are presented: $[M^{VI}O_{3-n}S_n(OSiR_2R')]^{1-}$ ($M = Mo, n = 0-3$; $M = W, n = 3$) and $[M^{VI}O_2(OSiR_2R')(bdt)]^{1-}$ ($M = Mo, W$; bdt = benzene-1,2-dithiolate). For both types, R_2R' are $Me_3, Pr^i_3, Ph_3, Me_2Bu^i$ and Ph_2Bu^i . The complete series of oxo/sulfido/silyloxo molybdenum complexes has been prepared. Complexes with $n = 0$ are readily prepared by the silylation of Ag_2MoO_4 and sustain mono- or disulfidation with Ph_3SiSH to form a species with $n = 1$ and $n = 2$, respectively. Complexes with $n = 3$ are accessible by the silylation of $[MOS_3]^{2-}$. Structures of the representative series members $[MoO_3(OSiPh_2Bu^i)]^{1-}$, $[MoO_2S(OSiPh_3)]^{1-}$, $[MoOS_2(OSiPr^i_3)]^{1-}$, $[MoS_3(OSiPh_2Bu^i)]^{1-}$, and also $[WS_3(OSiMe_2Bu^i)]^{1-}$, all with tetrahedral stereochemistry, are presented. Benzene-1,2-dithiolate complexes are prepared by the reaction of $[MoO_3(OSiR_2R')]^{1-}$ with the dithiol or by the silylation of previously reported $[MO_3(bdt)]^{2-}$. The structures of $[MoO_2(OSiPh_2Bu^i)(bdt)]^{1-}$ and $[WO_2(OSiPr^i_3)(bdt)]^{1-}$ conform to square-pyramidal stereochemistry with an oxo ligand in the apical position. The role of these complexes in the preparation of site analogues of the xanthine oxidoreductase enzyme family is noted. The sulfidation reactions reported here point to the utility of Ph_3SiSH and Pr^i_3SiSH as reagents for Mo^{VI} -based oxo-for-sulfido conversions.

Introduction

We are engaged in research on the biomimetic chemistry of molybdenum and tungsten directed at the attainment of structural and functional analogues of the catalytic centers of oxotransferase and hydroxylase enzymes. Results on oxotransferase analogues have been summarized¹ and are augmented by more recent developments.²⁻⁴ Hydroxylases are members of the xanthine oxidoreductase enzyme family,⁵ and utilize coordinated hydroxide as a nucleophile in reactions with purines, aldehydes, and other substrates of this family.^{6,7} A critical structural and mechanistic feature of the apparently common, oxidized catalytic site is the presence of a basal sulfido ligand in the square-pyramidal unprotonated and protonated sites $[Mo^{VI}O_2S(S_2pd)]$ and

$[Mo^{VI}O(OH)S(S_2pd)]$, respectively.⁸ Appropriate structural analogues of these sites are shown in Figure 1, together with actual or projected methods of synthesis. In these molecules, benzene-1,2-dithiolate models the cofactor ligand and silylation simulates protonation. The approximate square-pyramidal geometry with axial and basal oxo and basal sulfido coordination follows from the X-ray structure of *Pseudomonas putida* quinoline 2-oxidoreductase.⁹ Accurate bond distances are available from the molybdenum EXAFS analysis of bovine XO.¹⁰

The sulfidation method of synthesis has recently afforded, among others, the complexes $[WO_2S(bdt)]^{2-}$ and $[WOS(OSiPr^i_3)(bdt)]^{1-}$, which are the first structural analogues of the oxidized XO sites.¹¹ Tungsten is used without structural compromise to stabilize the M^{VI} state. In the course of this and recent work, we have investigated silylation and sulfidation reactions of oxomolybdates(VI) and oxotungstates-

* To whom correspondence should be addressed. E-mail: holm@chemistry.harvard.edu.

- (1) Enemark, J. H.; Cooney, J. J. A.; Wang, J.-J.; Holm, R. H. *Chem. Rev.* **2004**, *104*, 1175–1200.
- (2) Jiang, J.; Holm, R. H. *Inorg. Chem.* **2004**, *43*, 1302–1310.
- (3) Jiang, J.; Holm, R. H. *Inorg. Chem.* **2005**, *44*, 1068–1072.
- (4) Wang, J.-J.; Tessier, C.; Holm, R. H. *Inorg. Chem.* **2006**, *45*, 2979–2988.
- (5) Hille, R. *Chem. Rev.* **1996**, *96*, 2757–2816.
- (6) Hille, R. *Arch. Biochem. Biophys.* **2005**, *433*, 107–116.
- (7) Hille, R. *Eur. J. Inorg. Chem.* **2006**, 1913–1926.

(8) Abbreviations are given in the chart.

- (9) Bonin, I.; Martins, B. M.; Purvanov, V.; Fetzner, S.; Huber, R.; Dobbek, H. *Structure* **2004**, *12*, 1425–1435.
- (10) Doonan, C. J.; Stockert, A.; Hille, R.; George, G. N. *J. Am. Chem. Soc.* **2005**, *127*, 4518–4522.
- (11) Wang, J.-J.; Groysman, S.; Lee, S. C.; Holm, R. H. *J. Am. Chem. Soc.* **2007**, *129*, 7512–7513.






Figure 1. Synthetic analogues of the oxidized active sites in the xanthine oxidoreductase family of enzymes and their precursors.

(VI), leading to species of potential utility in the preparation of XO site analogues. For example, the ions $[\text{MO}_3\text{S}]^{2-}$ ($\text{M} = \text{Mo}, \text{W}$), difficult to obtain as pure R_4N^+ salts by the classical reaction of aqueous $[\text{MO}_4]^{2-}$ with H_2S , are readily prepared by the more easily controlled reactions 1 and 2.¹² Here, we describe the synthesis of a family of oxo/sulfido/silyloxo complexes of Mo^{VI} and W^{VI} comprised primarily of previously unreported or incompletely characterized types, and the structures of representative members. A later report will enlarge on the use of such complexes in the synthesis of oxo/sulfido site analogues.



Experimental Section

Preparation of Compounds. All of the reactions and manipulations were performed under a pure dinitrogen atmosphere using either an inert atmosphere box or standard Schlenk techniques. Acetonitrile, diethylether, dichloromethane, and THF were freshly purified using an Innovative Technology solvent purification system and stored over 4 Å molecular sieves. Benzene-1,2-dithiol ($\text{H}_2(\text{bdt})$) was prepared as described.¹³ ^1H NMR data for salts refer to anions in CD_3CN solutions. IR spectra were recorded in KBr pellets and absorption spectra in acetonitrile solutions. All of the compounds were characterized by a combination of IR, ^1H NMR, UV-vis, mass spectroscopy, and elemental analysis. Negative-ion electrospray mass spectra of all of the complex salts reveal a prominent peak for the parent anion (M^-). Certain compounds not subjected to elemental analysis were identified by X-ray structure determinations.

Oxo/Sulfido/Silyloxo Complexes. $(\text{Et}_4\text{N})[\text{MoO}_3(\text{OSiR}_2\text{R}')].$ These compounds were prepared on a 5–8 mmol scale, by a method analogous to that reported for $(\text{Et}_4\text{N})[\text{MoO}_3(\text{OSiPh}_3)]^{12}$ but with the use of $\text{R}_2\text{R}'\text{SiCl}$. A solution of $\text{R}_2\text{R}'\text{SiCl}$ (1.0 equiv) in dichloromethane was added dropwise to a suspension of Ag_2MoO_4 (2–3 g) in dichloromethane. The reaction mixture was stirred for 72 h and 1.0 equiv of Et_4NCl was added. The mixture was stirred for 8 h, filtered through Celite, and the filtrate was taken to dryness, leaving a solid white residue. This material was dissolved in 20–30 mL of dichloromethane, the solution was filtered, and the filtrate solvent was removed. This step was repeated with the residue, the resultant solid was extracted with acetonitrile, and the extract was filtered through Celite. The filtrate was reduced to half-volume, several volume equiv of ether were layered on the filtrate, and the mixture was allowed to stand for 2 d. Products were isolated as crystalline white solids in the indicated yields.

$\text{R} = \text{R}' = \text{Me}.$ Needlelike crystals, 49%. IR: 955, 902, 878 cm^{-1} . ES-MS: m/z 233 (M^-). ^1H NMR: δ 0.11 (s).

$\text{R} = \text{R}' = \text{Pr}^i.$ Platelike crystals, 59%. IR: 971, 906, 679 cm^{-1} . ES-MS: m/z 317 (M^-). ^1H NMR: δ 0.14 (sept, 3), 1.07 (d, 18). Anal. Calcd for $\text{C}_{17}\text{H}_{41}\text{MoNO}_4\text{Si}$: C, 45.62; H, 9.23; N, 3.13. Found: C, 45.48; H, 9.30; N, 3.09.

$\text{R} = \text{Me}, \text{R}' = \text{Bu}^t.$ Blocklike crystals, 55%. IR: 973, 907, 878, 678 cm^{-1} . ES-MS: m/z 275 (M^-). ^1H NMR: δ 0.06 (s, 6), 0.92 (s, 9).

$\text{R} = \text{Ph}, \text{R}' = \text{Bu}^t.$ Platelike crystals, 67%. IR: 972, 904, 885, 703 cm^{-1} . ^1H NMR: δ 1.03 (s, 9), 7.39 (m, 6), 7.74 (m, 4). ES-MS: m/z 399 (M^-). Anal. Calcd for $\text{C}_{24}\text{H}_{39}\text{MoNO}_4\text{Si}$: C, 54.43; H, 7.42; N, 2.64. Found: C, 54.38; H, 7.30; N, 2.69.

$(\text{Et}_4\text{N})[\text{MoO}_2\text{S}(\text{OSiPh}_2\text{Bu}^t)].$ The procedure was performed at -20°C . A solution of Ph_3SiSH (112 mg, 0.383 mmol) in 2 mL of THF was added dropwise to a solution of $(\text{Et}_4\text{N})[\text{MoO}_3(\text{OSiPh}_2\text{Bu}^t)]$ (201 mg, 0.380 mmol) in 8 mL of acetonitrile. The reaction mixture turned bright orange within 3 min, and stirring was continued for 10 min. Solvents were removed, and the yellow-orange solid was washed with hexanes and dissolved in 1.5 mL of acetonitrile. The solution was filtered, and the filtrate was layered with 20 mL of ether. Over 8 h, a solid separated, which was washed with ether (2×3 mL) and dried, to afford the product as 171 mg (83%) of yellow-orange crystals. Absorption spectrum: λ_{max} (ϵ_{M}) 256 (5820), 311 (6780), 399 (sh, 770), 455 (sh, 1100) nm. ES-MS: m/z 415 (M^-). ^1H NMR: δ 1.04 (s, 9), 7.42 (m, 6), 7.75 (m, 4). Anal. Calcd for $\text{C}_{24}\text{H}_{39}\text{MoNO}_3\text{SSi}$: C, 52.83; H, 7.20; N, 2.57; S, 5.88. Found: C, 52.68; H, 7.10; N, 2.48; S, 5.86.

$(\text{Et}_4\text{N})[\text{MoO}_2\text{S}(\text{OSiPr}^i_3)].$ The previous procedure was followed with the use of 0.40 mmol each of $(\text{Et}_4\text{N})[\text{MoO}_3(\text{OSiPr}^i_3)]$ and Pr^i_3SiSH . The product was obtained as 168 mg (90%) of yellow-orange crystals. IR: 949, 901, 881, 500 cm^{-1} . Absorption spectrum: λ_{max} (ϵ_{M}) 255 (5370), 310 (4440), 350 (1860), 472 (280) nm. ES-MS m/z 333 (M^-). ^1H NMR: δ 1.09 (d) (CH not detected). Anal. Calcd for $\text{C}_{17}\text{H}_{41}\text{MoNO}_3\text{SSi}$: C, 44.04; H, 8.91; N, 3.02; S, 6.92. Found: C, 43.87; H, 9.02; N, 2.97; S, 6.85.

$(\text{Et}_4\text{N})[\text{MoOS}_2(\text{OSiR}_2\text{R}')].$ To a solution of $(\text{Et}_4\text{N})[\text{MoO}_3(\text{OSiR}_2\text{R}')] (0.38\text{--}0.40$ mmol) in 6 mL of acetonitrile was added a solution of Ph_3SiSH (2.0 equiv) in 2 mL of THF. The reaction mixture turned bright orange within 1 min and was stirred for 10 min to give a deep-orange solution. The solution was filtered and the solvents were removed. The orange solid was washed with hexanes and ether and dissolved in 1.5 mL of acetonitrile. The solution was filtered, and the filtrate was layered with 20 mL of ether. The solid that separated over 8 h was washed with ether (2×3 mL) and dried to afford the products as orange platelike crystals in the indicated yields.

$\text{R} = \text{R}' = \text{Pr}^i.$ 84%. IR: 938, 899, 507 cm^{-1} . Absorption spectrum: λ_{max} (ϵ_{M}) 289 (4120), 300 (sh, 3950), 350 (4400), 404 (1260), 535 (190) nm. ES-MS: m/z 351 (M^-). ^1H NMR: δ 1.09

(12) Partyka, D. V.; Holm, R. H. *Inorg. Chem.* **2004**, *43*, 8609–8616.

(13) Giolando, D. M.; Kirschbaum, K. *Synthesis* **1992**, 451–452.

Table 1. Crystallographic Data for Compounds Containing $[\text{MO}_3 - n\text{S}_n(\text{OSiR}_2\text{R}')^{-1}]^{-}$ ($\text{M} = \text{Mo}$, $n = 0-3$; $\text{M} = \text{W}$, $n = 3$)^a

	(Et ₄ N)[1d]·CH ₂ Cl ₂	(Et ₄ N)[2b]	(Et ₄ N)[3a]	(Et ₄ N)[4e]	(Et ₄ N)[5c]
formula	C ₂₅ H ₄₁ Cl ₂ MoNO ₄ S _i	C ₂₆ H ₃₅ MoNO ₂ SSi	C ₁₇ H ₄₁ MoNO ₂ S ₂ Si	C ₂₄ H ₃₉ MoNOS ₃ Si	C ₁₄ H ₃₅ NOS ₃ SiW
fw	614.52	565.64	479.66	577.77	541.55
cryst syst	triclinic	triclinic	monoclinic	monoclinic	monoclinic
space group	<i>P</i> 1̄	<i>P</i> 1̄	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> 2 ₁ / <i>c</i>
Z	2	2	4	4	4
<i>a</i> (Å)	9.830(2)	8.5965(8)	17.7637(14)	20.6752(16)	18.678(4)
<i>b</i> (Å)	10.660(2)	9.2168(9)	8.4492(7)	10.3777(8)	8.1799(16)
<i>c</i> (Å)	15.920(3)	19.4188(18)	16.5049(14)	14.4721(11)	16.724(3)
α (deg)	107.21(3)	100.911(2)			
β (deg)	103.20(3)	92.755(2)	90.732(2)	110.422(1)	110.59(3)
γ (deg)	93.31(3)	113.094(2)			
<i>V</i> (Å ³)	1537.3(5)	1376.9(2)	2477.0(4)	2910.0(4)	2392.0(8)
<i>d</i> _{calcd} (g/cm ³)	1.328	1.364	1.286	1.319	1.504
μ (mm ⁻¹)	0.667	0.622	0.756	0.723	5.141
θ range (deg)	1.39–27.90	1.08–25.00	1.15–25.00	1.05–28.28	1.16–27.89
GOF (<i>F</i> ²)	1.042	1.080	1.089	1.107	1.176
R1 ^b (%)	3.10	4.90	4.79	3.50	3.22
wR2 ^c (%)	8.33	13.47	10.31	8.95	8.36

^a Mo Kα radiation, 193 K. ^b R1 = $\sum||F_o| - |F_c||/\sum|F_o|$. ^c wR2 = $[\sum w(F_o^2 - F_c^2)^2/\sum(F_o^2)^2]^{1/2}$.

(d) (CH not detected). Anal. Calcd for C₁₇H₄₁MoNO₂S₂Si: C, 42.57; H, 8.62; N, 2.92; S, 13.37. Found: C, 42.45; H, 8.76; N, 2.85; S, 13.44.

R = R' = Ph. 88%. IR: 964, 906, 885, 510 cm⁻¹. Absorption spectrum: λ_{max} (ε_M) 260 (9280), 313 (6530), 352 (2830), 406 (sh, 1530) nm. ES-MS: *m/z* 453 (M⁻). ¹H NMR: δ 7.43 (m, 9), 7.62 (m, 6).

R = Ph, R' = Bu^t. The procedure was conducted at 0 °C.; 82%. IR: 972, 904, 885, 703 cm⁻¹. Absorption spectrum: λ_{max} (ε_M) 256 (8500), 302 (sh, 4020), 350 (4400), 404 (sh, 1300), 470 (660) nm. ES-MS: *m/z* 431 (M⁻). ¹H NMR: δ 1.05 (s, 9), 7.42 (m, 6), 7.62 (m, 2), 7.76 (m, 2).

(Et₄N)[MoOS₂(OSiMe₂Bu^t)]. A solution of Me₂Bu^tSiCl (57 mg, 0.39 mmol) in 2 mL of THF was added to a suspension of (Et₄N)₂[MoO₃S]¹² (166 mg, 0.38 mmol) in 8 mL of THF. The reaction mixture became yellowish and then bright orange in 10 min and was stirred for 30 min, resulting in a deep-orange solution and a precipitate. The mixture was filtered, and the filtrate was taken to dryness. The solid residue was washed with ether (3 × 3 mL) and dissolved in 1.5 mL of acetonitrile. The solution was covered with 20 mL of ether, and the mixture was allowed to stand overnight. The product was isolated as 78 mg (43%) of orange platelike crystals. IR: 956, 882, 506 cm⁻¹. Absorption spectrum: λ_{max} (ε_M) 288 (4020), 301 (sh, 3890), 350 (4380), 406 (sh, 1240), 438 (780), 536 (140) nm. ES-MS: *m/z* 307 (M⁻). ¹H NMR: δ 0.06 (s, 6), 0.92 (s, 9).

(Et₄N)[MoS₃(OSiR₂R')]. These compounds were prepared by a method and on a scale similar to the preceding preparation. A solution of R₂R'SiCl (1.0 equiv) in THF was added to a suspension of (Et₄N)₂[MoOS₃]¹⁴ (1.0 equiv) in THF. The reaction mixture became orange within 5 min and was stirred for 30 min. Workup was performed as in the preceding preparation, to afford the products as red blocklike crystals in the indicated yields.

R = R' = Me. 49%. IR: 921, 510, 502 cm⁻¹. ES-MS: *m/z* 233 (M⁻). Absorption spectrum: λ_{max} (ε_M) 290 (4800), 347 (sh, 1300), 408 (2570), 535 (450) nm. ¹H NMR: δ 0.13 (s). This compound can also be prepared by the reaction of (Et₄N)[MoO₃(OSiMe₃)] and ≥3 equiv of (Me₃Si)₂S in acetonitrile.

R = R' = Prⁱ. 69%. IR: 922, 507 cm⁻¹. ES-MS: *m/z* 317 (M⁻). Absorption spectrum: λ_{max} (ε_M) 287 (5000), 349 (sh, 960), 407 (2280), 440 (sh, 900), 534 (600) nm. ¹H NMR: δ 1.06 (d) (CH

Table 2. Crystallographic Data for Compounds Containing $[\text{MO}_2(\text{OSiR}_2\text{R}')(\text{bdt})]^{-}$ ($\text{M} = \text{Mo}$, W) and $[\text{Mo}_2\text{O}_3\text{Cl}_2(\text{bdt})_2]^{2-}$ ^a

	(Et ₄ N)[8d]	(Et ₄ N)[9b]	(Et ₄ N) ₂ [10]
formula	C ₃₀ H ₄₃ MoNO ₃ S ₂ Si	C ₂₃ H ₄₅ NOS ₂ SiW	C ₂₈ H ₄₈ Cl ₂ Mo ₂ N ₂ O ₃ S ₄
fw	653.80	659.66	851.70
cryst syst	orthorhombic	triclinic	monoclinic
space group	<i>Pna</i> 2 ₁	<i>P</i> 1̄	<i>P</i> 2 ₁ / <i>c</i>
Z	4	4	2
<i>a</i> (Å)	23.374(6)	8.4161(4)	8.3500(17)
<i>b</i> (Å)	8.965(2)	17.3262(9)	16.477(3)
<i>c</i> (Å)	15.057(3)	19.9065(11)	13.535(3)
α (deg)		84.9760(10)	
β (deg)		88.8130(10)	104.00(3)
γ (deg)		89.7990(10)	
<i>V</i> (Å ³)	3155.3(13)	2891.0(3)	1806.9(6)
<i>d</i> _{calcd} (g/cm ³)	1.376	1.516	1.565
μ (mm ⁻¹)	0.616	4.204	1.104
θ range (deg)	1.74–27.90	1.18–27.91	1.98–27.89
GOF (<i>F</i> ²)	1.093	1.089	1.046
R1 ^b (%)	3.01	4.71	5.15
wR2 ^c (%)	6.38	9.54	11.95

^a Mo Kα radiation, 193 K. ^b R1 = $\sum||F_o| - |F_c||/\sum|F_o|$. ^c wR2 = $[\sum w(F_o^2 - F_c^2)^2/\sum(F_o^2)^2]^{1/2}$.

not detected). Anal. Calcd for C₁₇H₄₁MoNOS₃Si: C, 41.19; H, 8.34; N, 2.83; S, 19.40. Found: C, 41.31; H, 8.40; N, 2.76; S, 19.34. This compound can also be prepared by the reaction of (Et₄N)-[MoO₃(OSiPrⁱ₃)] and ≥3 equiv of Prⁱ₃SiSH in acetonitrile.

R = R' = Ph. 70%. ES-MS: *m/z* 467 (M⁻). IR: 946, 518, 500 cm⁻¹. Absorption spectrum: λ_{max} (ε_M) 273 (4130), 291 (sh, 3900), 445 (2200) nm. ¹H NMR: δ 7.42 (m, 9), 7.63 (m, 6). This compound can also be prepared by the reaction of (Et₄N)[MoO₃-(OSiPh₃)] and ≥3 equiv of Ph₃SiSH in acetonitrile.

R = Me, R' = Bu^t. 55%. IR: 924, 508 cm⁻¹. ES-MS: *m/z* 275 (M⁻). Absorption spectrum: λ_{max} (ε_M) 285 (5500), 347 (sh, 1360), 405 (2770), 437 (sh, 1200), 535 (600) nm. ¹H NMR: δ 0.08 (s, 6), 0.92 (s, 9).

R = Ph, R' = Bu^t. 68%. IR: 933, 508, 478 cm⁻¹. ES-MS: *m/z* 399 (M⁻). Absorption spectrum: λ_{max} (ε_M) 271 (4060), 300 (sh, 2720), 382 (sh, 630), 448 (520) nm. ¹H NMR: δ 1.06 (s, 9), 7.42 (m, 6), 7.74 (m, 4).

(Et₄N)[WS₃(OSiR₂R')]. These compounds were prepared on a 5–8 mmol scale in a procedure analogous to the preceding preparations, but with (Et₄N)₂[WOS₃]¹⁴ and were isolated as yellow crystalline solids in the indicated yields.

R = R' = Prⁱ. 89%. IR: 936, 901, 490, 482 cm⁻¹. ES-MS: *m/z* 453 (M⁻). Absorption spectrum: λ_{max} (ε_M) 254 (9000), 283 (sh,

(14) McDonald, J. W.; Friesen, G. D.; Rosenhein, L. D.; Newton, W. E. *Inorg. Chim. Acta* **1983**, *72*, 205–210.




Figure 3. Structures of $[\text{MoO}_3(\text{OSiPh}_2\text{Bu}')]^{1-}$ (left), $[\text{MoO}_2\text{S}(\text{OSiPh}_3)]^{1-}$ (right), and $[\text{MoOS}_2(\text{OSiPr}_3)]^{1-}$ (center). In this and the following structural figures, 50% probability ellipsoids, the atom-labeling schemes, and selected (mean) bond distances (angstroms) and angles (degrees) are given. Left: mean Mo–O_{1,2,3} 1.721(2), Mo–O₄ 1.876(2), O₄–Si 1.622(2), Si–C 1.876(2); O₁–Mo–O₂ 109.3(1), O₂–Mo–O₃ 108.8(1), O₁–Mo–O₃ 109.3(1), O₁–Mo–O₄ 110.1(1), O₂–Mo–O₄ 108.7(1), O₃–Mo–O₄ 110.7(1), Mo–O₄–Si 169.8(1). Right: Mo–O₁ 1.792(4), Mo–O₂ 1.719(3), Mo–S₁ 2.104(2), Mo–O₃ 1.883(3), O₃–Si 1.622(3), Si–C 1.865; O₁–Mo–O₂ 108.0(2), O₁–Mo–S₁ 108.8(2), O₂–Mo–S₁ 108.5(2), O₁–Mo–O₃ 110.7(2), O₂–Mo–O₃ 107.7(2), S₁–Mo–O₃ 113.0(1), Mo–O₃–Si 161.2(2). Center: Mo–O₁ 1.70(2), Mo–S₁ 2.150(1), Mo–S₂ 2.06(1), Mo–O₂ 1.88(1), O₂–Si 1.63(1), Si–C 1.87(1); S₁–Mo–S_{2A} 110.6(1), O_{1A}–Mo–S₁ 105.6(4), O_{1A}–Mo–S_{2A} 111.6(5), O_{1A}–Mo–O_{2A} 118.8(5), S₁–Mo–O_{2A} 106.4(4), S_{2A}–Mo–O_{2A} 103.7(4), Mo–O_{2A}–Si 156.3(8).

such that 971–1647 frames were collected for a hemisphere of data. The first 50 frames were recollected at the end of the data collection to monitor for crystal decay; no significant decay was detected for any compound. Cell parameters were retrieved using *SAINT* software and refined with *SAINT* on all of the reflections. Data integration was performed with *SAINT*, which corrects for Lorentz polarization and decay. Absorption corrections were applied with *SADABS*. Space groups were assigned unambiguously by analysis of symmetry, and systematic absences were determined by *XPREP*. All of the crystals were checked with *PLATON* for the possibility of higher symmetry; none was found. Structures were solved by direct methods and refined against all of the data by full-matrix least-squares techniques on F^2 using the *SHELXL-97* package. All of the non-hydrogen atoms were refined anisotropically. Isopropyl groups in anions **3a**, **4b**, **5a**, **8b**, and **9b** were disordered. The disorder was modeled effectively in all cases except **3a**, for which no satisfactory description of the disorder was obtained. Hydrogen atoms were placed in idealized positions on carbon atoms and refined as a riding model. Crystal parameters and agreement factors are collected in Tables 1 and 2.¹⁵

(15) See paragraph at the end of this article for Supporting Information available.

In addition to the compounds in Tables 1 and 2, several other structures were determined. Owing to the close structural similarities of their complexes to those described in full, only crystallographic parameters are reported.¹⁵ $(\text{Et}_4\text{N})[\text{MoS}_3(\text{OSiPr}_3)]$: monoclinic, $P2_1/c$, $a = 17.990(4)$ Å, $b = 8.490(2)$ Å, $c = 16.505(1)$ Å, $\beta = 90.79(3)^\circ$, $Z = 4$, $R_1(wR_2) = 0.1030(0.2617)$. $(\text{Et}_4\text{N})[\text{WS}_3(\text{OSiPr}_3)]$: monoclinic, $P2_1/c$, $a = 18.061(2)$ Å, $b = 8.523(1)$ Å, $c = 16.662(2)$ Å, $\beta = 91.032(2)^\circ$, $Z = 4$, $R_1(wR_2) = 0.0559(0.1242)$. $(\text{Et}_4\text{N})[\text{MoO}_2(\text{OSiPr}_3)(\text{bdt})]$: triclinic, $P\bar{1}$, $a = 8.400(2)$ Å, $b = 17.300(4)$ Å, $c = 19.860(4)$ Å, $\alpha = 85.08(3)^\circ$, $\beta = 88.78(3)^\circ$, $\gamma = 89.70(3)^\circ$, $Z = 4$, $R_1(wR_2) = 0.0495(0.1057)$. $(\text{Et}_4\text{N})[\text{WO}_2(\text{OSiMe}_2\text{Bu}')(\text{bdt})]$: monoclinic, $P2_1/c$, $a = 19.940(4)$ Å, $b = 7.830(2)$ Å, $c = 34.290(7)$ Å, $\beta = 92.17(3)^\circ$, $Z = 8$, $R_1(wR_2) = 0.0593(0.1329)$.

Other Physical Measurements. All of the measurements were made under anaerobic conditions. NMR spectra were recorded on a Varian Mercury 300 and 400 spectrometers. Infrared spectra were obtained on a Nicolet Nexus 470 FTIR spectrometer. Absorption spectra were measured with a Varian Cary 50 Bio spectrophotometer. Electrospray mass spectra were recorded on a Micromass

(16) Do, Y.; Simhon, E. D.; Holm, R. H. *Inorg. Chem.* **1985**, *24*, 1831–1838.

(17) Lim, B. S.; Willer, M. W.; Miao, M.; Holm, R. H. *J. Am. Chem. Soc.* **2001**, *123*, 8343–8349.




Figure 4. Structures of [MoS₃(OSiPh₂Bu^t)]¹⁻ (upper): mean Mo–S_{1,2,3} 2.150(1), Mo–O1 1.857(1), O1–Si 1.636(2), Si–C 1.875(2); S1–Mo–S2 109.17(3), S1–Mo–S3 109.29(3), S2–Mo–S3 109.23(3), S1–Mo–O1 108.91(5), S2–Mo–O1 109.93(5), S3–Mo–O1 110.30(6), Mo–O1–Si 171.4(1). [WS₃(OSiMe₂Bu^t)]¹⁻ (lower): mean W–S_{1,2,3} 2.154(1), W–O1 1.858(4), O1–Si 1.634(4), Si–C 1.861(8); S1–W–S2 109.33(5), S1–W–S3 109.33(6), S2–W–S3 109.40(6), S1–W–O1 110.0(2), S2–W–O1 110.4(1), S3–W–O1 108.4(1), W–O1–Si 155.2(3).

(Danvers, MA) Platform II quadrupole mass spectrometer. Elemental analyses were performed by H. Kolbe (Mülheim, Germany).

Results and Discussion

Reactions 3–8 leading to oxo/sulfido/silyloxy molybdates(VI) and tungstates(VI) are summarized by the scheme in Figure 2. All of the compounds were isolated as Et₄N⁺ salts. Structures of representative complexes are presented in Figures 3 and 4 together with selected metric data. In the sections that follow, complexes are numerically designated, according to Chart 1. Complexes **1**–**4** constitute the series [MoO_{3–n}S_n(OSiR₂R′)]¹⁻ (*n* = 0–3).


Oxosilyloxomolybdates. The scheme is initiated by the direct silylation reaction 3 of Ag₂MoO₄ in dichloromethane, affording colorless complexes [MoO₃(OSiR₂R′)]¹⁻ as **1a**–**1d** in yields of 49–67%. **1e** was prepared recently by this method.¹² The first examples of this type were obtained by the silylation of [MoO₂O₇]²⁻ with R₃SiOH.¹⁸ Subsequently, it was shown that excess Ph₃SiCl with Ag₂MoO₄ yielded the doubly silylated product [MoO₂(OSiPh₃)₂].¹⁹ The method also

Chart 1. Designation of Complexes and Abbreviations^a



[MoO ₃ (OSiR ₂ R′)] ¹⁻	R ₂ R′ = Me ₃ 1a , Pr ^{<i>i</i>} ₃ 1b , Me ₂ Bu ^{<i>t</i>} 1c , Ph ₂ Bu ^{<i>t</i>} 1d , Ph ₃ 1e ^{12,18}
[MoO ₂ S(OSiR ₂ R′)] ¹⁻	R ₂ R′ = Pr ^{<i>i</i>} ₃ 2a , Ph ₃ 2b , ¹² Ph ₂ Bu ^{<i>t</i>} 2c
[MoOS ₂ (OSiR ₂ R′)] ¹⁻	R ₂ R′ = Pr ^{<i>i</i>} ₃ 3a , Ph ₃ 3b , Me ₂ Bu ^{<i>t</i>} 3c , Ph ₂ Bu ^{<i>t</i>} 3d
[MoS ₃ (OSiR ₂ R′)] ¹⁻	R ₂ R′ = Me ₃ 4a , ¹⁶ Pr ^{<i>i</i>} ₃ 4b , Ph ₃ 4c , Me ₂ Bu ^{<i>t</i>} 4d , Ph ₂ Bu ^{<i>t</i>} 4e
[WS ₃ (OSiR ₂ R′)] ¹⁻	R ₂ R′ = Pr ^{<i>i</i>} ₃ 5a , Ph ₃ 5b , Me ₂ Bu ^{<i>t</i>} 5c , Ph ₂ Bu ^{<i>t</i>} 5d
[Mo ₃ (bdt)] ²⁻ ¹²	M = Mo 6 , W 7
[MoO ₂ (OSiR ₂ R′)(bdt)] ¹⁻	R ₂ R′ = Me ₃ 8a , Pr ^{<i>i</i>} ₃ 8b , Me ₂ Bu ^{<i>t</i>} 8c , Ph ₂ Bu ^{<i>t</i>} 8d , Ph ₃ 8e ¹⁷
[WO ₂ (OSiR ₂ R′)(bdt)] ¹⁻	R ₂ R′ = Me ₃ 9a , Pr ^{<i>i</i>} ₃ 9b , Me ₂ Bu ^{<i>t</i>} 9c , Ph ₂ Bu ^{<i>t</i>} 9d , Ph ₃ 9e ¹²
[Mo ₂ O ₂ Cl ₂ (bdt) ₂ (μ ₂ -O)] ²⁻	10

^a bdt = benzene-1,2-dithiolate(2-); Bu^{*t*}tach = 1,3,5-*tert*-butyl-1,3,5-triazacyclohexane; Cp* = pentamethylcyclopentadienide(1-); S₂pd = pyranopterindithiolate(2-) cofactor ligand; Tp = tris(pyrazolyl)hydroborate(1-); XO = xanthine oxidoreductase.

succeeds with AgMoO₄, resulting in [Me₃SiOMO₃] (M = Tc,²⁰ Re²¹). The structure of **1d** (Figure 3) reveals the expected tetrahedral stereochemistry at the molybdenum atom (O–Mo–O angles of 108.9(1)–110.7(1)°, a mean Mo=O distance of 1.721(2) Å, a Mo–O_{Si} distance of 1.876(2) Å, and a nearly linear Mo–O–Si angle of 169.8(1)°. These complexes are precursors to monosulfido species. Despite repeated attempts under different conditions, no reaction was observed between Ag₂WO₄ and silylchlorides. A route to the complexes [WO₃(OSiR₂R′)]¹⁻ remains to be developed.



Oxothiosilyloxomolybdates. Monosulfidation reaction 4a was first used to prepare **2b** from **1e**.¹² Application of this reaction to **1d** was successful, yielding **2c** (83%). However, this procedure was unsuccessful with the trialkylsilyloxides **1a**, **1b**, and **1c**, affording instead **2b** in good yield. This species was identified by mass spectrometry and a crystal structure determination (Figure 3). In these reactions, sulfidation occurs but with apparent protonation of the bound silyloxy group and its departure as the silanol, followed by the coordination of Ph₃SiS⁻, nucleophilic attack of an oxo ligand at silicon, and Si–S bond breaking to generate the product. Such a sequence is avoided in the formation of **2c** because the capacious Ph₂Bu^{*t*}SiO group hinders the encroachment of also bulky Ph₃SiSH; additionally, the silyloxy group contains only one electron-releasing alkyl substituent. This situation can be avoided by the use of a less acidic silylthiol and/or the same silyl substituents, as in reaction 4b, which gives yellow-orange **2a** (90%). The structure of **2b** resembles that of **1d**, but with a Mo=O distance of 2.144(3) Å and a Mo–O–Si angle of 160.4(5)°.




(18) Klemperer, W. G.; Mainz, V. V.; Wang, R.-C.; Shum, W. *Inorg. Chem.* **1985**, *24*, 1970–1971.

(19) Huang, M.; DeKock, C. W. *Inorg. Chem.* **1993**, *32*, 2287–2291.

(20) Nugent, W. A. *Inorg. Chem.* **2007**, *22*, 965–969.

(21) Schmidt, M.; Schmidbaur, H. *Inorg. Synth.* **1967**, *9*, 149–151.



[WO₂(OSiPrⁱ)₃(bdt)]¹⁻




Figure 5. Structures of $[\text{MoO}_2(\text{OSiPh}_2\text{Bu}'(\text{bdt}))]^{1-}$ (upper): Mo–O1 1.700(2), Mo–O2 1.723(2), Mo–O3 1.911(2), Mo–S1 2.447(1), Mo–S2 2.472(1), S1–Mo–S2 79.47(2), $\theta = 132.1$, $\delta = 0.56$. $[\text{WO}_2(\text{OSiPr}^i_3)(\text{bdt})]^{1-}$ (lower): W–O1 1.724(4), W–O2 1.731(4), W–O3 1.886(4), W–S1 2.434(2), W–S2 2.476(1), S1–W–S2 80.07(5), $\theta = 131.0$, $\delta = 0.57$.

[Mo₂O₃Cl₂(bdt)₂]²⁻





Figure 6. Structure of centrosymmetric $[\text{Mo}_2\text{O}_3\text{Cl}_2(\text{bdt})_2]^{2-}$: Mo–O1 1.689(3), Mo–O2 1.877(1), Mo–Cl 2.395(1), Mo–S1 2.385(1), Mo–S2 2.393(1), S1–Mo–S2 83.14(4), $\theta = 130.7$, $\delta = 0.68$.

Attempts to prepare pure monosulfido **2** by the silylation of $[\text{MoO}_3]^{2-}$ were not productive. In one case, reaction 5 with $\text{Me}_2\text{Bu}'\text{SiCl}$ (Figure 2) afforded the disulfido **3c** (43%). The course of this reaction is unclear. Subsequently, disulfido **3a**, **3b**, and **3d** were obtained by double-sulfidation reaction 6 as orange solids in 82–88% yield. The obvious intermediate is a monosulfido complex, as demonstrated by the clean conversions $2\text{b} \rightarrow 3\text{b}$ and $2\text{c} \rightarrow 3\text{d}$ with 1.0 equiv of Ph_3SiSH in reaction 7 (Figure 2). The structure of **3a** (Figure 3) closely resembles others in the series, with $\text{Mo}=\text{O}$ and

average $\text{Mo}=\text{S}$ distances of 1.880(4) and 2.13 Å, respectively, and a $\text{Mo}-\text{O}-\text{Si}$ angle of $159.1(4)^\circ$.



Trisulfido complexes are readily accessible by the silylation of the anions $[\text{MOS}_3]^{2-}$ ($\text{M} = \text{Mo}, \text{W}$) in reaction 8. Molybdenum complexes **4a–4e** were isolated as red crystals (49–70%) and tungsten complexes **5a–5d** as yellow crystals (75–89%). **4a**, **4b**, and **4c** were also prepared in high purity and yield by the reaction of the corresponding trioxo complexes with ≥ 3 equiv of $(\text{Me}_3\text{Si})_2\text{S}$, Pr^i_3SiSH , and Ph_3SiSH , respectively, in acetonitrile. The first of these reactions is preceded by the reaction of $[\text{Mo}_2\text{O}_7]^{2-}$ with excess $(\text{Me}_3\text{Si})_2\text{S}$ to afford **4a**.¹⁶ The MS_3OSi fragments of **4e** and **5c** are isostructural and nearly isometric (Figure 4). Parameters of **4e** include a mean $\text{Mo}=\text{S}$ bond length of 2.150(1) Å and a $\text{Mo}-\text{O}-\text{Si}$ angle of $171.4(1)^\circ$. For **5c**, the mean $\text{W}=\text{S}$ bond distance is 2.155(1) Å and the $\text{W}-\text{O}-\text{Si}$ angle is $155.2(3)^\circ$ because of the smaller steric bulk of the silyloxy substituents.



Oxosilyloxobenzenedithiolate Complexes. Monobenzenedithiolate complexes were obtained by two related methods (Figure 2). In reaction 9, bdt is introduced with the elimination of an oxo ligand as water, to yield molybdenum complexes **8a–8d** as orange solids (72–83%). This reaction is similar to the preparation of $[\text{MoO}(\text{bdt})(\mu_2\text{-S})_2\text{CuL}]^{2-}$ from $[\text{MoO}_2(\mu_2\text{-S})_2\text{CuL}]^{2-}$ and the dithiol.²² Alternatively, trioxo complexes **6** and **7** are silylated to form **8a–8d** (47–62%) and tungsten complexes **9a–9d** (green-brown or light-yellow, 42–68%), respectively, in reaction 10.



The structures of **8d** and **9b** are set out in Figure 5 with selected dimensions. These complexes present isostructural $\text{MO}_3(\text{bdt})$ fragments with distorted square-pyramidal coordination indicated by two basal angles $\text{S1}-\text{M}-\text{O3}$ and $\text{S2}-\text{M}-\text{O2}$ in the range of 138 – 152° , dihedral angles θ between MS1S2 and MO2O3 coordination planes, and displacements δ of the metal atom toward the apical oxo ligand O1. Values of the latter two parameters for **8d/9b** are $\theta = 132.1^\circ/131.0^\circ$ and $\delta = 0.56 \text{ \AA}/0.57 \text{ \AA}$. Other bond distances and angles are unexceptional. When trioxo complex **6** was treated with 2 equiv of Me_3SiCl in THF, the only isolable product was the Mo^{V} μ_2 -oxo dimer **10**. The reaction proceeds with the initial formation of **8a**. The second equiv of the silyl chloride apparently removes the silyloxy as the siloxane with

(22) Takuma, M.; Ohki, Y.; Tatsumi, K. *Inorg. Chem.* **2005**, *44*, 6034–6043.




Figure 7. UV-vis absorption spectra of 1.0 mM $[\text{Mo}^{\text{VI}}\text{O}_{3-n}\text{S}_n(\text{OSiPr}^i_3)]^{1-}$ ($n = 3$, red solid line; $n = 2$, blue solid line; $n = 1$, green broken line; $n = 0$, brown broken line) in acetonitrile solutions.

coordination of chloride. The likely reductant is bdt. As seen in Figure 6, the complex is centrosymmetric and contains the ubiquitous $\text{O}=\text{Mo}^{\text{V}}-\text{O}-\text{Mo}^{\text{V}}=\text{O}$ fragment. Coordination at the molybdenum sites is approximately square pyramidal; other metric features are normal.

Summary. The principal results of this work are summarized in terms of two main classes of compounds. Multiple examples within each class are provided to support the generality of the synthetic procedures and the scope of the stable compounds.

• $[\text{MO}_3-n\text{S}_n(\text{OSiR}_2\text{R}')]^{1-}$. Examples of all of the members of the tetrahedral $[\text{MoO}_3-n\text{S}_n(\text{OSiR}_2\text{R}')]^{1-}$ series ($n = 0-3$) have now been isolated and structurally characterized. Colors range from nearly colorless ($n = 0$) to bright red ($n = 3$), as seen by the progressive low-energy shifts of the LMCT bands in Figure 7 as the sulfur content increases. The $n = 0$ complexes **1** are obtained by silylation of Ag_2MoO_4 . These complexes undergo monosulfidation to give the $n = 1$ complexes **2**. These can be converted by further sulfidation to the $n = 2$ species **3**, one of which can be obtained by the silylation of $[\text{MoO}_3\text{S}]^{2-}$. Complexes **4** and **5** with $n = 3$ are accessible by the silylation of $[\text{MOS}_3]^{2-}$. The $n = 0$ and 3 members are preceded by three examples (**1e**,^{12,18} $[\text{MoO}_3(\text{OSiBu}^t_3)]^{1-}$,¹⁸ and **4a**¹⁶), two of which were prepared by different methods than employed here. The only previous example of an $n = 1$ complex is **2b**.¹² There has been no prior report of an $n = 2$ complex. Concerning the multiply bonded pyramidal fragment MO_3-nS_n ($M = \text{Mo}, \text{W}$) in other molecular types but excluding members of the set $[\text{MO}_4-n\text{S}_n]^{2-}$,^{12,23} there are no prior structurally characterized examples of $n = 1$ and of $n = 2$ other than $[\text{Cp}^*\text{WOS}_2]^{1-24}$

and $[(\text{Bu}^t\text{tach})\text{WOS}_2]$.²⁵ Pyramidal MS_3 units are more common, having been found in $[(\text{Bu}^t\text{tach})\text{MoS}_3]$,²⁵ $[(\text{Tp})\text{WS}_3]^{1-}$,²⁶ $[\text{Cp}^*\text{MS}_3]^{1-}$,^{27,28} and $[\text{WS}_3(\text{SR})]^{1-}$.²⁹⁻³²

• $[\text{MO}_2(\text{OSiR}_2\text{R}')(\text{bdt})]^{1-}$. The complexes **8** are accessible from **1** by the reaction with $\text{H}_2(\text{bdt})$; silylation of trioxo complexes **6** and **7** affords **8** and **9**, respectively. They are square pyramidal with an oxo ligand in the apical position. As such, they are structural analogues of the inactive protonated sites $[\text{MoO}_2(\text{OH})(\text{S}_2\text{pd})]$ of the XO family. By sulfidation with Ph_3SiSH , they are immediate precursors to square-pyramidal $[\text{WOS}(\text{OSiR}_2\text{R}')(\text{bdt})]^{1-}$ with apical oxo and basal sulfido ligation and are structural analogues of the active protonated sites $[\text{MoOS}(\text{OH})(\text{S}_2\text{pd})]$.^{11,33} Further, **8** and **9** react with thiols to give $[\text{MO}_2(\text{SR})(\text{bdt})]^{1-}$,³³ which like the molybdenum complex with $\text{R} = \text{C}_6\text{H}_2-2,4,6-\text{Pr}^i_3$,¹⁷ are analogues of the oxidized active site of sulfite oxidase.

The sulfidation reactions reported here call attention to the utility of silylthiols, here Ph_3SiSH and Pr^i_3SiSH , commercial compounds of moderate cost, as reagents for oxo-for-sulfido conversions. In the exhaustive compilation of such

(23) Müller, A.; Diemann, E.; Jostes, R.; Bögge, H. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 934-955.

(24) Kawaguchi, H.; Tatsumi, K. *Angew. Chem., Int. Ed.* **2001**, *40*, 1266-1267.

(25) Partyka, D. V.; Staples, R. J.; Holm, R. H. *Inorg. Chem.* **2003**, *42*, 7877-7886.

(26) Seino, H.; Arai, Y.; Iwata, N.; Nagao, S.; Mizobe, Y.; Hidai, M. *Inorg. Chem.* **2001**, *40*, 1677-1682.

(27) Rau, M. S.; Kretz, C. M.; Geoffroy, G. N. *Organometallics* **1993**, *12*, 3447-3460.

(28) Kawaguchi, H.; Yamada, K.; Lang, J.-P.; Tatsumi, K. *J. Am. Chem. Soc.* **1997**, *119*, 10346-10358.

(29) Boorman, P. M.; Wang, M.; Parvez, M. *J. Chem. Soc., Chem. Commun.* **1994**, 999-1000.

(30) Kruhlik, N. L.; Wang, M.; Boorman, P. M.; Parvez, M.; McDonald, R. *Inorg. Chem.* **2001**, *40*, 3141-3148.

(31) Lang, J.-P.; Kawaguchi, H.; Tatsumi, K. *J. Chem. Soc., Dalton Trans.* **2002**, 2573-2580.

(32) We note in passing that pyramidal MS_3 groups are not confined to $M = \text{Mo}, \text{W}$. Complexes of the type $[\text{MS}_3(\text{SR})]^{2-}$ ($M = \text{Nb}, \text{Ta}$) have been prepared: Coucouvanis, D.; Chen, S.-J.; Mandimutsira, B. S.; Kim, C. G. *Inorg. Chem.* **1994**, *33*, 4429-4430.

(33) Wang, J.-J. Ph.D. Thesis, Harvard University, 2007.

reactions by Donahue,³⁴ only one compound is cited for only one reaction, Ph₃SiSH in the conversion **1e** → **2b**. The minimal representation of these conversions is M^{VI}=O + Ph₃SiSH → M^{VI}=S + Ph₃SiOH. On an enthalpic basis, the reaction is favored in part by the ca. 25–40 kcal/mol difference in Si–O versus Si–S bond dissociation energies.^{35–37} Experimental M=O and M=S bond energies for M = Mo and W at constant structure and ligation are lacking. Density functional theory results for [M^{VI}QCl₄] (Q = O, S) with optimized M=Q bond lengths close to those observed in the present compounds indicate bond energy differences for M=

O versus M=S of 47 kcal/mol for M = Mo and 42 kcal/mol for M = W.³⁸ These results would appear to suggest that enthalpy changes for the minimal reaction based on bond energies are unfavorable. Experimental bond energies and reaction enthalpies would be helpful in resolving this issue.

Acknowledgment. This research was supported by NSF Grant CHE 0547734. We appreciate expert crystallographic assistance from Dr. Douglas M. Ho.

Supporting Information Available: X-ray crystallographic files in CIF format for the compounds in Tables 1 and 2. This material is available free of charge via the Internet at <http://pubs.acs.org>.

IC701294Y

(34) Donahue, J. P. *Chem. Rev.* **2006**, *106*, 4747–4783.

(35) Luo, Y.-R. *Handbook of Bond Dissociation Energies in Organic Compounds*; CRC Press: New York, 2003, pp 287–299.

(36) Basch, H. *Inorg. Chim. Acta* **1996**, *252*, 265–279.

(37) Leroy, G.; Tamsamani, D. R.; Wilante, C. *J. Mol. Struct.* **1994**, *306*, 21–39.

(38) González-Blanco, O.; Branchadell, V.; Monteyne, K.; Ziegler, T. *Inorg. Chem.* **1998**, *37*, 1744–1748.