

Metal-Mediated Transformation of a Triazinephenanthridinium Ligand Leading to a {Pd₅} Coordination Complex Observed Crystallographically and by Cryospray Mass Spectrometry

Phil J. Kitson,[†] Yu-Fei Song,[†] Patrick Gamez,[‡] Paul de Hoog,[‡] De-Liang Long,[†] Alexis D. C. Parenty,[†] Jan Reedijk,[‡] and Leroy Cronin^{*†}

WestChem Department of Chemistry, University of Glasgow, Glasgow G12 8QQ, U.K., and Leiden Institute of Chemistry, Leiden University, P.O. Box 9502, 2300 RA Leiden, The Netherlands


Received November 10, 2007

The formation of a pentanuclear palladium(II) complex with a phenanthridinonetriazine-based ligand system, which itself is formed by a metal-mediated rearrangement of a triazinephenanthridinium proligand, is described.

Triazine-based compounds are widely used, with applications ranging from medicinal chemistry,¹ herbicides,² catalysis³ and polymer chemistry.⁴ *s*-Triazine derivatives have recently shown to be of great interest in supramolecular chemistry, because of the ease with which the successive chloride substitution of 2,4,6-trichloro-1,3,5-triazine can be achieved under controlled conditions, yielding trisubstituted ligands for the construction of multinuclear coordination frameworks.⁵ We are interested in combining the flexibility of the triazine-based ligand design with a novel heterocyclic system that forms upon reaction between a primary amine⁶ and (2-bromoethyl)phenanthridinium bromide (PhenBR), which leads to the formation of a dihydroimidazophenanthridinium (DIP) heterocyclic cation (Scheme 1).^{6a}

DIPs are interesting because they exhibit tunable structures^{6b} and biological activity and have affinity for DNA as well as resistance to reduction and hydrolysis.⁷ It has been shown that the cationic nature of the phenanthridinium is

Scheme 1. Synthesis of Compound 3^a



^a Compound 2 was isolated in 90% yield and compound 3 in a 58% yield.
>>

vital to its efficacy as a potential anticancer molecule.⁸ Metal complexes of ligands that contain DNA-intercalating properties/phenanthridinium moieties have been used as DNA sensors⁹ and as possible chemotherapeutic agents.¹⁰

Herein, we describe a new ligand design approach that exploits the versatile, “modular” reactivity of 2,4,6-trichloro-1,3,5-triazine to integrate the DIP moiety, producing a ligand that could form multinuclear coordination complexes that integrate pendent-delocalized aromatic frameworks. By using this strategy as a basis, we design complexes that display interesting properties such as supramolecular self-assembly or that act as potential multivalent DNA binding agents;

* To whom correspondence should be addressed. E-mail: L.Cronin@chem.gla.ac.uk. Fax: +44 (0)141 330 4888.

[†] University of Glasgow.

[‡] Leiden University.

(1) Ronchi, S.; Prosperi, D.; Compostella, F.; Panza, L. *Synth. Lett.* **2004**, 6, 1007.

(2) Oliva, J. M.; Azenha, E. M. D. G.; Burrows, H. D.; Coimbra, R.; de Melo, J. S. S.; Canle, M. L.; Fernandez, M. I.; Santaballa, J. A.; Serrano-Andres, L. *ChemPhysChem* **2005**, 6, 306.

(3) Hu, X. P.; Chen, H. L.; Zheng, Z. *Adv. Synth. Catal.* **2005**, 347, 541.

(4) Pedroso, L. M.; Castro, M. M. C. A.; Simoes, P.; Portugal, A. *Polymer* **2005**, 46, 1766.

(5) Gamez, P.; Reedijk, J. *Eur. J. Inorg. Chem.*, **2006**, 1, 29.

(6) (a) Parenty, A. D. C.; Smith, L. V.; Pickering, A. L.; Long, D. L.; Cronin, L. *J. Org. Chem.* **2004**, 69, 5934. (b) Parenty, A. D. C.; Smith, L. V.; Guthrie, K. M.; Long, D. L.; Plumb, J.; Brown, R.; Cronin, L. *J. Med. Chem.* **2005**, 48, 4504.

(7) Nakanishi, T.; Masuda, A.; Suwa, M.; Akiyama, Y.; Hoshino-Abe, N.; Suzuki, M. *Bioorg. Med. Chem. Lett.* **2000**, 10, 2321.

(8) Zee-Cheng, R. K. Y.; Yan, S. J.; Cheng, C. C. *J. Med. Chem.* **1978**, 21, 199.

(9) Mullins, S. T.; Sammes, P. G.; West, R. M.; Yahioğlu, G. *J. Chem. Soc., Perkin Trans.* **1996**, 1, 75.

(10) Whittaker, J.; McFayden, W. D.; Baguley, B. C.; Murray, V. *Anti-Cancer Drug Des.* **2001**, 16, 81.





Figure 1. Ball-and-stick representation of $[\text{Pd}_5(\text{L})_2(\text{OAc})_8]^{2+}$. The solvent and anions of crystallization, as well as the H atoms, are omitted for clarity: C, gray; N, blue; O, red; Pd, green. Pd3 rests on an inversion center.

Scheme 2. Transformation of **3** into **L** upon Coordination to $\text{Pd}(\text{OAc})_2$ in DCM via a Ring-Opening Process, Occurring in a 33% Yield



therefore, a new ligand, compound **3**, was designed and synthesized. This compound integrates one DIP moiety along with two bidentate binding sites (Scheme 1). As shown in Scheme 1, 2-chloro[4,6-(dipyridin-2-ylamino)]-1,3,5-triazine¹¹ was coupled with ethylenediaminedihydroimidazophenanthridinium by nucleophilic substitution on the 2 position of the triazine; as a result, compound **3** is obtained with a good yield. The reaction of 2 equiv of palladium(II) acetate with 1 equiv of compound **3** in dichloromethane (DCM) yields light-yellow, needle-shaped single crystals of compound **4** after 2 weeks, whose crystal structure reveals that the initial compound **3** has been hydrolyzed upon coordination to give a new ligand, **L** (Scheme 2).

The structure of compound **4**, which crystallizes in the space group $P\bar{1}$, can be formulated as a pentanuclear Pd^{II} complex containing two of the new ligands, **L**, with a formula of $[\text{Pd}_5(\text{L})_2(\text{OAc})_8](\text{Br})_2$, and the structure is shown in Figure 1. The structure contains three different Pd^{II} ion positions identified as Pd1, Pd2, and Pd3. The coordination




Figure 2. Perspective view showing the one-dimensional supramolecular chains assembled by means of π - π stacking interactions. The mean distance d between the two phenanthridine planes amounts to 3.391 Å.

geometries around all Pd atoms are square-planar, typical for d^8 Pd^{II} centers. Pd1 and Pd2 both have a N_2O_2 donor set resulting from the coordination of two pyridine units and two acetato ligands but with minor deviations. The structure is well-defined, but there is some disorder relating to the position of the two bromide anions that compensate for the charge of the cationic species (Figure 1). In addition, complex **4** crystallizes with water and DCM molecules in the crystal lattice.

The Pd-N and Pd-O bond lengths are in normal ranges, as compared with analogous complexes.¹² Pd3 is coordinated by four amine N atoms, belonging to two different **L** ligands, thus connecting two Pd1/Pd2 entities and generating the title pentanuclear compound (Figure 1). The coordinating ethylenediamine moieties are formed through the alteration of the original compound **3** upon reaction with palladium(II) acetate. The Pd-N bond distances are typical for a square-planar $\text{Pd}(\text{N}_{\text{amine}})_4$ coordination environment involving bidentate amine ligands.¹³ The angles around the Pd^{II} ion reflect a significant distortion from the ideal square-planar geometry, most likely as the result of the small bite angle of the chelating diethylenediamine group ($\text{N}_4\text{-Pd}_3\text{-N}_8 = 80.21^\circ$), combined with steric hindrance due to the phenanthridines.

The crystal packing of **4** shows interesting supramolecular features. The pentanuclear units are connected to two neighbors via π - π stacking interactions (the distance between the aromatic rings amounts to ca. 3.4 Å), giving rise to a one-dimensional chain (Figure 2). These supramolecular chains are further stabilized through $\text{C-H}\cdots\pi$ binding contacts ($d_{\text{H-arene centroid}} = 2.694$ Å), between phenanthridine H atoms and the triazine rings of adjacent pentanuclear complexes. Such $\text{C-H}\cdots\pi$ interactions with the *s*-triazine ring have been reported.¹⁴ It is interesting to note that these electron-deficient arenes appear to be interacting with the

(11) de Hoog, P.; Gamez, P.; Driessen, W. L.; Reedijk, J. *Tetrahedron Lett.* **2002**, *43*, 6783.

(12) (a) Seward, C.; Jia, W. L.; Wang, R. Y.; Wang, S. *Inorg. Chem.*, **2004**, *43*, 978. (b) Okabe, N.; Mizubayashi, Y.; Odoko, M. *Acta Crystallogr., Sect. E* **2006**, *62*, 2747.

(13) (a) Macdougall, J. J.; Nelson, J. H.; Fultz, W. C.; Burmeister, J. L.; Holt, E. M.; Alcock, N. W. *Inorg. Chim. Acta* **1982**, *63*, 75. (b) Newkome, G. R.; Frere, Y. A.; Fronczek, F. R.; Gupta, V. K. *Inorg. Chem.* **1985**, *24*, 1001.




Figure 3. Cryospray mass spectrum (negative ion mode) of compound **4** in a DCM/MeCN (1:1) mixture at $-40\text{ }^{\circ}\text{C}$ shown in black. The simulated spectrum for $[\text{Pd}_5(\text{C}_{40}\text{H}_{34}\text{N}_{12}\text{O})_2(\text{CH}_3\text{CO}_2)_5\text{Cl}_6]^-$ is shown in red.


chlorine lone pairs¹⁵ ($d_{\text{Cl-arene centroid}} = 3.503\text{ \AA}$) from the DCM solvent (lp), which can be formulated as a type of $\text{C-H}\cdots\pi\cdots\text{lp}$ motif; although this is only a crystallographic observation, a search of CSD indicates that this observation is unique. Furthermore, the interplay between cation- π , π - π , and anion- π interactions^{14c} has been clearly demonstrated by density functional theory calculations, which have predicted that such interactions could exist.^{14b} Therefore, we tentatively suggest that the present system corresponds to the first crystallographic example of a $\text{C-H}\cdots\pi\cdots\text{lp}$ (which can also be symbolized as a $\delta+\cdots\pi\cdots\delta-$ supramolecular complex, corroborating the recent results of Quinonero et al.¹⁶

Solution investigations of **4** show that the Pd_5L_2 core of compound **4** can be observed using cryospray mass spectrometry as the $[\text{Pd}_5(\text{C}_{40}\text{H}_{34}\text{N}_{12}\text{O})_2(\text{CH}_3\text{CO}_2)_5\text{Cl}_6]^-$ anion, where the chloride anions originate from NaCl added to the solvent mixture (see Figure 3).

L forms from compound **3** (Scheme 2) via a ring-opening process with the introduction of a carbonyl group to give the ligand observed in the crystal structure upon coordination to $\text{Pd}(\text{OAc})_2$. While the precise mechanism of this ligand transformation has to await detailed experimental and theoretical investigation, one's first thought turns to a hydrolytic process during coordination, as shown in Scheme 3.

It is proposed that the Pd^{II} cation acts as a Lewis acid, which, upon coordination to the DIP N atom, activates the α position of the phenanthridinium moiety to nucleophilic attack from ambient water molecules in the reaction solvents. Elimination of HBr from this intermediate results in the formation of a neutral hydroxydihydroimidazophenanthridinium pseudobase molecule, which then undergoes a ring-opening rearrangement to form the crystallographically observed phenanthridinone moiety. This ring opening sig-

Scheme 3. Proposed Mechanism for the Ligand Transformation upon Coordination of Compound **3** by $\text{Pd}(\text{OAc})_2$



nificantly reduces the steric bulk around the Pd^{II} coordination environment, which may contribute to the driving force for this reaction. Preliminary NMR experiments (see the Supporting Information) show that the first step of this process is that compound **3** coordinates to Pd^{II} ion, after which hydrolysis by water completes the transformation.

In conclusion, a new triazine-based building block linked by a phenanthridinone molecule through an ethylenediamine unit has been designed and synthesized successfully. A novel pentanuclear Pd^{II} complex has been obtained by the reaction of compound **3** with $\text{Pd}(\text{OAc})_2$, in which a Pd^{II} -mediated ligand transformation from compound **3** to a phenanthridinone-based ligand in compound **4** has been observed. This observation represents the potential for the development of a new route to the synthesis of N-derivatized phenanthridinone moieties under mild conditions. The possible observation of a $\text{C-H}\cdots\pi\cdots\text{lp}$ interaction is interesting,¹⁷ and we will investigate this further with both theoretical and other experiments to gather further evidence giving insight into the nature of this interaction.

Acknowledgment. We thank the EPSRC and University of Glasgow for funding of this work. Support by the Graduate Research School Combination "NRSC Catalysis", a joint activity of the graduate research schools NIOK, HRSMC, and PTN is appreciated.

Supporting Information Available: Full synthetic information and characterization of compounds **1–3** and complex **4**, CIF file of compound **4**, along with mass spectra and initial NMR experiments on ligand transformation. This material is available free of charge via the Internet at <http://pubs.acs.org>.

IC7022192

- (14) (a) Thalladi, V. R.; Boese, R.; Brasselet, S.; Ledoux, I.; Zyss, J.; Jetli, R. K.; Desiraju, G. R. *Chem. Commun.* **1999**, 17, 1639. (b) Gamez, P.; Mooibroek, T. J.; Teat, S. J.; Reedijk, J. *Acc. Chem. Res.* **2007**, *40*, 435. (c) Frontera, A.; Quinonero, D.; Costa, A.; Ballester, P.; Deya, P. M. *New J. Chem.* **2007**, *4*, 556.
- (15) Lu, Z.; Gamez, P.; Mutikainen, I.; Turpeinen, U.; Reedijk, J. *Cryst. Growth Des.* **2007**, *7*, 1669–1671.
- (16) Quinonero, D.; Frontera, A.; Garau, C.; Ballester, P.; Costa, A.; Deya, P. M. *ChemPhysChem.* **2006**, *7*, 2487.

- (17) DiLabio, G. A.; Johnson, E. R. *J. Am. Chem. Soc.* **2007**, *129*(19), 6199.