

Reaction Chemistry of the U³⁺ Metallocene Amidinate (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr]U Including the Isolation of a Uranium Complex of a Monodentate Acetate

William J. Evans,* Justin R. Walensky, and Joseph W. Ziller


Department of Chemistry, University of California Irvine, California 92697-2025

Received October 28, 2009

The reductive chemistry of U³⁺ in the metallocene amidinate coordination environment of (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']U, **1**, has been explored. Two equivalents of **1** react with PhSSPh and 2,2'-dithiopyridine (pySSpy) to produce (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']U(SPh), **2**, and (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']U(Spy), **3**, respectively. Complexes **2** and **3** can also be synthesized through insertion of ⁱPrN=C=NⁱPr into the methyl group in (C₅Me₅)₂UMe(SPh) and (C₅Me₅)₂UMe(Spy), **4**, respectively. Complex **1** readily reduces the Cu¹⁺ reagents, CuBr, CuI, and CuO₂CMe, to produce the corresponding (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']UX complexes (X=Br, **5**; I, **6**; O₂CMe, **7**). X-ray crystallography established complex **7** as the first f element complex containing a monodentate acetate anion. Complex **7** can also be obtained by reaction of (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']U with CO₂ at 80 psi. In contrast to the reactions above, **1** reduces TiC₅H₅ with the unusual loss of (C₅Me₅)⁻ to form (C₅Me₅)(C₅H₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']U, **8**.

Introduction

Recent studies of carbodiimide insertion with the actinide metallocene dialkyl complexes (C₅Me₅)₂AnMe₂ (An = U, Th) have produced monoalkyl products (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']AnMe, **1**, eq 1, that contain an ancillary ligand set, {(C₅Me₅)₂[ⁱPrNC(Me)NⁱPr]}³⁻, alternative to the



heavily studied (C₅Me₅)₂⁻ organoactinide coordination environment. This ligand set allows for examination of U-element bond reactivity in a metallocene complex with one “action” ligand, X or R, rather than the two found in (C₅Me₅)₂AnX₂ and (C₅Me₅)₂AnR₂ complexes.²

Complexes of U⁴⁺ with this heteroleptic cyclopentadienyl amidinate ligand set were found to undergo reduction to form the U³⁺ complex, (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']U, **1**. Specifically, {(C₅Me₅)₂[ⁱPrNC(Me)NⁱPr-κ²N,N']U}{BPh₃Me}

*To whom correspondence should be addressed. E-mail: wevans@uci.edu. Fax: 949-824-2210.

(1) Evans, W. J.; Walensky, J. R.; Ziller, J. W.; Rheingold, A. L. *Organometallics* **2009**, *28*, 3350.

(2) (a) Fagan, P. J.; Manriquez, J. M.; Maatta, E. A.; Seyam, A. M.; Marks, T. J. *J. Am. Chem. Soc.* **1981**, *103*, 6650. (b) Ephritikhine, M. *Dalton Trans.* **2006**, 2501. (c) Sharma, M.; Eisen, M. S. *Struct. Bonding (Berlin)* **2008**, *127*, 1.

reacts with KC₅Me₅ to form **1**.³ Complex **1** provides an opportunity to explore reductive U³⁺ chemistry⁴ with a new ligand set and to expand the range of uranium compounds with {(C₅Me₅)₂[ⁱPrNC(Me)NⁱPr]}³⁻ ligation. Representative reactions of the reductive chemistry of **1** with inorganic substrates are reported here including the isolation of the first monodentate acetate complex of uranium and an unusual (C₅Me₅)⁻ displacement reaction.⁵ As emphasized in several recent reviews,⁶ exploration of new coordination environments and identification of new coordination modes and reactivity patterns are essential to fully defining the fundamental chemistry of the actinides.

Experimental Section

The syntheses and manipulations described below were conducted with rigorous exclusion of air and water using

(3) Evans, W. J.; Walensky, J. R.; Ziller, J. W. *Chem.—Eur. J.* **2009**, *15*, 12204.

(4) (a) Cloke, F. G. N.; Hitchcock, P. B. *J. Am. Chem. Soc.* **2002**, *124*, 9352. (b) Castro-Rodriguez, I.; Olsen, K.; Gantzel, P.; Meyer, K. *J. Am. Chem. Soc.* **2003**, *125*, 4565. (c) Karmazin, L.; Mazzanti, M.; Bezombes, J.-P.; Gateau, C.; Pecaut, J. *Inorg. Chem.* **2004**, *43*, 5147. (d) Evans, W. J.; Kozimor, S. A. *Coord. Chem. Rev.* **2006**, *250*, 911. (e) Korobkov, I.; Gorelsky, S.; Gambarotta, S. *J. Am. Chem. Soc.* **2009**, *131*, 10406.

(5) See for example: (a) Evans, W. J.; Kozimor, S. A.; Ziller, J. W.; Kaltsoyannis, N. *J. Am. Chem. Soc.* **2004**, *126*, 14533. (b) Evans, W. J.; Mueller, T. J.; Ziller, J. W. *J. Am. Chem. Soc.* **2009**, *131*, 2678.

(6) (a) Burns, C. J.; Eisen, M. S. In *The Chemistry of the Actinide and Transactinide Elements*; Morss, L. R., Edelstein, N. M., Fuger, J., Eds.; Springer: Dordrecht, The Netherlands, **2006**; Vol. 5, p 2911. (b) Fox, A. R.; Bart, S. C.; Meyer, K.; Cummins, C. C. *Nature* **2008**, *455*, 341. (c) Ephritikhine, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 4998.

Schlenk, vacuum line, and glovebox techniques. All reactions were conducted in a Vacuum Atmospheres inert-atmosphere (Ar) glovebox free of coordinating solvents. Solvents were sparged with UHP argon, dried by passage through columns containing Q-5 and molecular sieves, and delivered directly to the glovebox through stainless steel tubing. All reactions were conducted in an Ar glovebox free of coordinating solvents. Benzene- d_6 (Cambridge Isotope Laboratories) was dried over NaK alloy and benzophenone, degassed by three freeze-pump-thaw cycles, and vacuum transferred before use. $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$, **1**,³ $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ Me, **1** and $(C_5Me_5)_2U$ Me(SPh)⁷ were prepared as previously described. 2,2'-Dithiodipyridine, CuSPh, CuBr, CuI, CuO₂CMe (Aldrich) and cyclopentadienylium (Strem, 99% sublimed) were used as received. ${}^iPrN=C=N^iPr$ (Aldrich) was dried over molecular sieves and degassed by three freeze-pump-thaw cycles. PhSSPh (Aldrich) was sublimed prior to use. CO₂ (Airgas) was used as received. NMR experiments were conducted with Bruker 400 and 500 MHz spectrometers. Because of the paramagnetism of uranium, only resonances that could be unambiguously assigned are reported.⁸ Infrared spectra were recorded as KBr pellets on a Perkin-Elmer Spectrum One FT-IR spectrometer. Elemental analyses were performed by Analytische Laboratorien (Lindlar, Germany) or with a Perkin-Elmer 2400 CHN elemental analyzer.

Synthesis of $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ (SPh), **2.** **Method A.** PhSSPh (29 mg, 0.13 mmol) was added to a stirred solution of **1** (175 mg, 0.269 mmol) in hexanes (6 mL). After 12 h, the solvent was removed, and **2** was isolated as a red powder (163 mg, 80%). Crystals suitable for X-ray diffraction were grown from a saturated pentane/toluene solution at $-35^\circ C$. ¹H NMR (C_6D_6 , 298 K): δ 13.0 (s, 6H, CHMe₂), 9.4 (s, 1H, CHMe₂), 5.3 (s, 30H, C₅Me₅), 1.3 (s, 3H, Me), 0.4 (s, 1H, CHMe₂). IR: 3058s, 2980s, 1648w, 1578 m, 1489 m, 1435 m, 1342s, 1196s, 1141s, 1084s, 1058 m, 1024s, 789s, 737s cm⁻¹. Anal. Calcd. for C₃₄H₅₂N₂SU: C, 53.81; H, 6.91; N, 3.69. Found: C, 53.81; H, 7.14, N, 3.19.

Method B. Complex **2** can also be prepared by addition of ${}^iPrN=C=N^iPr$ (35 μ L, 0.23 mmol) to a stirred solution of $(C_5Me_5)_2U$ Me(SPh) (160 mg, 0.225 mmol) in hexanes (8 mL). After 1 h, the solvent was removed under vacuum to yield **2** as a microcrystalline red solid (181 mg, 94%).

Method C. Reaction of **1** (20 mg, 0.031 mmol) with CuSPh (6 mg, 0.03 mmol) in an NMR tube showed the quantitative conversion to **2** by ¹H NMR spectroscopy.

$(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ (Spy), **3**. **Method A.** 2,2'-Dithiodipyridine (18 mg, 0.082 mmol) was added to a stirred solution of **1** (97 mg, 0.149 mmol) in methylcyclohexane (10 mL). The color turned from green-brown to orange. After 12 h, the solvent was removed under vacuum to yield an orange oil. Crystallization from a saturated pentane solution at $-35^\circ C$ gave orange crystals of **3** (74 mg, 66%). ¹H NMR (C_6D_6 , 298 K): δ 28.4 (s, 1H, py), 12.3 (s, 30H, C₅Me₅), 4.8 (t, 1H, py), 4.0 (s, 1H, CHMe₂), -7.6 (s, 1H, py), -9.7 (s, 3H, Me). IR: 2963s, 1585s, 1543 m, 1513s, 1443s, 1413s, 1378s, 1260 m, 1195s, 1137s, 1087 m, 785s, 756s, 729s cm⁻¹. Anal. Calcd for C₃₃H₅₁N₃SU: C, 52.44; H, 6.80; N, 5.56. Found: C, 52.15; H, 6.87; N, 5.96.

Method B. Complex **3** can also be prepared from $(C_5Me_5)_2U$ Me(Spy), **4**, described below. In an NMR tube, 15 mg of **4** and 5 μ L of ${}^iPrN=C=N^iPr$ were combined. The quantitative conversion of **4** to **3** was observed by ¹H NMR spectroscopy.

$(C_5Me_5)_2U$ Me(Spy), **4**. After addition of pySSpy (165 mg, 0.749 mmol) to a stirred solution of $(C_5Me_5)_2U$ Me₂ (203 mg, 0.377 mmol) in toluene (10 mL), the solution was allowed to stir

for 16 h. The solvent was then removed under vacuum to yield a red-orange oil. Upon crystallization from a saturated hexanes solution, **4** was obtained as orange crystals suitable for X-ray diffraction (175 mg, 85%). ¹H NMR (C_6D_6 , 298 K): δ 15.1 (s, 3H, Me), 8.2 (d, 1H, py), 7.4 (d, 1H, py), 6.8 (t, 1H, py), 6.3 (t, 1H, py), 2.8 (s, 15H, C₅Me₅), 1.8 (s, 15H, C₅Me₅). Anal. Calcd for C₂₆H₃₇NSU: C, 49.28; H, 5.89; N, 2.21. Found: C, 49.52; H, 5.99; N, 2.13.

$(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ Br, **5**,⁹ **from 1 and CuBr.** In an NMR tube, CuBr (6 mg, 0.04 mmol) was added to a solution of **1** (20 mg, 0.031 mmol) in C_6D_6 (0.5 mL). After 3 h, the ¹H NMR resonances corresponding to **1** disappeared, and the previously characterized **5** was the only observable product with the formation of a dark precipitate. ¹H NMR (C_6D_6 , 298 K): δ 5.8 (s, 30H, C₅Me₅), -5.1 (s, 3H, Me). IR: 2931s, 2893s, 1642 m, 1476s, 1442s, 1418s, 1383s, 1347s, 1339s, 1203s, 1143s, 1126s, 1047 m, 1019 m, 993s, 790 m cm⁻¹. Anal. Calcd for C₂₈H₄₇N₂BrU: C, 46.09; H, 6.49; N, 3.84. Found: C, 46.15; H, 6.68; N, 3.76.

$(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ I, **6**,⁹ **from 1 and CuI.** In an NMR tube, CuI (6 mg, 0.03 mmol) was added to a solution of **1** (20 mg, 0.031 mmol) in C_6D_6 (0.5 mL). Over 3 h, the ¹H NMR resonances corresponding to **1** disappeared, and the previously characterized **6** was the only observable product with the formation of a dark precipitate. ¹H NMR (C_6D_6 , 298 K): δ 8.3 (s, 30H, C₅Me₅), -4.8 (s, 3H, Me). IR: 2927s, 2896s, 2720 m, 1638w, 1494s, 1436s, 1414s, 1379s, 1359s, 1349s, 1195s, 1140s, 1126s, 1055 m, 1018 m, 1000 m, 808w, 790 m cm⁻¹. Anal. Calcd for C₂₈H₄₇N₂IU: C, 43.30; H, 6.10; N, 3.61; I, 16.34; U, 30.65. Found: C, 43.73; H, 5.68; N, 3.34; I, 16.03; U, 30.84.

$(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ (O₂CMe), **7**. **Method A.** CuO₂CMe (34 mg, 0.28 mmol) was added to a stirred solution of **1** (123 mg, 0.189 mmol) in toluene (8 mL). After 16 h, insoluble material was removed by centrifugation, and the solvent was removed under vacuum to yield **7** as a red-brown oil. Upon extraction with hexanes and removal of solvent, **7** was obtained as a red-brown powder (114 mg, 85%). Crystals suitable for X-ray crystallography were grown from a saturated pentane/toluene solution at $-35^\circ C$. ¹H NMR (C_6D_6 , 298 K): δ 32.0 (s, 3H, Me), 27.9 (s, 6H, CHMe₂), 3.9 (s, 1H, CHMe₂), 0.04 (s, 30H, C₅Me₅), -9.8 (s, 3H, Me), -19.7 (s, 6H, CHMe₂). IR: 2955s, 2859s, 1657s, 1584s, 1514s, 1436 m, 1419s, 1376s, 1356s, 1197s, 1130s, 1007 m, 922 m, 791 m, 677 m cm⁻¹. Anal. Calcd for C₃₀H₅₀N₂O₂U: C, 52.37; H, 8.05; N, 3.70. Found: C, 51.93; H, 7.94; N, 3.88.

Method B. In a glovebox, $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]U$ Me (150 mg, 0.226 mmol) was dissolved in toluene (5 mL) and placed into a Fischer-Porter vessel. The reaction vessel was attached to a high pressure line and CO₂ (80 psi) was introduced. Over 24 h, the color of the mixture changed from yellow-brown to orange. The reaction vessel was transferred to a glovebox where the solvent removed under vacuum to yield **7** (identified by ¹H NMR spectroscopy) as an orange microcrystalline solid in quantitative yield.

$(C_5Me_5)(C_5H_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$, **8**. TIC₅H₅ (48 mg, 0.31 mmol) was added to a stirred solution of **1** (98 mg, 0.15 mmol) in toluene (8 mL). The solution turned from green to red after several minutes. After 12 h, insoluble material was removed by centrifugation, and the solvent was removed under vacuum to yield **8** as a red powder (80 mg, 82%). Crystals suitable for X-ray crystallography were grown from a saturated toluene solution at $-35^\circ C$. ¹H NMR (C_6D_6 , 298 K): δ 4.5 (s, 6H, CHMe₂), 2.3 (s, 6H, CHMe₂), 2.2 (s, 15H, C₅Me₅), -11.3 (s, 10H, C₅H₅). IR: 2971s, 1469s, 1440 m, 1377s, 1359s, 1199s, 1142s, 1060 m, 1011s, 995s, 903w, 791s, 716s, 693s cm⁻¹. Anal. Calcd for C₂₈H₄₂N₂U: C, 52.17; H, 6.37; N, 4.21. Found: C, 52.38; H, 6.49; N, 4.10.

X-ray Data Collection, Structure Solution, and Refinement for 2-4, 7, and 8. This information is available in the Supporting

(7) Evans, W. J.; Miller, K. A.; Ziller, J. W.; DiPasquale, A. G.; Heroux, K. J.; Rheingold, A. L. *Organometallics* **2007**, *26*, 4287.

(8) Gaunt, A. J.; Scott, B. L.; Neu, M. P. *Inorg. Chem.* **2006**, *45*, 7401.

(9) Evans, W. J.; Walensky, J. R.; Ziller, J. W. *Organometallics* **2010**, *29*, 101.

Table 1. X-ray Data Collection Parameters for Complexes **2–4**, **7**, and **8**


	2	3	4	7	8
empirical formula	C ₃₄ H ₅₂ N ₂ SU	C ₃₃ H ₅₁ N ₃ SU • 1/2(C ₇ H ₈)	C ₂₆ H ₃₇ NSU	C ₃₀ H ₅₀ N ₂ O ₂ U	C ₂₈ H ₄₂ N ₂ U
formula weight	758.87	805.93	633.66	708.75	644.67
temperature (K)	103(2)	103(2)	153(2)	143(2)	103(2)
crystal system	monoclinic	monoclinic	monoclinic	monoclinic	triclinic
space group	<i>P</i> 2 ₁ / <i>c</i>	<i>C</i> 2/ <i>c</i>	<i>P</i> 2 ₁	<i>P</i> 2 ₁ / <i>n</i>	<i>P</i> $\bar{1}$
<i>a</i> (Å)	11.5156(6)	34.4356(18)	8.343(3)	9.5795(6)	8.2845(4)
<i>b</i> (Å)	17.1703(9)	12.3473(7)	15.068(5)	17.4341(11)	11.1001(6)
<i>c</i> (Å)	16.0926(8)	16.1941(9)	9.730(3)	17.8520(11)	14.2310(7)
α (deg)	90	90	90	90	81.9930(5)
β (deg)	90.2320(10)	90.6820(10)	96.247(4)	101.4380(10)	76.1897(5)
γ (deg)	90	90	90	90	81.9746(6)
volume Å ³	3181.9(3)	6885.0(7)	1216.0(7)	2992(9)	1250.72(11)
<i>Z</i>	4	8	2	4	2
ρ_{calcd} (Mg/m ³)	1.584	1.555	1.731	1.611	1.712
μ (mm ⁻¹)	5.191	4.804	6.772	5.582	6.506
R1 [<i>I</i> > 2.0 σ (<i>I</i>)] ^a	0.0295	0.0181	0.0431	0.0240	0.0166
wR2 (all data) ^a	0.0622	0.0468	0.1111	0.0604	0.0427

^a Definitions: $wR2 = [\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)]^{1/2}$, $R1 = \sum ||F_o| - |F_c|| / \sum |F_o|$.


Information. Selected X-ray collection parameters are found in Table 1.

Results

RSSR Reactions. PhSSPh is one of the substrates examined with new f element reducing agents since the S–S bond is readily reduced ($E = -1.7$ vs SCE)¹⁰ and the (SPh)[−] ligands formed on reduction frequently form isolable complexes.¹¹ Two equivalents of **1** reduce PhSSPh to produce the U⁴⁺ complex (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr- κ^2 N,N']U(SPh), **2**, eq 2. Complex **2** was characterized by analytical and spectroscopic methods, but did not initially



crystallize to allow identification by X-ray crystallography. To provide an extra coordination option, the pyridyl (py) analogue, pySSpy, was treated with **1** and (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr- κ^2 N,N']U(Spy), **3**, was isolated, eq 3.



Subsequently, both **2** and **3** were obtained in crystalline form suitable for X-ray crystallography, Figures 1 and 2. Hence, the (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr]^{3−} ligand set can




Figure 1. Thermal ellipsoid plot of (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr- κ^2 N,N']U(SPh), **2**, shown at the 30% probability level. Hydrogen atoms have been omitted for clarity.

accommodate both a mono- and bidentate arylsulfide ligand. Table 2 compares the metrical parameters of these closely related complexes.

Complex **2** displays U–(C₅Me₅ ring centroid) distances of 2.510 and 2.515 Å that are between the 2.498 and 2.500 Å distances in (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr]U¹ and the 2.527 and 2.538 Å in (C₅Me₅)₂[ⁱPrNC(Me)NⁱPr]U^{Me}.¹ The 2.7665(9) Å U–S bond is longer than the 2.7060(14) Å distance in (C₅Me₅)₂U^{Me}(SPh)⁷ and the 2.6845(7) and 2.6967(7) Å lengths in (C₅Me₄H)₂U(SPh)₂¹² as might be expected since the [ⁱPrNC(Me)NⁱPr][−] ligand is larger than (Me)[−] and (SPh)[−].

In complex **3**, the 2.565 and 2.608 Å U–(C₅Me₅ ring centroid) distances are longer than those in **2** and in any previously characterized U⁴⁺ metallocene amidinate complexes.^{1,3,9} This is consistent with the increased coordination number. Interestingly, the 2.408(1) and 2.510(1) Å U–N (amidinate) bond distances and

(10) Bradbury, J. R.; Masters, A. F.; McDonell, A. C.; Brunette, A. A.; Bond, A. M.; Wedd, A. C. *J. Am. Chem. Soc.* **1981**, *103*, 1959.

(11) See for example: (a) Diaconescu, P. L.; Arnold, P. L.; Baker, T. A.; Mindiola, D. J.; Cummins, C. C. *J. Am. Chem. Soc.* **2000**, *122*, 6108. (b) Graves, C. R.; Scott, B. L.; Morris, D. E.; Kiplinger, J. L. *Chem. Commun.* **2009**, 776.

(12) Evans, W. J.; Miller, K. A.; Hillman, W. R.; Ziller, J. W. *J. Organomet. Chem.* **2007**, *692*, 3649.




Figure 2. Thermal ellipsoid plot of $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N^i]U(Spy)$, **3**, shown at the 50% probability level. Hydrogen atoms have been omitted for clarity.

Table 2. Selected Bond Lengths [Å] and Angles [deg] for $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N^i]U(SPh)$, **2**, and $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N^i]U(Spy)$, **3**

bond distance/angle	2	3
U(1)–(C ₅ Me ₅ ring centroid)	2.510, 2.515	2.565, 2.608
U(1)–N(1)	2.482(3)	2.510(2)
U(1)–N(2)	2.380(3)	2.408(2)
U(1)–C(21)	2.902(3)	2.926(2)
U(1)–N(3)		2.572(2)
U(1)–S(1)	2.7665(9)	2.7997(6)
(C ₅ Me ₅ ring centroid)–U(1)– (C ₅ Me ₅ ring centroid)	129.7	118.7
(C ₅ Me ₅ ring centroid)–U(1)–N(1)	101.1, 100.1	93.4, 146.2
(C ₅ Me ₅ ring centroid)–U(1)–N(2)	122.3, 108.9	109.0, 101.9
(C ₅ Me ₅ ring centroid)–U(1)–N(3)		146.4, 89.1
(C ₅ Me ₅ ring centroid)–U(1)–S(1)	91.2, 102.4	95.3, 101.7
U(1)–S(1)–C(29)	126.46(18)	84.24(8)
N(1)–U(1)–N(2)	54.41(10)	53.84(6)

2.7997(6) Å U–S bond length in **3** are not dissimilar to the distances seen in **2**: 2.380(3)/2.482(3) Å and 2.7996(9) Å, respectively. Two other U⁴⁺ complexes bearing the thio-pyridine ligand are known, U(Spy)₄(THF)⁸ and [U{(SiMe₂NPh)₃-tacn}(Spy)]¹³ but comparisons are more complicated since they are not metallocenes. The 2.572(1) Å U–N(3) bond in **3** is similar to the 2.504–(2)–2.575(2) Å U–N(py) analogues in these compounds as is the 2.7997(6) Å U–S distance (2.8222(8)–2.880(2) Å in the other compounds).

Since complex **3** exhibited long U–(C₅Me₅ ring centroid) distances compared to the other metallocene amidinate complexes, the displacements of the methyl groups from the plane of the (C₅Me₅)[–] ligand were measured. Displacements of this type have been used as a calibration of steric crowding in f element complexes.¹⁴ Displacements of 0.27, 0.21, 0.43, 0.11, and 0.22 Å were found for C(6)–C(10), and values of 0.28, 0.33, 0.12, 0.52, and 0.32 Å were measured for C(16)–C(20), respectively. The 0.52 Å value measured for C(19), Figure 3, is in the range of extreme values displayed by sterically crowded (C₅Me₅)₃M complexes





Figure 3. Two methyl groups with the greatest displacement from the cyclopentadienyl ring plane, C(8) and C(19), in complex **3** are shown. Thermal ellipsoids are shown at the 50% probability level.


that display unusual (C₅Me₅)[–] reactivity like sterically induced reduction (SIR).^{14,15}

Although **3** has a methyl group with an extreme displacement, it does not display unusual (C₅Me₅)[–] reactivity. For example, it does not reduce phenazine even at 100 °C in toluene. This is consistent with the finding that the sterically crowded complexes that have SIR reactivity are globally crowded at all the ligand sites.¹⁶

Since **2** could be formed by insertion of ⁱPrN=C=NⁱPr into the methyl group of the U⁴⁺ complex, (C₅Me₅)₂–U–Me(SPh),⁷ this reaction was examined and a second route to **2** was identified, eq 4. To determine if **3** could also form via insertion of a carbodiimide, the necessary



precursor, (C₅Me₅)₂U–Me(Spy), **4**, was synthesized by the reaction of (C₅Me₅)₂U–Me₂ with pySSpy, eq 5. The byproduct of this σ bond metathesis reaction, MeSpy,



was identified by ¹H NMR spectroscopy.¹⁷ The structure of **4** was established by X-ray crystallography, Figure 4, although the data were not of high enough quality for a detailed discussion. As shown in eq 6, insertion of a carbodiimide into the U–Me bond in **4** is also successful and forms **3**. Hence, the extra coordination of the pyridyl

(13) Antunes, M. A.; Dias, M.; Monteiro, B.; Domingos, A.; Santos, I. C.; Marques, N. *Dalton Trans.* **2006**, 3368.

(14) Evans, W. J.; Kozimor, S. A.; Ziller, J. W. *Inorg. Chem.* **2005**, *44*, 7960.

(15) Evans, W. J.; Davis, B. L. *Chem. Rev.* **2002**, *102*, 2119.

(16) Evans, W. J.; Walensky, J. R.; Furche, F.; Ziller, J. W.; DiPasquale, A. G.; Rheingold, A. L. *Inorg. Chem.* **2008**, *47*, 10169.

(17) Doudouh, A.; Woltermann, C.; Gros, P. C. *J. Org. Chem.* **2007**, *72*, 4978.






Figure 4. Thermal ellipsoid plot of $(C_5Me_5)_2U(Me)(Spy)$, **4**, shown at the 30% probability level. Hydrogen atoms have been omitted for clarity.

substituent in **4** versus $(C_5Me_5)_2U(Me)(SPh)$ did not prevent this insertion from occurring.



Attempts to reduce $1,3,5,7-C_8H_8$ ($E = -1.85$ V and -1.9 V vs SCE),¹⁸ with **1** were unsuccessful even at $100^\circ C$ in toluene. Although the reductions of PhSSPh ($E = -1.7$ V vs SCE)⁹ and pySSpy were facile, complex **1** showed no reactivity with phenazine ($E = -0.36$ V vs SCE),¹⁹ benzaldehyde azine,²⁰ and benzophenone.²¹

Reactions of **1** with copper salts were also of interest since Kiplinger and co-workers recently showed that it was possible to convert U^{3+} precursors to U^{4+} halides and pseudohalides.²² Complex **1** reacts similarly with CuSPh, CuBr, and CuI to form **2**, $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']UBr$, **5**,⁹ and $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']UI$, **6**,⁹ eq 7, respectively. Complexes **5** and **6** were previously synthesized by reaction of $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ Me with CuBr and CuI, respectively.⁹ The previous synthesis is a more facile route to **5** and **6** than eq 7, since $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ Me is directly obtainable from $(C_5Me_5)_2U$ Me₂, eq 1.¹



The reaction of **1** with copper acetate provided the desired result. Complex **1** reacts with CuO_2CMe to make the expected acetate complex, $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-$






Figure 5. Thermal ellipsoid plot of $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U(O_2CMe)$, **7**, shown at the 30% probability level. Hydrogen atoms have been omitted for clarity.

$\kappa^2N,N']U(O_2CMe)$, **7**, eq 8, but X-ray crystallography showed that the acetate attached to uranium in a monodentate



coordination mode, $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U(\eta^1-O_2CMe)$, **7**, Figure 5. Although monodentate acetate coordination has been observed in transition metal complexes,²³ this is a new mode of acetate binding in f element chemistry where the large, highly charged, electropositive metals would be expected to interact with both acetate oxygen atoms. Given the flexibility in the coordination sphere exhibited by complexes **2** and **3**, it is surprising that this small bite angle chelate did not form a bidentate complex.

Just as **2** and **3** could be made from ${}^iPrN=C=N^iPr$ insertion reactions, **7** can also be obtained via CO_2 insertion, eq 9. Since $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-\kappa^2N,N']U$ Me is a precursor to **1**, eq 9 is a more efficient way to form **7**.



Only one other U^{4+} acetate structure was found in the literature, $[HB(3,5-Me_2Pz)_3]U(O_2CMe)_3$,²⁴ although many UO_2^{2+} acetate structures are known.²⁵ The U—(C_5Me_5 ring centroid) distances of 2.495 and 2.504 Å and

(23) See for example: (a) Cotton, F. A.; Darensbourg, D. J.; Kothammer, B. W. S.; Kudasroski, R. *Inorg. Chem.* **1982**, *21*, 1656. (b) Li, S.; Lin, Y.; Cao, J.; Zhang, S. *J. Org. Chem.* **2007**, *72*, 4067.

(24) Domingos, A.; Marcalo, J.; Marques, N.; Pires De Matos, A. *Polyhedron* **1992**, *11*, 501.

(25) See for example: (a) Zalkin, A.; Ruben, H.; Templeton, D. H. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1982**, *B38*, 610. (b) Templeton, D. H.; Zalkin, A.; Ruen, H.; Templeton, L. K. *Acta Crystallogr., Sect. C: Struct. Commun.* **1985**, *C41*, 1439. (c) Navaza, A.; Charpin, P.; Vignier, D.; Heger, G. *Acta Crystallogr., Sect. C: Struct. Commun.* **1991**, *C47*, 1842. (d) Silva, M. R.; Beja, A. M.; Paizao, J. A.; Da Veiga, L. A.; Martin-Gil, J. *Acta Crystallogr., Sect. C: Struct. Commun.* **1999**, *C55*, 2039.

(18) de Boer, E. *Adv. Organomet. Chem.* **1964**, *2*, 115.

(19) Nechaeva, O. N.; Pushkareva, Z. V. *Zh. Obshch. Khim.* **1958**, *28*, 2693.

(20) Evans, W. J.; Drummond, D. K. *J. Am. Chem. Soc.* **1989**, *111*, 3329.

(21) Hou, Z.; Fujita, A.; Zhang, Y.; Miyano, T.; Yamakazi, H.; Wakatsuki, Y. *J. Am. Chem. Soc.* **1998**, *120*, 754.



(22) Graves, C. R.; Schelter, E. J.; Cantat, T.; Scott, B. L.; Kiplinger, J. L. *Organometallics* **2008**, *27*, 5371.

Table 3. Selected Bond Lengths [Å] and Angles [deg] for $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr-{}^{\kappa^2}N,N']U(O_2CMe)_7$


bond distance/angle	7
U(1)–(C ₅ Me ₅ ring centroid)	2.495, 2.504
U(1)–N(1)	2.503(3)
U(1)–N(2)	2.370(3)
U(1)–C(21)	2.895(3)
U(1)–O(1)	2.213(2)
(C ₅ Me ₅ ring centroid)–U(1)–(C ₅ Me ₅ ring centroid)	135.4
(C ₅ Me ₅ ring centroid)–U(1)–N(1)	99.8, 99.2
(C ₅ Me ₅ ring centroid)–U(1)–N(2)	112.9, 111.0
(C ₅ Me ₅ ring centroid)–U(1)–O(1)	95.9, 96.2
N(1)–U(1)–O(1)	138.18(9)
N(2)–U(1)–O(1)	83.72(10)
N(1)–U(1)–N(2)	54.46(9)

the U–N(1) and U–N(2) distances of 2.503(3) and 2.370(3) Å in **7** (Table 3) are not unusual: they are similar to those observed in **2** (2.482(3) and 2.380(3) Å) and **6** (2.480(3) and 2.371(4) Å). The 2.213(2) Å U–O(1) bond is similar to the 2.117(9) Å of the U–O(OH) bond in $(C_5Me_5)_2UCl(OH)(HNSPh)_2$,²⁶ but much shorter than the average U–O(acetate) bond distance of 2.41(1) Å in $[HB(3,5-Me_2Pz)_3]U(O_2CMe)_3$,²⁴ in which the acetates are coordinated in a bidentate fashion. The 1.302(4) and 1.221(4) Å bond distances for C(29)–O(1) and C(29)–O(2), respectively, show the localization in the acetate ligand. Examining the crystal packing in the unit cell, a 2.55 Å distance was found between the uncoordinated oxygen of the acetate ligand and a hydrogen atom on a nearby $(C_5Me_5)^-$ ligand of an adjacent molecule (see Figure 1 in Supporting Information). This distance is much less than the sum of the covalent radii of oxygen and hydrogen.^{27,28}

The reaction of **1** with TiC_5H_5 was examined to determine if an unsubstituted cyclopentadienyl ligand could coordinate along with the $\{(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]\}^{3-}$ ligand set. Previously, it had been shown that $(C_5Me_5)^-$ was too large to add to actinide complexes with this ligation and led to an unexpected C–H bond, eq 10, and reduction reactivity.³ As shown in eq 11, an unexpected $(C_5Me_5)^-$ displacement occurred to form



$(C_5Me_5)(C_5H_5)_2[{}^iPrNC(Me)N^iPr-{}^{\kappa^2}N,N']U$, **8**. Attempts to react just 1 equiv of TiC_5H_5 with **1** only gave a 50%

**Figure 6.** Thermal ellipsoid plot of $(C_5Me_5)(C_5H_5)_2[{}^iPrNC(Me)N^iPr-{}^{\kappa^2}N,N']U$, **8**, shown at the 50% probability level. Hydrogen atoms have been omitted for clarity.**Table 4.** Selected Bond Lengths [Å] and Angles [deg] for $(C_5Me_5)(C_5H_5)_2[{}^iPrNC(Me)N^iPr-{}^{\kappa^2}N,N']U$, **8**

bond distance/angle	7
U(1)–(C ₅ Me ₅ ring centroid)	2.565
U(1)–(C ₅ H ₅ ring centroid)	2.581, 2.591
U(1)–N(1)	2.480(2)
U(1)–N(2)	2.502(2)
U(1)–C(21)	2.950(2)
(C ₅ Me ₅ ring centroid)–U(1)–(C ₅ H ₅ ring centroid)	114.1, 112.6
(C ₅ H ₅ ring centroid)–U(1)–(C ₅ H ₅ ring centroid)	105.2
(C ₅ Me ₅ ring centroid)–U(1)–N(1)	101.9
(C ₅ Me ₅ ring centroid)–U(1)–N(2)	94.2
(C ₅ H ₅ ring centroid)–U(1)–N(1)	130.9, 89.3
(C ₅ H ₅ ring centroid)–U(1)–N(2)	90.5, 138.6
N(1)–U(1)–N(2)	53.37(6)

yield of **8** with starting material remaining. Interestingly, this same $[(C_5Me_5)(C_5H_5)_2U]^{1+}$ unit is obtained cleanly from $(C_5Me_5)U(CH_2Ph)_3$ and cyclopentadiene.²⁹ Complex **8** was identified by X-ray crystallography, Figure 6. The ¹H NMR spectrum of **8** showed the presence of TiC_5Me_5 (δ 2.28 ppm in C_6D_6) as a byproduct.³⁰

The U–(C₅Me₅ ring centroid) distance of 2.565 Å in **8** (Table 4) is identical to that of the smaller of the two centroid distances in **3**, but **8** has methyl displacements (0.22–0.36 Å) in the sterically normal range. The 2.480(2) and 2.502(2) Å U–N bond lengths are also similar to those in **3** (2.408(2) and 2.510(2) Å). The average U–C–(C₅H₅) distances of 2.84(1) and 2.86(1) Å are significantly longer than those in $(C_5Me_5)(C_5H_5)_2U(CH_2C_6H_5)$ (2.74(1) and 2.75(1) Å),²⁹ $(C_5H_5)_3UCl$ (2.74(1) Å),³¹ and $(C_5Me_5)_2(C_5H_5)UMe$ (2.76(1) Å).¹⁶ The (C₅H₅ ring centroid)–U–(C₅H₅ ring centroid) angle of 105.2° is smaller than the 113.8° in $(C_5Me_5)(C_5H_5)_2U(CH_2C_6H_5)$ as well.

Discussion

$(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]U$, **1**, readily reduces disulfide substrates to make tetravalent $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]U(SR)$ complexes, **2** and **3**, eq 2 and 3. Complex **1** also reduces copper halides to produce $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]UX$

(26) Ariyaratne, K. A. N. S.; Cramer, R. E.; Jameson, G. B.; Gilje, J. W. *J. Organomet. Chem.* **2004**, *689*, 2029.

(27) Klein, R. A. *Chem. Phys. Lett.* **2006**, *425*, 128.

(28) (a) Desiraju, G. R. *The Weak Hydrogen Bond*; Oxford University Press: New York, 2001. (b) Hamilton, W. C.; Ibers, J. A. *Hydrogen Bonding in Solids*; W. A. Benjamin: New York, 1968.

(29) Kiplinger, J. L.; Morris, D. E.; Scott, B. L.; Burns, C. J. *Organometallics* **2002**, *21*, 5978.

(30) Werner, H.; Otto, H.; Kraus, H. J. *J. Organomet. Chem.* **1986**, *315*, C57.

(31) Wong, C. H.; Yen, T. M.; Lee, T. Y. *Acta Crystallogr.* **1965**, *18*, 340.

species, **5** and **6**, eq 7, as is typical for trivalent uranium metallocenes.²² Complex **1** does not react with phenazine,³² benzaldehyde azine,²⁰ benzophenone,²¹ and cyclooctatetrene,³³ substrates generally reducible with low-valent f element metallocenes. It is possible that steric crowding from the amidinate ligand may prevent coordination of these substrates that could inhibit electron transfer if an inner sphere electron mechanism is involved. However, the sterically crowded trivalent uranium complex $(C_5Me_5)_3U$ is known to reduce C_8H_8 ³⁴, and $(C_5Me_5)_3U$ is known to reduce first via U^{3+} in some cases.³⁵ Moderation of reactivity with the $\{(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]\}^{3-}$ ligand set has previously been observed with U^{4+} alkyl complexes,⁹ and was explained on the basis of steric protection. It is possible that these three ancillary ligands can protect U^{3+} as well as U^{4+} alkyls.

The formation of the monohapto acetate complex, $(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]U(O_2CMe)$ **7**, eq 8, demonstrates the type of steric variation that can be achieved with the $\{(C_5Me_5)_2[{}^iPrNC(Me)N^iPr]\}^{3-}$ ligand set. Clearly, coordination modes of common ions can be changed just by adding an amidinate to a metallocene coordination environment. Since coordination chemistry is used to control speciation in f element separation chemistry,³⁶ it is important to remember that unusual coordination modes may be accessible in the presence of the certain collections of ligands as shown in **7**.

Interestingly, although this heteroleptic ligand set leads to a monohapto acetate ligand, it can accommodate both monodentate (SPh)⁻ and bidentate (Spy)⁻ ligation in **2** and **3**, respectively. An electronic argument can be made for the difference between the thiopyridyl and acetate ligands. Since acetate is a hard-donor ligand, it may satisfy the uranium

center in a monodentate binding mode while the softer sulfur ligand requires the binding to the harder nitrogen donor atom in the pyridyl substituent.

The formation of $(C_5Me_5)(C_5H_5)_2[{}^iPr]NC(Me)N^iPr]U$, **8**, eq 11, is another unusual result from this heteroleptic ligand set, since a usually inert $(C_5Me_5)^-$ ligand is displaced from uranium. If this reaction initially involves U^{3+} reduction of Tl^{1+} , an intermediate such as $“(C_5Me_5)_2(C_5H_5)U[{}^iPrNC(Me)N^iPr]”$ could be envisioned. If the cyclopentadienyl ligands are pentahapto, this would be expected to be extremely crowded and could lose $(C_5Me_5)^-$ as previously observed in the reaction of the sterically crowded $[(C_5Me_5)_2U]_2(C_6H_6)$ with $[N(SiMe_3)_2]^-$.^{5a} The isolation of **8** suggests that several combinations of amidinate and cyclopentadienyl ligands of various substitutions should be accessible.

Conclusion

The heteroleptic ligand set $\{(C_5Me_5)_2[{}^iPr]NC(Me)N^iPr]\}^{3-}$ provides an alternative coordination environment with which to explore U^{3+} chemistry. $(C_5Me_5)_2[{}^iPr]NC(Me)N^iPr]U$, **1**, displays selective reductive chemistry that appears to be dependent on the specific substrate. Selective chemistry is also observed in terms of the types of coordination modes accessible with these ancillary ligands. This ligand set can accommodate variable coordination modes with (SPh)⁻ and (Spy)⁻ ligands, but enforces an unusual (η^1 -O₂CMe) binding mode with acetate. The $(C_5Me_5)_2[{}^iPr]NC(Me)N^iPr]U$ unit appears to be sufficiently sterically saturated that addition of ligands such as $(C_5H_5)^-$ can lead to unusual $(C_5Me_5)^-$ displacement.

Acknowledgment. We thank the Chemical Sciences, Geosciences, and Biosciences Division of the Office of Basic Energy Sciences of the Department of Energy for support of this research. We thank Michael K. Takase for assistance with X-ray crystallography and the reviewers for helpful suggestions.

Supporting Information Available: X-ray diffraction details (CIF) and X-ray data collection, structure, solution, and refinement of compounds **2–4**, **7**, and **8**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(32) Evans, W. J.; Gonzales, S. L.; Ziller, J. W. *J. Am. Chem. Soc.* **1994**, *116*, 2600.

(33) Evans, W. J.; Gonzales, S. L.; Ziller, J. W. *J. Am. Chem. Soc.* **1991**, *113*, 7423.

(34) Evans, W. J.; Nyce, G. W.; Ziller, J. W. *Angew. Chem., Int. Ed.* **2000**, *39*, 240.

(35) Evans, W. J.; Nyce, G. W.; Johnston, M. A.; Ziller, J. W. *J. Am. Chem. Soc.* **2000**, *122*, 12019.

(36) Rizkalla, E. N.; Choppin, G. R. In *Handbook on the Physics and Chemistry of Rare Earths*; Gschneidner, K. A., Eyring, L., Jr., Choppin, G. R., Lander, G. H., Eds.; Elsevier Science: New York, 1994; Vol. 18, p 529.