

that of the five-coordinate cobalt(I) *tert*-butyl isocyanide complex  $[(t\text{-BuNC})_5\text{Co}][\text{PF}_6]$  discussed above. Unchanged  $[(t\text{-BuNC})_5\text{Mn}][\text{PF}_6]$  was the only manganese(I) complex recovered from its reactions with several of the trivalent phosphorus and arsenic ligands in boiling ethanol. Even in boiling diglyme the  $[(t\text{-BuNC})_5\text{Mn}][\text{PF}_6]$  was recovered unchanged from its attempted reactions with all of the trivalent phosphorus ligands tried except for triphenylphosphine. This lower reactivity of  $[(t\text{-BuNC})_5\text{Mn}][\text{PF}_6]$  relative to  $[(t\text{-BuNC})_5\text{Co}][\text{PF}_6]$  is the expected trend for an octahedral complex relative to a trigonal-bipyramidal complex and is similar to the lower reactivity of  $\text{Cr}(\text{CO})_6$  relative to  $\text{Fe}(\text{CO})_5$ .<sup>19</sup>

The reaction between  $[(t\text{-BuNC})_5\text{Mn}][\text{PF}_6]$  and triphenylphosphine in boiling diglyme gave a low yield of the yellow crystalline monosubstituted derivative  $[(t\text{-BuNC})_5\text{MnP}(\text{C}_6\text{H}_5)_3][\text{PF}_6]$  (IX) which had to be separated from unchanged  $[(t\text{-BuNC})_5\text{Mn}][\text{PF}_6]$  by hand picking the crystals. The infrared spectrum of

$[(t\text{-BuNC})_5\text{MnP}(\text{C}_6\text{H}_5)_3][\text{PF}_6]$  (IX) exhibited three  $\nu(\text{CN})$  frequencies in accord with the  $C_{4v}$  symmetry of the complex cation. The weak  $\nu(\text{CN})$  band at  $2166\text{ cm}^{-1}$  may be assigned to the  $A_1$  mode of the  $C_{4v}$  system. The remaining  $\nu(\text{CN})$  bands at  $2093$  and  $2059\text{ cm}^{-1}$  are strong and may be assigned to the other  $A_1$  mode and the E mode. The proton nmr spectrum of  $[(t\text{-BuNC})_5\text{MnP}(\text{C}_6\text{H}_5)_3][\text{PF}_6]$  exhibited two methyl resonances with a 1:4 relative intensity ratio. The less intense resonance can be assigned to the nine methyl protons of the single axial *tert*-butyl isocyanide ligand and the more intense resonance can be assigned to the 36 methyl protons of the four equivalent equatorial *tert*-butyl isocyanide ligands.

**Acknowledgment.**—We are indebted to the Air Force Office of Scientific Research for partial support of this work under Grant AF-AFOSR-71-2000. We also acknowledge the collaboration of Miss Melanie J. Loots and Miss Susan J. Redding as students in the Secondary Science Training Program sponsored by the National Science Foundation for the summers 1970 and 1971, respectively.

(19) R. B. King, "Transition Metal Organometallic Chemistry: An Introduction," Academic Press, New York, N. Y., 1969.

CONTRIBUTION FROM THE DEPARTMENT OF CHEMISTRY,  
UNIVERSITY OF GEORGIA, ATHENS, GEORGIA 30601

## Organometallic Transition Metal Derivatives Containing Fluorine.

### V. Some Perfluoro-1-methylpropenyl Derivatives of Metal Carbonyls and Metal Cyclopentadienyls<sup>1,2</sup>

BY R. B. KING\* AND W. C. ZIPPERER

Received November 24, 1971

Reactions of various transition metal halide derivatives with perfluoro-1-methylpropenylsilver,  $\text{C}_4\text{F}_7\text{Ag}$ , in dichloromethane solution at room temperature give the corresponding perfluoro-1-methylpropenyl transition metal derivatives. Thus reactions of the metal pentacarbonyl bromides  $\text{M}(\text{CO})_5\text{Br}$  ( $\text{M} = \text{Mn}$  and  $\text{Re}$ ) with  $\text{C}_4\text{F}_7\text{Ag}$  give the volatile white crystalline  $\text{C}_4\text{F}_7\text{M}(\text{CO})_5$  ( $\text{M} = \text{Mn}$  and  $\text{Re}$ ). Similarly, treatment of  $\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2\text{I}$  with  $\text{C}_4\text{F}_7\text{Ag}$  gives the volatile yellow crystalline  $\text{C}_4\text{F}_7\text{Fe}(\text{CO})_2\text{C}_5\text{H}_5$ . Reaction of  $\text{C}_5\text{H}_5\text{Cr}(\text{NO})_2\text{Cl}$  with  $\text{C}_4\text{F}_7\text{Ag}$  gives the volatile green crystalline  $\text{C}_4\text{F}_7\text{Cr}(\text{NO})_2\text{C}_5\text{H}_5$ . Reactions of the fluorocarbon transition metal halides  $\text{R}_t\text{Fe}(\text{CO})_4\text{I}$  ( $\text{R}_t = \text{C}_2\text{F}_5$ ,  $\text{CF}_3\text{CF}_2\text{CF}_2$ , and  $(\text{CF}_3)_2\text{CF}$ ) and  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{R}_t)\text{I}$  ( $\text{R}_t = \text{C}_2\text{F}_5$  and  $\text{CF}_3\text{CF}_2\text{CF}_2$ ) with  $\text{C}_4\text{F}_7\text{Ag}$  give volatile pale yellow  $\text{R}_t\text{Fe}(\text{CO})_4\text{C}_4\text{F}_7$  ( $\text{R}_t = \text{C}_2\text{F}_5$ ,  $\text{CF}_3\text{CF}_2\text{CF}_2$ , and  $(\text{CF}_3)_2\text{CF}$ ) and volatile yellow  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{R}_t)(\text{C}_4\text{F}_7)$  ( $\text{R}_t = \text{C}_2\text{F}_5$  and  $\text{CF}_3\text{CF}_2\text{CF}_2$ ), respectively. The infrared, fluorine nmr, and mass spectra of these new compounds are discussed.

#### Introduction

Within the last decade fluorocarbon transition metal derivatives have received considerable attention largely because of their stability relative to their hydrocarbon analogs.<sup>3</sup> Such compounds are prepared by the following general methods: (1) reactions of metal carbonyl anions with perfluoroacyl derivatives followed by decarbonylation; (2) nucleophilic substitution of fluoride with metal carbonyl groups by reactions of fluorinated olefins or aromatic compounds with metal carbonyl anions;<sup>4</sup> (3) addition of fluoroolefins, fluorin-

ated alkynes, or perfluoroalkyl iodides to appropriate transition metal systems, particularly those in lower oxidation states; (4) reactions of metal halide derivatives with fluoroalkyl or fluoroaryl derivatives of electropositive metals such as lithium or magnesium. This repertoire of preparative methods in fluorocarbon transition metal chemistry has the following limitations: (1) the need of a sufficiently nucleophilic metal carbonyl anion for the reactions with perfluoroacyl derivatives, fluoroolefins, and fluorinated aromatic compounds (methods 1 and 2 above) or the need of an appropriately reactive low-oxidation-state derivative for the reactions with fluoroolefins, fluorinated alkynes, or perfluoroalkyl iodides (method 3 above); (2) the instability of many fluorocarbon derivatives of electropositive metals such as lithium or magnesium coupled with the high reactivity of organometallic derivatives of such electropositive metals which makes them able to effect undesirable side reactions (method 4). For these

(1) For part IV of this series see R. B. King, R. N. Kapoor, and K. H. Pannell, *J. Organometal. Chem.*, **20**, 187 (1969).

(2) Portions of this work were presented to the Division of Fluorine Chemistry at the 162nd National Meeting of the American Chemical Society, Washington, D. C., Sept 1971; see Abstract FLUO-7.

(3) For a review article discussing fluorocarbon transition metal derivatives, see P. M. Treichel and F. G. A. Stone, *Advan. Organometal. Chem.*, **1**, 143 (1964).

(4) For a review of nucleophilic reactions of metal carbonyl anions with fluorocarbons, see M. I. Bruce and F. G. A. Stone, *Angew. Chem., Int. Ed. Engl.*, **7**, 747 (1968).


TABLE I  
 REACTIONS OF PERFLUORO-1-METHYLPROPENYLSILVER WITH VARIOUS TRANSITION METAL HALIDE DERIVATIVES

Halide (g, mmol)	Quantity of C <sub>4</sub> F <sub>7</sub> Ag, g (mmol)	Solvent (ml)	Time, hr	Product	Yield, g (%)
Mn(CO) <sub>5</sub> Br (1.0, 3.9)	1.42 (4.9)	CH <sub>2</sub> Cl <sub>2</sub> (50)	4	C <sub>4</sub> F <sub>7</sub> Mn(CO) <sub>5</sub>	0.67 (46)
Re(CO) <sub>5</sub> Br (1.1, 2.8)	1.0 (3.5)	CH <sub>2</sub> Cl <sub>2</sub> (50)	5	C <sub>4</sub> F <sub>7</sub> Re(CO) <sub>5</sub>	1.02 (73)
C <sub>5</sub> H <sub>5</sub> Fe(CO) <sub>2</sub> I (0.84, 2.8)	1.0 (3.5)	CH <sub>2</sub> Cl <sub>2</sub> (50)	5	C <sub>4</sub> F <sub>7</sub> Fe(CO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	0.7 (71)
C <sub>5</sub> H <sub>5</sub> Fe(CO) <sub>2</sub> I (0.84, 2.8)	1.0 (3.5)	THF (50)	1	C <sub>4</sub> F <sub>7</sub> Fe(CO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	0.11 (11)
C <sub>5</sub> H <sub>5</sub> Cr(NO) <sub>2</sub> Cl (0.58, 2.8)	1.0 (3.5)	CH <sub>2</sub> Cl <sub>2</sub> (50)	3	C <sub>4</sub> F <sub>7</sub> Cr(NO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	0.68 (68)
C <sub>2</sub> F <sub>5</sub> Fe(CO) <sub>4</sub> I (4.60, 9.9)	4.0 (13.8)	CH <sub>2</sub> Cl <sub>2</sub> (100)	3	(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>	1.23 (27)
( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub> I (1.81, 4.9)	1.42 (4.9)	CH <sub>2</sub> Cl <sub>2</sub> (50)	10	( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>	0.89 (44)
( <i>i</i> -C <sub>3</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub> I (1.29, 2.8)	1.0 (3.5)	CH <sub>2</sub> Cl <sub>2</sub> (50)	6	( <i>i</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>	0.44 (29)
C <sub>5</sub> H <sub>5</sub> Co(CO)(C <sub>2</sub> F <sub>5</sub> )I (0.9, 2.3)	0.82 (2.8)	CH <sub>2</sub> Cl <sub>2</sub> (40)	6	C <sub>5</sub> H <sub>5</sub> Co(CO)(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )	0.25 (20)
C <sub>5</sub> H <sub>5</sub> Co(CO)( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )I (0.4, 0.9)	0.34 (1.2)	CH <sub>2</sub> Cl <sub>2</sub> (50)	6	C <sub>5</sub> H <sub>5</sub> Co(CO)( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )	0.07 (16)
(C <sub>5</sub> H <sub>5</sub> ) <sub>2</sub> TiCl <sub>2</sub> (0.69, 2.8)	2.0 (6.9)	CH <sub>2</sub> Cl <sub>2</sub> (50)	23	(C <sub>5</sub> H <sub>5</sub> ) <sub>2</sub> TiF <sub>2</sub>	0.06 (8)
(C <sub>5</sub> H <sub>5</sub> ) <sub>2</sub> ZrCl <sub>2</sub> (0.8, 2.8)	2.0 (6.9)	CH <sub>2</sub> Cl <sub>2</sub> (50)	14	(C <sub>5</sub> H <sub>5</sub> ) <sub>2</sub> ZrF <sub>2</sub>	0.09 (12)

reasons the development of new and distinctly different methods for the synthesis of fluorocarbon transition metal derivatives would allow a considerable extension of the range of fluorocarbon transition metal derivatives that can be prepared.

A significant recent development in fluorocarbon chemistry has been the discovery of stable fluorocarbon derivatives of the coinage metals. Stable compounds of this type include pentafluorophenylcopper (C<sub>6</sub>F<sub>5</sub>Cu),<sup>5</sup> perfluoro-*tert*-butylcopper [(CF<sub>3</sub>)<sub>3</sub>CCu],<sup>5</sup> perfluoro-1-methylpropenylsilver (C<sub>4</sub>F<sub>7</sub>Ag),<sup>6</sup> and pentafluorophenylsilver (C<sub>6</sub>F<sub>5</sub>Ag).<sup>7</sup> Reactions of such fluorocarbon derivatives of coinage metals with transition metal halide derivatives appeared to provide a method for the preparation of fluorocarbon transition metal derivatives not obtainable by previous methods. Fluorocarbon derivatives of coinage metals would have a high tendency to react with transition metal halide derivatives because of the insolubility of the coinage metal halide but would have a much lower tendency to effect undesirable side reactions than similar fluorocarbon derivatives of relatively electropositive metals such as lithium or magnesium.

These considerations led us to investigate reactions of coinage metal fluorocarbon derivatives with transition metal halide derivatives as possible routes to new fluorocarbon transition metal derivatives. We selected perfluoro-1-methylpropenylsilver (I) as the coinage metal fluorocarbon derivative for our studies for the following reasons: (1) its relatively facile preparation<sup>6</sup> from silver(I) fluoride and the commercially available hexafluorobutylene-2; (2) its stability which allows purification by vacuum sublimation. This paper reports the details of our work in this area. Of particular interest is the preparation of the first known derivatives with two different fluorocarbon groups bonded to the same transition metal atom.



### Experimental Section

A nitrogen atmosphere was always provided for the following three operations: (a) carrying out reactions, (b) handling all filtered solutions of organometallic compounds, and (c) admitting to evacuated vessels containing organometallic compounds.

**Materials.**—Perfluoro-1-methylpropenylsilver (I) was prepared

(5) A. Cairncross and W. A. Sheppard, *J. Amer. Chem. Soc.*, **90**, 2186 (1968).

(6) W. T. Miller, R. H. Snider, and R. J. Hummel, *ibid.*, **91**, 6532 (1969).

(7) K. K. Sun and W. T. Miller, *ibid.*, **92**, 6985 (1970).

by the published procedure<sup>6</sup> which employs the reaction of silver(I) fluoride with hexafluorobutylene-2 in acetonitrile at room temperature. We purified the acetonitrile used in this preparation by distillation over phosphorus pentoxide. The perfluoro-1-methylpropenylsilver (I) was always vacuum sublimed (~100° (0.1 mm)) before use.

The following transition metal derivatives were purchased from the indicated commercial sources: Fe(CO)<sub>5</sub> (GAF Corp., New York, N. Y.), Co<sub>2</sub>(CO)<sub>8</sub> (Strem Chemical Co., Danvers, Mass.), CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>Mn(CO)<sub>3</sub> (Ethyl Corp., New York, N. Y.), Mo(CO)<sub>6</sub> and Re<sub>2</sub>(CO)<sub>10</sub> (Pressure Chemical Corp., Pittsburgh, Pa.), and (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>MCl<sub>2</sub> (M = Ti and Zr) (Arapahoe Chemical Corp., Boulder, Colo.). The compounds Mn<sub>2</sub>(CO)<sub>10</sub>,<sup>8</sup> Mn(CO)<sub>5</sub>Br,<sup>9a</sup> C<sub>5</sub>H<sub>5</sub>Fe(CO)<sub>2</sub>I,<sup>9b</sup> C<sub>5</sub>H<sub>5</sub>Cr(NO)<sub>2</sub>Cl,<sup>9c</sup> R<sub>f</sub>Fe(CO)<sub>4</sub>I (R<sub>f</sub> = C<sub>2</sub>F<sub>5</sub>,<sup>10</sup> CF<sub>3</sub>CF<sub>2</sub>CF<sub>2</sub>,<sup>9d</sup> and (CF<sub>3</sub>)<sub>2</sub>CF<sub>2</sub>),<sup>11</sup> C<sub>5</sub>H<sub>5</sub>Co(CO)<sub>2</sub>Fe,<sup>9e</sup> C<sub>5</sub>H<sub>5</sub>Co(CO)(R<sub>f</sub>)I (R<sub>f</sub> = C<sub>2</sub>F<sub>5</sub> and CF<sub>3</sub>CF<sub>2</sub>CF<sub>2</sub>),<sup>12</sup> [C<sub>5</sub>H<sub>5</sub>Mo(NO)I]<sub>2</sub>,<sup>13</sup> and Fe(CO)<sub>4</sub>I<sub>2</sub><sup>14</sup> were prepared by the cited published procedures.

**Reactions of Transition Metal Halide Derivatives with Perfluoro-1-methylpropenylsilver (Table I).**—The indicated quantities of the transition metal halide derivative and perfluoro-1-methylpropenylsilver were stirred in the indicated quantity of dichloromethane (redistilled over phosphorus pentoxide) for the indicated period of time (see Table I). After the reaction period was over, the precipitated silver(I) halide was removed by filtration. Solvent was removed from the dichloromethane filtrate at 25° (25 mm). The product was isolated from the residue in the indicated yield (Table I) by vacuum sublimation (or distillation in the case of (C<sub>2</sub>F<sub>5</sub>)(C<sub>4</sub>F<sub>7</sub>)Fe(CO)<sub>4</sub>).

Analytical data and other properties of the new perfluoro-1-methylpropenyl transition metal derivatives are given in Table II.

The product (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>TiF<sub>2</sub> from the reaction of (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>TiCl<sub>2</sub> and C<sub>4</sub>F<sub>7</sub>Ag was identified by comparison of its infrared and mass spectra with those reported for an authentic sample.<sup>15</sup> The white crystalline (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrF<sub>2</sub> was identified by its mass spectrum (see below) and by elemental analyses. *Anal.* Calcd for C<sub>10</sub>H<sub>10</sub>F<sub>2</sub>Zr: C, 46.4; H, 3.9; F, 14.7; Zr, 35.1. Found: C, 46.4; H, 3.7; F, 14.6; Zr, 34.9. The fluorine nmr spectrum of (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrF<sub>2</sub> exhibited a single resonance at  $\phi$  -177.4.

**Infrared Spectra (Table III).**—The infrared spectra of the solid compounds listed in Table III were taken in potassium bromide pellets and recorded on a Perkin-Elmer Model 621 spectrometer. The infrared spectrum of the liquid (C<sub>2</sub>F<sub>5</sub>)(C<sub>4</sub>F<sub>7</sub>)Fe(CO)<sub>4</sub> was measured as a liquid film. The  $\nu$ (CO) regions of the metal carbonyl derivatives and the  $\nu$ (NO) region of C<sub>4</sub>F<sub>7</sub>Cr(NO)<sub>2</sub>C<sub>5</sub>H<sub>5</sub> (Table III) were also measured in cyclo-

(8) R. B. King, J. C. Stokes, and T. F. Korenowski, *J. Organometal. Chem.*, **11**, 641 (1968).

(9) R. B. King, "Organometallic Syntheses," Vol. 1, Academic Press, New York, N. Y., 1965: (a) p 174; (b) p 175; (c) p 161; (d) p 177; (e) p 118, but using no ultraviolet irradiation and using boiling dichloromethane as the reaction solvent.

(10) R. B. King, S. L. Stafford, P. M. Treichel, and F. G. A. Stone, *J. Amer. Chem. Soc.*, **83**, 3604 (1961).

(11) R. B. King, R. N. Kapoor, and L. W. Houk, *J. Inorg. Nucl. Chem.*, **31**, 2179 (1969).

(12) R. B. King, P. M. Treichel, and F. G. A. Stone, *J. Amer. Chem. Soc.*, **83**, 3593 (1961).

(13) R. B. King, *Inorg. Chem.*, **6**, 30 (1967).

(14) W. Hieber and G. Bader, *Ber.*, **61**, 1717 (1928).

(15) R. B. King and N. Welcman, *Inorg. Chem.*, **8**, 2540 (1969).

TABLE II  
 PROPERTIES OF THE TRANSITION METAL PERFLUORO-1-METHYLPROPENYL DERIVATIVES

Compound <sup>a</sup>	Color	Mp, <sup>b</sup> °C	Sublimation conditions, °C (mm)	Analyses, % <sup>c</sup>				
				C	H	F	Metal	Other
C <sub>4</sub> F <sub>7</sub> Mn(CO) <sub>5</sub>	White	38	25 (0.1)	28.8 (28.8)		35.3 (35.4)	14.7 (14.6) (Mn)	
C <sub>4</sub> F <sub>7</sub> Re(CO) <sub>5</sub>	White	63	25 (0.05)	21.3 (21.4)	0.2 (0.0)	26.0 (26.2)	36.6 (36.7) (Re)	16.0 (15.8) (O)
C <sub>4</sub> F <sub>7</sub> Fe(CO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	Yellow	75	40 (0.05)	36.6 (36.9)	1.4 (1.4)	37.0 (37.2)	15.4 (15.6) (Fe)	
C <sub>4</sub> F <sub>7</sub> Cr(NO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	Green	80	50 (10)	30.1 (30.2)	1.5 (1.4)	37.2 (37.2)	14.6 (14.6) (Cr)	
(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>	Pale yellow	Liquid	40 (0.1)	24.9 (25.1)	0.2 (0.0)	48.6 (48.8)	11.9 (12.0) (Fe)	
( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>	Pale yellow	51-53	44 (0.1)	23.8 (25.4)	0.4 (0.0)	50.2 (51.4)	11.2 (10.8) (Fe)	12.7 (12.4) (O)
( <i>i</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>	Pale yellow	55	25 (0.1)	25.4 (25.4)	0.1 (0.0)	51.3 (51.4)	10.9 (10.8) (Fe)	12.4 (12.4) (O)
C <sub>5</sub> H <sub>5</sub> Co(CO)(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )	Yellow	75	35 (0.05)	31.7 (31.9)	1.1 (1.1)	50.6 (50.7)	13.1 (13.1) (Co)	
C <sub>5</sub> H <sub>5</sub> Co(CO)( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )	Yellow	94	25 (0.05)	31.2 (31.1)	1.1 (1.0)	52.7 (53.0)	11.7 (11.8) (Co)	

<sup>a</sup> The following abbreviations were used: C<sub>4</sub>F<sub>7</sub> = perfluoro-1-methylpropenyl; *n*-C<sub>3</sub>F<sub>7</sub> = CF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>; *i*-C<sub>3</sub>F<sub>7</sub> = (CF<sub>3</sub>)<sub>2</sub>CF. <sup>b</sup> Melting points were taken in capillaries and are uncorrected. <sup>c</sup> Analyses were performed by Meade Microanalytical Laboratory, Amherst, Mass. Calculated values in parentheses.

 TABLE III  
 INFRARED SPECTRA OF THE TRANSITION METAL PERFLUORO-1-METHYLPROPENYL DERIVATIVES

Compound	Infrared spectrum, cm <sup>-1</sup>				
	ν(CH) <sup>a</sup>	ν(CO) <sup>b</sup>	ν(C=C) <sup>a</sup>	ν(CF) <sup>a</sup>	Other bands <sup>a</sup>
C <sub>4</sub> F <sub>7</sub> Mn(CO) <sub>5</sub>		2142 (w, A <sub>1</sub> ), 2056 (vs, E), 2025 (s, A <sub>1</sub> )	1618 (m)	1311 (s), 1224 (vs), 1190 (vs), 1153 (vs), 1126 (vs), 1052 (m)	889 (s), 846 (s), 740 (s), 698 (s), 659 (vs), 641 (vs), 619 (s)
C <sub>4</sub> F <sub>7</sub> Re(CO) <sub>5</sub>		2155 (w, A <sub>1</sub> ), 2048 (vs, E), 2015 (s, A <sub>1</sub> )	1620 (m)	1320 (s), 1227 (vs), 1205 (vs), 1137 (vvs), 1055 (m)	961 (m), 900 (s), 847 (s), 741 (s), 698 (w), 667 (s), 610 (s), 590 (s)
C <sub>4</sub> F <sub>7</sub> Fe(CO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	3150 (vw)	2054 (s), 2012 (s)	1630 (w)	1320 (m), 1228 (s), 1194 (s), 1148 (m), 1134 (s), 1113 (m, sh), 1054 (w)	1432 (w), 1401 (w), 1020 (w), 1010 (vw), 897 (m), 854 (w), 844 (w), 734 (w), 654 (w)
C <sub>4</sub> F <sub>7</sub> Cr(NO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	3135 (w)	1820 (s), <sup>c</sup> 1720 (s) <sup>c</sup>	1636 (w)	1323 (s), 1235 (s), 1192 (s), 1138 (s), 1112 (m), 1054 (w)	1439 (w), 1434 (w), 1400 (w), 1024 (w), 1015 (vw) 936 (vw), 905 (m), 848 (m), 842 (m), 748 (w), 664 (m), 645 (w), 613 (w)
(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub> <sup>d</sup>		2167 (vw), 2131 (w), 2103 (s), 2071 (w)	1621 (m)	1324 (s), 1310 (s), 1227 (vs), 1200 (vs), 1163 (vs), 1052 (s), 919 (s)	902 (s), 849 (s), 742 (s), 733 (s), 670 (s), 638 (s)
( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>		2166 (vw), 2129 (vw, sh), 2102 (s), 2072 (w)	1620 (w)	1320 (s), 1226 (s), 1204 (s), 1162 (s), 1098 (s), 1066 (w), 1028 (s)	897 (m), 858 (m), 813 (s), 738 (m), 728 (m), 667 (m), 636 (s)
( <i>i</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>		2135 (vw), 2107 (s), 2077 (vw), 2067 (vw)	1615 (w)	1328 (s), 1292 (m), 1266 (s), 1220-1194 (vs, br), 1167 (vs), 1130 (m), 1079 (w), 1023 (s), 943 (s)	900 (m), 878 (s), 849 (m), 742 (s), 710 (s), 668 (s)
C <sub>5</sub> H <sub>5</sub> Co(CO)(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )	3131 (w)	2108 (m), 2100 (m)	1614 (w)	1314 (s), 1285 (s), 1229 (s), 1203 (s, sh), 1191 (s), 1153 (s, sh), 1140 (s), 1059 (m), 1045 (m), 1037 (m), 892 (s)	1437 (w), 1422 (w), 962 (w), 908 (m), 862 (s), 848 (m), 837 (w), 741 (m), 732 (s), 663 (s)
C <sub>5</sub> H <sub>5</sub> Co(CO)( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )	3146 (w)	2105 (s)	1624 (w)	1321 (s), 1252 (m, sh), 1231 (s), 1201 (s), 1181 (m), 1158 (s), 1129 (m), 1093 (m), 1077 (w), 1060 (w), 1044 (w), 1030 (m), 1011 (w)	1448 (w), 1431 (w), 964 (vw), 907 (m), 867 (s), 850 (m), 841 (w), 807 (m), 743 (w), 725 (s), 668 (m), 663 (m)

<sup>a</sup> These frequencies were determined in potassium bromide pellets. <sup>b</sup> These frequencies were determined in cyclohexane solution. Assignments of the ν(CO) frequencies for the C<sub>4</sub>F<sub>7</sub>M(CO)<sub>5</sub> (M = Mn or Re) derivatives are given in parentheses. <sup>c</sup> ν(NO) frequency. <sup>d</sup> A liquid film rather than a KBr pellet was used for the infrared spectrum of this liquid compound.

 TABLE IV  
 FLUORINE NMR SPECTRA OF THE TRANSITION METAL PERFLUORO-1-METHYLPROPENYL DERIVATIVES<sup>a</sup>

Compound	Perfluoro-1-methylpropenyl fluorines					Group	Other fluorines <sup>b</sup>		
	β-CF <sub>3</sub>	γ-CF <sub>3</sub>	β-CF	<sup>3</sup> J(FF)	<sup>4</sup> J(FF) <sup>5</sup> J(FF)		α(CF or CF <sub>2</sub> )	β(CF <sub>2</sub> or CF <sub>3</sub> )	γ(CF <sub>3</sub> )
C <sub>4</sub> F <sub>7</sub> Ag	50.7	70.0	95.1	12.5	15.1 1.7				
C <sub>4</sub> F <sub>7</sub> Mn(CO) <sub>5</sub> <sup>c</sup>	48.8	65.0	71.9	7	20.7				
C <sub>4</sub> F <sub>7</sub> Re(CO) <sub>5</sub>	48.4	63.3	80.5	~6	19.0				
C <sub>4</sub> F <sub>7</sub> Fe(CO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	47.8	63.3	71.8	~6	27.3				
C <sub>4</sub> F <sub>7</sub> Cr(NO) <sub>2</sub> C <sub>5</sub> H <sub>5</sub>	50.8	65.3	79.7	~6	23.0				
(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub> <sup>d</sup>	47.1	64.2	68.9			C <sub>2</sub> F <sub>5</sub>	~64?	83.6	
( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub> <sup>e</sup>	49.0	65.4	69.2	7	22.3 1.7	<i>n</i> -C <sub>3</sub> F <sub>7</sub>	61.3 q (10.5)	115.5	80.2 t (10.5)
( <i>i</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> )Fe(CO) <sub>4</sub>	46.6	63.7	66?		25.9	<i>i</i> -C <sub>3</sub> F <sub>7</sub>	68.6 d (~4)		
C <sub>5</sub> H <sub>5</sub> Co(CO)(C <sub>2</sub> F <sub>5</sub> )(C <sub>4</sub> F <sub>7</sub> )	50.1	62.6	68.3 d (103) <sup>e</sup>			C <sub>2</sub> F <sub>5</sub>	?	80.9	
C <sub>5</sub> H <sub>5</sub> Co(CO)( <i>n</i> -C <sub>3</sub> F <sub>7</sub> )(C <sub>4</sub> F <sub>7</sub> ) <sup>e</sup>	50.0	63.8	69.8 d (186) <sup>e</sup>		25.0	<i>n</i> -C <sub>3</sub> F <sub>7</sub>	53.4 br	115	80.9 t (8.5)

<sup>a</sup> These spectra were obtained at 56.456 MHz on a Perkin-Elmer Model R-20 spectrometer unless otherwise indicated. <sup>b</sup> The following abbreviations are used: d = doublet, t = triplet, q = quartet, br = broad. Separations in Hz are given in parentheses. <sup>c</sup> This spectrum was obtained at 94.1 MHz on a Varian HA-100 spectrometer. <sup>d</sup> The nmr spectrum of this compound exhibited broad resonances apparently because of the presence of paramagnetic impurities. <sup>e</sup> This "doublet" appears to arise from a chemical shift difference rather than spin-spin coupling since in the fluorine nmr spectrum of C<sub>5</sub>H<sub>5</sub>Co(CO)(*n*-C<sub>3</sub>F<sub>7</sub>)(C<sub>4</sub>F<sub>7</sub>) the separation was ~120 Hz at 56.456 MHz and ~185 Hz at 94.1 MHz. This must arise from the presence of some type of isomerism.

hexane solution. All spectra were calibrated against the 1601.4-cm<sup>-1</sup> band of polystyrene film.

**Fluorine Nmr Spectra (Table IV).**—The fluorine nmr spectra listed in Table IV were taken in dichloromethane solution with either fluorotrichloromethane ( $\phi$  0.0) or 1,2-difluoro-1,1,2,2-tetrachloroethane ( $\phi$  67.8) as an internal standard and recorded either at 56.456 MHz on a Perkin-Elmer Hitachi Model R-20 spectrometer or at 94.1 MHz on a Varian HA-100 spectrometer. All fluorine nmr chemical shift data are given in the  $\phi$  scale of Filipovich and Tiers.<sup>16</sup>

**Mass Spectra.**—The mass spectra listed below were taken at the indicated temperatures at 70 eV on a Perkin-Elmer Hitachi RMU-6 mass spectrometer. Relative intensities are given in parentheses.

**A. C<sub>4</sub>F<sub>7</sub>Re(CO)<sub>5</sub> (Chamber Temperature 100°; Sample Temperature ~70°).**—C<sub>4</sub>F<sub>7</sub>Re(CO)<sub>5</sub><sup>+</sup> (37), C<sub>4</sub>F<sub>7</sub>Re(CO)<sub>4</sub><sup>+</sup> (14), C<sub>4</sub>F<sub>7</sub>Re(CO)<sub>3</sub><sup>+</sup> (11), C<sub>4</sub>F<sub>7</sub>Re(CO)<sub>2</sub><sup>+</sup> (13), C<sub>4</sub>F<sub>7</sub>ReCO<sup>+</sup> (35), C<sub>4</sub>F<sub>6</sub>ReCO<sup>+</sup> (4), C<sub>4</sub>F<sub>6</sub>Re<sup>+</sup> (69), C<sub>4</sub>F<sub>6</sub>Re<sup>+</sup> (6), Re(CO)<sub>5</sub><sup>+</sup> (44), Re(CO)<sub>4</sub>F<sup>+</sup> (58), Re(CO)<sub>4</sub><sup>+</sup> (60), Re(CO)<sub>3</sub>F<sup>+</sup> (100), C<sub>3</sub>F<sub>3</sub>Re<sup>+</sup> (13), CF<sub>4</sub>Re<sup>+</sup> (2), Re(CO)<sub>3</sub><sup>+</sup> (12), C<sub>2</sub>F<sub>3</sub>Re<sup>+</sup> (10), Re(CO)<sub>2</sub>F<sup>+</sup> (17), C<sub>3</sub>F<sub>2</sub>Re<sup>+</sup> (6), CF<sub>3</sub>Re<sup>+</sup> (4), C<sub>2</sub>F<sub>2</sub>Re<sup>+</sup> (11), ReF<sub>3</sub><sup>+</sup> (14), Re(CO)<sub>2</sub><sup>+</sup> (4), CF<sub>2</sub>Re<sup>+</sup> (3), ReCOF<sup>+</sup> (8), ReC<sub>2</sub>F<sup>+</sup> (7), ReF<sub>2</sub><sup>+</sup> (7), CFRe<sup>+</sup> (3), C<sub>4</sub>F<sub>6</sub>Re(CO)<sub>2</sub><sup>+</sup> (1), ReCO<sup>+</sup> (2), ReC<sub>2</sub><sup>+</sup> (3), ReF<sup>+</sup> (5), ReC<sup>+</sup> (3), C<sub>4</sub>F<sub>6</sub>ReCO<sup>+</sup> (1), Re<sup>+</sup> (6), C<sub>4</sub>F<sub>7</sub>H<sup>+</sup> (89), C<sub>4</sub>F<sub>6</sub>Re<sup>+</sup> (3), C<sub>4</sub>F<sub>6</sub>H<sup>+</sup> (190), Re(CO)<sub>4</sub><sup>+</sup> (2), C<sub>4</sub>F<sub>5</sub><sup>+</sup> (190), C<sub>3</sub>F<sub>3</sub>H<sup>+</sup> (19), C<sub>4</sub>F<sub>4</sub><sup>+</sup> (42), C<sub>3</sub>F<sub>4</sub>H<sup>+</sup> (440), C<sub>4</sub>F<sub>3</sub><sup>+</sup> (8), C<sub>2</sub>F<sub>4</sub><sup>+</sup> (3), C<sub>3</sub>F<sub>3</sub><sup>+</sup> (360), C<sub>2</sub>F<sub>3</sub>H<sup>+</sup> (20), C<sub>3</sub>F<sub>2</sub>H<sup>+</sup> (43), C<sub>3</sub>F<sub>2</sub><sup>+</sup> (48), and CF<sub>3</sub><sup>+</sup> (270).

**B. C<sub>4</sub>F<sub>7</sub>Fe(CO)<sub>5</sub>C<sub>5</sub>H<sub>5</sub> (Chamber Temperature 100°; Sample Temperature 100°).**—C<sub>4</sub>F<sub>7</sub>Fe(CO)<sub>5</sub>C<sub>5</sub>H<sub>5</sub><sup>+</sup> (53), C<sub>4</sub>F<sub>7</sub>FeCOC<sub>5</sub>H<sub>5</sub><sup>+</sup> (88), C<sub>4</sub>F<sub>7</sub>FeC<sub>5</sub>H<sub>5</sub><sup>+</sup> (100), C<sub>4</sub>F<sub>6</sub>FeC<sub>5</sub>H<sub>4</sub><sup>+</sup> (4), *m/e* 207 (7), C<sub>4</sub>F<sub>7</sub>H<sup>+</sup> (7), C<sub>5</sub>H<sub>5</sub>Fe(CO)<sub>2</sub><sup>+</sup> (79), *m/e* 169 (10), C<sub>5</sub>H<sub>5</sub>FeCOF<sup>+</sup> (7), C<sub>4</sub>F<sub>6</sub>H<sup>+</sup> (13), C<sub>4</sub>F<sub>6</sub><sup>+</sup> (12), C<sub>5</sub>H<sub>5</sub>FeCO<sup>+</sup> (24), C<sub>4</sub>F<sub>5</sub><sup>+</sup> (125), C<sub>5</sub>H<sub>5</sub>FeF<sup>+</sup> (>1000), C<sub>4</sub>F<sub>4</sub><sup>+</sup> (26), C<sub>5</sub>H<sub>4</sub>Fe<sup>+</sup> (44), C<sub>3</sub>F<sub>4</sub>H<sub>2</sub><sup>+</sup> (22), C<sub>3</sub>F<sub>4</sub>H<sup>+</sup> (28), C<sub>3</sub>F<sub>3</sub><sup>+</sup> (145), FeF<sup>+</sup> (180), C<sub>5</sub>H<sub>5</sub><sup>+</sup> (125), Fe<sup>+</sup> (62), and C<sub>3</sub>H<sub>3</sub><sup>+</sup> (130). Metastable ions at *m/e* 304 s [C<sub>4</sub>F<sub>7</sub>Fe(CO)<sub>5</sub>C<sub>5</sub>H<sub>5</sub><sup>+</sup> → C<sub>4</sub>F<sub>7</sub>FeCOC<sub>5</sub>H<sub>5</sub><sup>+</sup> + CO], *m/e* 278 s [C<sub>4</sub>F<sub>7</sub>FeCOC<sub>5</sub>H<sub>5</sub><sup>+</sup> → C<sub>4</sub>F<sub>7</sub>FeC<sub>5</sub>H<sub>5</sub><sup>+</sup> + CO], *m/e* 102.9 m [C<sub>5</sub>H<sub>5</sub>FeF<sup>+</sup> → C<sub>5</sub>H<sub>4</sub>Fe<sup>+</sup> + HF], *m/e* 65.0 [C<sub>4</sub>F<sub>7</sub>FeC<sub>5</sub>H<sub>5</sub><sup>+</sup> → C<sub>5</sub>H<sub>5</sub>FeF<sup>+</sup> + C<sub>4</sub>F<sub>6</sub>], *m/e* 60.5 w [C<sub>4</sub>F<sub>5</sub><sup>+</sup> → C<sub>3</sub>F<sub>3</sub><sup>+</sup> + CF<sub>2</sub>], and *m/e* 23.5 w [C<sub>5</sub>H<sub>5</sub><sup>+</sup> → C<sub>3</sub>H<sub>3</sub><sup>+</sup> + C<sub>2</sub>H<sub>2</sub>].

**C. C<sub>4</sub>F<sub>7</sub>Cr(NO)<sub>2</sub>C<sub>5</sub>H<sub>5</sub> (Chamber Temperature 220°; Sample Temperature 80°).**—C<sub>4</sub>F<sub>7</sub>Cr(NO)<sub>2</sub>C<sub>5</sub>H<sub>5</sub><sup>+</sup> (8), C<sub>4</sub>F<sub>7</sub>CrNOCC<sub>5</sub>H<sub>5</sub><sup>+</sup> (45), C<sub>4</sub>F<sub>7</sub>CrC<sub>5</sub>H<sub>5</sub><sup>+</sup> (4), C<sub>4</sub>F<sub>7</sub>H<sup>+</sup> (33), C<sub>5</sub>H<sub>5</sub>Cr(NO)<sub>2</sub><sup>+</sup> (27), C<sub>5</sub>H<sub>5</sub>CrNOF<sup>+</sup> (14), C<sub>4</sub>F<sub>6</sub>H<sup>+</sup> (72), C<sub>4</sub>F<sub>6</sub><sup>+</sup> (~400), C<sub>5</sub>H<sub>5</sub>CrF<sup>+</sup> (4), C<sub>3</sub>F<sub>3</sub><sup>+</sup> (3), C<sub>5</sub>H<sub>5</sub>CrNO<sup>+</sup> (5), C<sub>4</sub>F<sub>5</sub><sup>+</sup> (170), C<sub>5</sub>H<sub>5</sub>CrF<sup>+</sup> (100), C<sub>4</sub>F<sub>4</sub><sup>+</sup> (29), C<sub>3</sub>F<sub>4</sub>H<sup>+</sup> (48), C<sub>3</sub>F<sub>3</sub><sup>+</sup> (124), C<sub>2</sub>F<sub>3</sub>H<sup>+</sup> (12), C<sub>2</sub>F<sub>3</sub>H<sup>+</sup> (33), C<sub>2</sub>F<sub>2</sub><sup>+</sup> (46), CrF<sup>+</sup> (53), CF<sub>3</sub><sup>+</sup> (~50), C<sub>5</sub>H<sub>5</sub><sup>+</sup> (66), C<sub>5</sub>H<sub>5</sub> (25), Cr<sup>+</sup> (23), and C<sub>3</sub>H<sub>3</sub><sup>+</sup> (29). Metastable ions at *m/e* 300.5 w [C<sub>4</sub>F<sub>7</sub>Cr(NO)<sub>2</sub>C<sub>5</sub>H<sub>5</sub><sup>+</sup> → C<sub>4</sub>F<sub>7</sub>CrNOCC<sub>5</sub>H<sub>5</sub><sup>+</sup> + NO], *m/e* 60.5 w [C<sub>4</sub>F<sub>5</sub><sup>+</sup> → C<sub>3</sub>F<sub>3</sub><sup>+</sup> + CF<sub>2</sub>], and *m/e* 53.5 w [C<sub>4</sub>F<sub>6</sub><sup>+</sup> → C<sub>3</sub>F<sub>3</sub><sup>+</sup> + CF<sub>3</sub>].

**D. C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>2</sub>F<sub>5</sub>)(C<sub>4</sub>F<sub>7</sub>) (Chamber Temperature 80°).**—C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>2</sub>F<sub>5</sub>)(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (3.2), C<sub>5</sub>H<sub>5</sub>Co(C<sub>2</sub>F<sub>5</sub>)(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (3.5), C<sub>5</sub>H<sub>5</sub>CoC<sub>6</sub>F<sub>11</sub><sup>+</sup> (1.9), C<sub>5</sub>H<sub>5</sub>CoC<sub>6</sub>F<sub>10</sub><sup>+</sup> (9), C<sub>5</sub>H<sub>5</sub>CoC<sub>6</sub>F<sub>9</sub><sup>+</sup> (1.6), C<sub>5</sub>H<sub>5</sub>CoC<sub>6</sub>F<sub>8</sub><sup>+</sup> (1.9), C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (51), C<sub>5</sub>H<sub>5</sub>CoF(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (1.4), C<sub>5</sub>H<sub>4</sub>CoF(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (0.5), C<sub>5</sub>H<sub>5</sub>CoC<sub>4</sub>F<sub>7</sub><sup>+</sup> (100), C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>2</sub>F<sub>5</sub>)<sup>+</sup> (20), C<sub>5</sub>H<sub>5</sub>CoF(C<sub>2</sub>F<sub>5</sub>)<sup>+</sup> (1.9), C<sub>5</sub>H<sub>5</sub>CoC<sub>3</sub>F<sub>5</sub><sup>+</sup> (9), C<sub>5</sub>H<sub>5</sub>Co(CO)CF<sub>3</sub><sup>+</sup> (1.1), C<sub>4</sub>F<sub>6</sub>H<sup>+</sup> (2.2), C<sub>4</sub>F<sub>5</sub><sup>+</sup> (2.2), *m/e* 207 (15), C<sub>4</sub>F<sub>3</sub><sup>+</sup> (2.2), C<sub>5</sub>H<sub>5</sub>CoCF<sub>3</sub><sup>+</sup> (3.2), C<sub>4</sub>F<sub>7</sub>H<sup>+</sup> (4), C<sub>5</sub>H<sub>5</sub>CoCF<sub>2</sub><sup>+</sup> (3.2), C<sub>3</sub>F<sub>7</sub><sup>+</sup> (3.2), C<sub>4</sub>F<sub>6</sub>H<sup>+</sup> (7), C<sub>4</sub>F<sub>5</sub><sup>+</sup> (3.5), C<sub>5</sub>H<sub>5</sub>CoCF<sup>+</sup> (4), C<sub>5</sub>H<sub>5</sub>CoCO<sup>+</sup> (14), C<sub>5</sub>H<sub>5</sub>CoF<sup>+</sup> and/or C<sub>4</sub>F<sub>5</sub><sup>+</sup> (320), *m/e* 127 (9), C<sub>5</sub>H<sub>5</sub>Co<sup>+</sup> and/or C<sub>4</sub>F<sub>4</sub><sup>+</sup> (110), C<sub>5</sub>H<sub>4</sub>Co<sup>+</sup> (95), C<sub>3</sub>F<sub>4</sub>H<sup>+</sup> (16), C<sub>2</sub>F<sub>4</sub><sup>+</sup> (7), C<sub>2</sub>H<sub>3</sub>Co<sup>+</sup> (14), C<sub>3</sub>H<sub>2</sub>Co<sup>+</sup> (17), C<sub>3</sub>HCO<sup>+</sup> (7), C<sub>3</sub>F<sub>3</sub><sup>+</sup> (46), CoF<sup>+</sup> (15), C<sub>3</sub>F<sub>2</sub>H<sup>+</sup> (5), C<sub>3</sub>F<sub>2</sub><sup>+</sup> (6), CF<sub>3</sub><sup>+</sup> (23), C<sub>5</sub>H<sub>5</sub><sup>+</sup> (72), C<sub>5</sub>H<sub>3</sub><sup>+</sup> (14), Co<sup>-</sup> (55), C<sub>4</sub>H<sub>3</sub><sup>+</sup> (5), C<sub>4</sub>H<sub>2</sub><sup>+</sup> (2.7), and C<sub>3</sub>H<sub>3</sub><sup>+</sup> (~54). Metastable ions at *m/e* 347 w [C<sub>5</sub>H<sub>5</sub>CoC<sub>6</sub>F<sub>10</sub><sup>+</sup> → C<sub>5</sub>H<sub>5</sub>CoC<sub>6</sub>F<sub>9</sub><sup>+</sup> + HF], *m/e* 279 s [C<sub>5</sub>H<sub>5</sub>Co(CO)C<sub>4</sub>F<sub>7</sub><sup>+</sup> → C<sub>5</sub>H<sub>5</sub>CoC<sub>4</sub>F<sub>7</sub><sup>+</sup> + CO], *m/e* 217.5 w [C<sub>5</sub>H<sub>5</sub>Co(CO)C<sub>6</sub>F<sub>8</sub><sup>+</sup> → C<sub>5</sub>H<sub>5</sub>CoC<sub>6</sub>F<sub>7</sub><sup>+</sup> + CO], *m/e* 105.9 vs [C<sub>5</sub>H<sub>5</sub>CoF<sup>+</sup> → C<sub>5</sub>H<sub>4</sub>Co<sup>+</sup> + HF], *m/e* 67.0 vw [C<sub>5</sub>H<sub>5</sub>CoC<sub>4</sub>F<sub>7</sub><sup>+</sup> → C<sub>5</sub>H<sub>5</sub>CoF<sup>+</sup> + C<sub>4</sub>F<sub>6</sub>], and *m/e* 23.5 w [C<sub>5</sub>H<sub>5</sub><sup>+</sup> → C<sub>3</sub>H<sub>3</sub><sup>+</sup> + C<sub>2</sub>H<sub>2</sub>].

**E. C<sub>5</sub>H<sub>5</sub>Co(CO)(CF<sub>2</sub>CF<sub>2</sub>CF<sub>3</sub>)(C<sub>4</sub>F<sub>7</sub>) (Chamber Temperature 80°).**—C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>3</sub>F<sub>7</sub>)(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (1.8), C<sub>5</sub>H<sub>5</sub>Co(C<sub>3</sub>F<sub>7</sub>)(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (4.4), C<sub>5</sub>H<sub>5</sub>CoC<sub>7</sub>F<sub>13</sub><sup>+</sup> (2.1), C<sub>5</sub>H<sub>5</sub>CoC<sub>7</sub>F<sub>12</sub><sup>+</sup> (7), C<sub>5</sub>H<sub>5</sub>CoC<sub>7</sub>F<sub>11</sub><sup>+</sup> (0.6), C<sub>5</sub>H<sub>4</sub>CoC<sub>7</sub>F<sub>11</sub><sup>+</sup> (0.3), C<sub>5</sub>H<sub>5</sub>Co(CO)C<sub>5</sub>F<sub>9</sub><sup>+</sup> (1.2), C<sub>5</sub>H<sub>5</sub>CoC<sub>5</sub>F<sub>8</sub><sup>+</sup> (1.8), C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> (42), C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>3</sub>F<sub>7</sub>)<sup>+</sup> (1.5), C<sub>5</sub>H<sub>5</sub>CoC<sub>4</sub>F<sub>7</sub><sup>+</sup> (100), C<sub>5</sub>H<sub>5</sub>CoC<sub>3</sub>F<sub>7</sub><sup>+</sup> (12), C<sub>5</sub>H<sub>5</sub>CoC<sub>2</sub>F<sub>5</sub><sup>+</sup> (2.9), C<sub>4</sub>F<sub>6</sub>H<sup>+</sup> (1.2), C<sub>4</sub>F<sub>5</sub><sup>+</sup> (1.2), *m/e* 207 (14), *m/e* 195 (5.3), C<sub>4</sub>F<sub>7</sub><sup>+</sup> (4.4), C<sub>5</sub>H<sub>5</sub>CoCF<sub>2</sub><sup>+</sup> (4.4), C<sub>5</sub>H<sub>5</sub>Co(CO)F<sup>+</sup> (1.2), C<sub>3</sub>F<sub>7</sub><sup>+</sup> (3.5), C<sub>4</sub>F<sub>6</sub>H<sup>+</sup> (7), C<sub>4</sub>F<sub>6</sub><sup>+</sup> (4.7), C<sub>5</sub>H<sub>5</sub>CoCF<sup>+</sup> (3.8), C<sub>5</sub>H<sub>5</sub>CoCO<sup>+</sup>

(17), C<sub>5</sub>H<sub>5</sub>CoF<sup>+</sup> and/or C<sub>4</sub>F<sub>5</sub><sup>+</sup> (~1050), C<sub>3</sub>F<sub>5</sub><sup>+</sup> (9), *m/e* 127 (15), C<sub>5</sub>H<sub>5</sub>Co<sup>+</sup> and/or C<sub>4</sub>F<sub>4</sub><sup>+</sup> (105), C<sub>4</sub>H<sub>4</sub>Co<sup>+</sup> (91), C<sub>3</sub>F<sub>4</sub>H<sup>+</sup> (17), C<sub>2</sub>F<sub>4</sub><sup>+</sup> (6), C<sub>3</sub>H<sub>3</sub>Co<sup>+</sup> (14), C<sub>3</sub>H<sub>2</sub>Co<sup>+</sup> (18), C<sub>3</sub>HCO<sup>+</sup> (7), C<sub>2</sub>F<sub>3</sub><sup>+</sup> (39), CoF<sup>+</sup> (12), C<sub>3</sub>F<sub>2</sub>H<sup>+</sup> (4.7), C<sub>3</sub>F<sub>2</sub><sup>+</sup> (5.3), CF<sub>3</sub><sup>+</sup> (38), C<sub>5</sub>H<sub>5</sub><sup>+</sup> (~38), C<sub>5</sub>H<sub>3</sub><sup>+</sup> (13), Co<sup>+</sup> (~38), C<sub>4</sub>H<sub>3</sub><sup>+</sup> (6), C<sub>4</sub>H<sub>2</sub><sup>+</sup> (2.6), and C<sub>3</sub>H<sub>3</sub><sup>+</sup> (~38). Metastable ions at *m/e* 279 s [C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>4</sub>F<sub>7</sub>)<sup>+</sup> → C<sub>5</sub>H<sub>5</sub>CoC<sub>4</sub>F<sub>7</sub><sup>+</sup> + CO], *m/e* 267.5 m [C<sub>5</sub>H<sub>5</sub>Co(CO)(C<sub>3</sub>F<sub>7</sub>)<sup>+</sup> → C<sub>5</sub>H<sub>5</sub>CoC<sub>3</sub>F<sub>7</sub><sup>+</sup> + CO], *m/e* 105.9 s [C<sub>5</sub>H<sub>5</sub>CoF<sup>+</sup> → C<sub>5</sub>H<sub>4</sub>Co<sup>+</sup> + HF], 67.0 w [C<sub>5</sub>H<sub>5</sub>CoC<sub>4</sub>F<sub>7</sub><sup>+</sup> → C<sub>5</sub>H<sub>5</sub>CoF<sup>+</sup> + C<sub>4</sub>F<sub>6</sub>], and *m/e* 23.5 w [C<sub>5</sub>H<sub>5</sub><sup>+</sup> → C<sub>3</sub>H<sub>3</sub><sup>+</sup> + C<sub>2</sub>H<sub>2</sub>].

**F. (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrF<sub>2</sub> (Chamber Temperature 220°; Sample Temperature ~140°).**—(C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrF<sub>2</sub><sup>+</sup> (24), (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrF<sup>+</sup> (5), C<sub>5</sub>H<sub>5</sub>ZrF<sub>2</sub><sup>+</sup> (100), C<sub>5</sub>H<sub>5</sub>ZrF<sub>2</sub><sup>+</sup> (16), C<sub>5</sub>H<sub>5</sub>ZrF<sup>+</sup> (2), C<sub>10</sub>H<sub>10</sub><sup>+</sup> (38), C<sub>10</sub>H<sub>9</sub><sup>+</sup> (42), C<sub>10</sub>H<sub>8</sub><sup>+</sup> (28), C<sub>9</sub>H<sub>7</sub><sup>+</sup> (27), C<sub>8</sub>H<sub>6</sub><sup>+</sup> (145), C<sub>8</sub>H<sub>5</sub><sup>+</sup> (98), C<sub>5</sub>H<sub>3</sub><sup>+</sup> (24), and C<sub>3</sub>H<sub>3</sub><sup>+</sup> (54).

## Discussion

The syntheses of perfluoro-1-methylpropenyl transition metal derivatives from corresponding metal halide derivatives and perfluoro-1-methylpropenylsilver by metathetical reactions resemble previously reported metathetical syntheses of perfluorocarboxylate<sup>17</sup> and trifluoromethylthio<sup>15</sup> transition metal derivatives using the silver perfluorocarboxylates and trifluoromethylthiosilver, respectively. However, the metathetical reactions of perfluoro-1-methylpropenylsilver with transition metal halide derivatives reported in this paper are the first reported examples of the formation of a metal-carbon bond from a metathetical reaction involving a silver derivative. The only previous examples of the use of an organosilver compound to form a metal-carbon bond are the very recently reported<sup>18</sup> preparations of the perfluoro-1-methylpropenylmetal pentacarbonyl anions [C<sub>4</sub>F<sub>7</sub>M(CO)<sub>5</sub>]<sup>-</sup> (M = Cr, Mo, and W) by oxidation of the corresponding decacarbonyl-dimetalates [M<sub>2</sub>(CO)<sub>10</sub>]<sup>2-</sup> (M = Cr, Mo, and W) with perfluoro-1-methylpropenylsilver.

The reactions of perfluoro-1-methylpropenylsilver with transition metal halide derivatives to form the corresponding perfluoro-1-methylpropenyl transition metal derivatives appear to be limited to transition metal halide derivatives with only one replaceable halogen atom. Thus attempts to prepare derivatives with two perfluoro-1-methylpropenyl groups attached to a single transition metal atom by reactions of perfluoro-1-methylpropenylsilver with the transition metal dihalide derivatives Fe(CO)<sub>4</sub>I<sub>2</sub>, C<sub>5</sub>H<sub>5</sub>Co(CO)I<sub>2</sub>, and [C<sub>5</sub>H<sub>5</sub>Mo(NO)I<sub>2</sub>]<sub>2</sub> gave no identifiable fluorocarbon transition metal derivatives. Reactions of the bis(cyclopentadienyl)metal dichlorides of titanium and zirconium, (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>MCl<sub>2</sub> (M = Ti and Zr), with perfluoro-1-methylpropenylsilver gave only the corresponding bis(cyclopentadienyl)metal difluorides (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>MF<sub>2</sub> (M = Ti or Zr). Apparently, the high tendency for fluorine to bond to the relatively electropositive 4+ titanium and zirconium results in a fluoride shift from carbon to the titanium or zirconium possibly with elimination of hexafluorobutene-2 in a reaction of a reverse type to the formation of perfluoro-1-methylpropenylsilver from silver fluoride and hexafluorobutene-2. A similar fluoride shift has previously<sup>15</sup> been observed in the reaction of (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>TiCl<sub>2</sub> with trifluoromethylthiosilver to give (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>TiF<sub>2</sub>.

The perfluoro-1-methylpropenyl group in all of the new transition metal derivatives prepared in this work

(17) R. B. King and R. N. Kapoor, *J. Organometal. Chem.*, **15**, 457 (1968).

(18) W. J. Schlientz and J. K. Ruff, *ibid.*, **33**, C64 (1971).

(16) G. Filipovich and G. V. D. Tiers, *J. Phys. Chem.*, **63**, 761 (1959).

exhibits a characteristic infrared  $\nu(\text{C}=\text{C})$  frequency in the range  $1615\text{--}1630\text{ cm}^{-1}$  and characteristic  $\nu(\text{CF})$  frequencies of medium to very strong intensities in the five ranges  $1310\text{--}1328$ ,  $1224\text{--}1235$ ,  $1190\text{--}1205$ ,  $1137\text{--}1163$ , and  $1052\text{--}1055\text{ cm}^{-1}$  (see Table III). The range of  $\nu(\text{C}=\text{C})$  frequencies for the transition metal perfluoro-1-methylpropenyl derivatives falls considerably below the  $1675\text{-cm}^{-1}$   $\nu(\text{C}=\text{C})$  frequency in perfluoro-1-methylpropenylsilver and the  $1739\text{-cm}^{-1}$   $\nu(\text{C}=\text{C})$  frequency in the fluoroolefin *trans*- $\text{CF}_3\text{CF}=\text{CHCF}_3$ .<sup>6</sup> The lower  $\nu(\text{C}=\text{C})$  frequencies in the transition metal perfluoro-1-methylpropenyl derivatives can be taken as an indication of the reduction of the carbon-carbon bond order in the perfluoro-1-methylpropenyl group by back-donation of electrons from the filled transition metal d orbitals into the antibonding orbitals of the carbon-carbon double bond of the perfluoro-1-methylpropenyl group. A similar lowering of the  $\nu(\text{C}=\text{C})$  frequency of a polyfluoroalkenyl group upon bonding to a transition metal has been observed in other organometallic derivatives.<sup>19</sup>


The fluorine nmr spectra of the transition metal perfluoro-1-methylpropenyl derivatives (Table IV) exhibit the expected two trifluoromethyl doublets in the ranges  $\delta$  46.6–50.8 and 62.3–65.4 and a broad resonance (not always readily and unequivocally observed) in the range  $\delta$  66–80.5 arising from the single olefinic fluorine atom. By analogy with the nmr spectrum of perfluoro-1-methylpropenylsilver<sup>6</sup> the lower field trifluoromethyl doublet is assigned to the trifluoromethyl group bonded to the carbon atom which is also bonded to the transition metal (designated as  $\beta\text{-CF}_3$  in Table IV because the fluorines are bonded to a  $\beta$  carbon atom relative to the transition metal) and the higher field trifluoromethyl doublet is assigned to the trifluoromethyl group bonded to the carbon atom which is also bonded to the single olefinic fluorine atom (designated as  $\gamma\text{-CF}_3$  in Table IV because the fluorines are bonded to a  $\gamma$  carbon atom relative to the transition metal).

The couplings of the single olefinic fluorine atom to the two different trifluoromethyl groups ( $^3J(\text{FF})$  and  $^4J(\text{FF})$ ) were observed in the fluorine nmr spectra of most of the perfluoro-1-methylpropenyl transition metal derivatives (Table IV). However, the coupling between the fluorine atoms in the two different trifluoromethyl groups ( $^5J(\text{FF})$ ) was only 1.7 Hz and was thus observed only in a few of the better spectra. Comparison with the published<sup>19–21</sup>  $^5J(\text{FF})$  coupling constants between the fluorine atoms in the two trifluoromethyl groups in the various 1-trifluoromethyl-3,3,3-trifluoropropenyl derivatives obtained by the addition of metal hydrides to hexafluorobutyn-2 indicates that the two trifluoromethyl groups in the perfluoro-1-methylpropenyl transition metal derivatives prepared in this work are in *trans* positions around the carbon-carbon double bond. Thus in the previous work<sup>19–21</sup> with 1-trifluoromethyl-3,3,3-trifluoropropenyl derivatives, the compounds with a *trans* configuration of the trifluoromethyl groups around the carbon-carbon double bond exhibited a  $^5J(\text{FF})$  coupling constant between the fluorine atoms of the two trifluoromethyl groups of

1 to 3 Hz, whereas the compounds with a *cis* configuration of the trifluoromethyl groups around the carbon-carbon double bond exhibited a much larger  $^5J(\text{FF})$  coupling constant between the fluorine atoms of the two trifluoromethyl groups of 12 to 15 Hz. These nmr observations on the small magnitude of the  $^5J(\text{FF})$  coupling constant in the perfluoro-1-methylpropenyl transition metal derivatives thus indicate that the *trans* configuration of the trifluoromethyl groups around the carbon-carbon bond in perfluoro-1-methylpropenylsilver(I) is retained upon reactions with transition metal halides. A similar retention of the *trans* configuration of the trifluoromethyl groups around the carbon-carbon double bond has been observed in previously reported<sup>6</sup> reactions of perfluoro-1-methylpropenylsilver with other halogen derivatives such as methyl iodide.

A further feature of interest in the fluorine nmr spectra (Table IV) of the perfluoro-1-methylpropenyl transition metal derivatives is the observation that the  $^3J(\text{FF})$  coupling constant ( $\sim 7$  Hz) is much less than the more remote (in terms of number of bonds)  $^4J(\text{FF})$  coupling constant (18 to 28 Hz). A similar effect was observed several years ago<sup>22</sup> in the fluorine nmr spectra of transition metal *n*-heptafluoropropyl derivatives where the  $^3J(\text{FF})$  coupling constants ( $< 2$  Hz) were found to be much less than the more remote  $^4J(\text{FF})$  coupling constants (11 to 13 Hz).

The work described in this paper resulted in the preparation and characterization of nine new perfluoro-1-methylpropenyl transition metal derivatives. Of these nine derivatives the three compounds  $\text{C}_4\text{F}_7\text{M}(\text{CO})_5$  (II, M = Mn and Re) and  $\text{C}_4\text{F}_7\text{Fe}(\text{CO})_2\text{C}_5\text{H}_5$  (III, M = Fe; L = CO) resemble numerous previously reported compounds of the types  $\text{R}_t\text{M}(\text{CO})_5$  (M = Mn and Re)<sup>23</sup> and  $\text{R}_t\text{Fe}(\text{CO})_2\text{C}_5\text{H}_5$ ,<sup>4,24</sup> respectively, in such respects as color, volatility, and infrared spectra of the metal carbonyl group in the  $\nu(\text{CO})$  region. The compound  $\text{C}_4\text{F}_7\text{Cr}(\text{NO})_2\text{C}_5\text{H}_5$  (III, M = Cr; L = NO) is apparently the first known neutral fluorocarbon chro-



mium derivative and also apparently the first fluorocarbon derivative of a metal nitrosyl system. However, analogous  $\text{RCr}(\text{NO})_2\text{C}_5\text{H}_5$  derivatives, but only with hydrocarbon groups bonded to the chromium atom (*e.g.*, R = methyl, ethyl, chloromethyl, and  $\sigma$ -cyclopentadienyl), have been known for some time<sup>25</sup> being preparable by reactions of the halides  $\text{C}_5\text{H}_5\text{Cr}(\text{NO})_2\text{X}$  (X = Cl and I) with organomagnesium halides or sodium cyclopentadienide. The compound  $\text{C}_4\text{F}_7\text{Cr}(\text{NO})_2\text{C}_5\text{H}_5$  (III, M = Cr; L = NO), like the previously reported  $\text{RCr}(\text{NO})_2\text{C}_5\text{H}_5$  derivatives,<sup>25</sup> is a very

(19) H. C. Clark and W. S. Tsang, *J. Amer. Chem. Soc.*, **89**, 533 (1967).

(20) W. R. Cullen, D. S. Dawson, and G. E. Styan, *Can. J. Chem.*, **43**, 3892 (1965).

(21) H. C. Clark and R. J. Puddephatt, *Inorg. Chem.*, **9**, 2670 (1970).

(22) E. Pitcher, A. D. Buckingham, and F. G. A. Stone, *J. Chem. Phys.*, **36**, 124 (1962).


(23) J. B. Wilford and F. G. A. Stone, *Inorg. Chem.*, **4**, 389 (1965), and references cited therein.

(24) R. B. King and M. B. Bisnette, *J. Organometal. Chem.*, **2**, 15 (1964).

(25) T. S. Piper and G. Wilkinson, *J. Inorg. Nucl. Chem.*, **3**, 104 (1956).

volatile green solid which exhibits two strong  $\nu(\text{NO})$  frequencies in its infrared spectrum.

The remaining five compounds prepared in this work are of the types  $\text{R}_f\text{Fe}(\text{CO})_4\text{C}_4\text{F}_7$  (IV,  $\text{R}_f = \text{C}_2\text{H}_5$ ,  $\text{CF}_3\text{CF}_2\text{CF}_2$ , and  $(\text{CF}_3)_2\text{CF}$ ) and  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{R}_f)-(\text{C}_4\text{F}_7)$  (V,  $\text{R}_f = \text{C}_2\text{H}_5$  or  $\text{CF}_3\text{CF}_2\text{CF}_2$ ) and represent the first examples of fluorocarbon transition metal derivatives with two different fluorocarbon groups bonded to a single transition metal atom. The fluorine nmr spectrum (Table IV) of each of these five compounds exhibited resonances with chemical shifts and fine structures corresponding to those expected both for the perfluoro-1-methylpropenyl and the other perfluoroalkyl groups. This provides evidence supporting the proposed structures IV and V and excludes otherwise unlikely possibilities of coupling and/or rearrangement of the two fluorocarbon groups bonded to the transition metal to form a more complex system.



The iron compounds of the type  $\text{C}_4\text{F}_7\text{Fe}(\text{CO})_4\text{R}_f$  (IV,  $\text{R}_f = \text{C}_2\text{F}_5$ ,  $\text{CF}_3\text{CF}_2\text{CF}_2$ , and  $(\text{CF}_3)_2\text{CF}$ ) exhibit a single strong  $\nu(\text{CO})$  frequency around  $2100\text{ cm}^{-1}$  in their infrared spectra in addition to several much weaker  $\nu(\text{CO})$  frequencies (see Table III). This pattern is tentatively interpreted as suggestive of the indicated isomers IV where the two different fluorocarbon groups occupy trans positions of the iron octahedron. If the two fluorocarbon groups in the iron derivatives IV were equivalent and axially symmetric, these compounds would have  $D_{4h}$  symmetry which would give rise to only one infrared-active  $\nu(\text{CO})$  frequency, a strong  $E_u$  mode. Nonequivalence of the fluorocarbon groups in IV reduces the symmetry to  $C_{4v}$  which would add a weaker infrared-active  $A_1$  mode to the already very strongly infrared-active E mode. Lack of axial symmetry of the fluorocarbon groups in IV (a possible consequence of restricted rotation around the metal-carbon ( $\text{C}_4\text{F}_7$ ) bond in the perfluoro-1-methylpropenyl derivatives arising from retrodonative bonding) would further reduce the symmetry of the system which would lead to slight infrared activity of the normally only Raman-active  $B_1$  mode<sup>23</sup> and to splitting of the doubly degenerate E mode. The weak  $\nu(\text{CO})$  frequencies accompanying the single strong  $\nu(\text{CO})$  frequency in the  $\text{R}_f\text{Fe}(\text{CO})_4\text{C}_4\text{F}_7$  derivatives most likely arise from these effects associated with deviation of the system from strict  $D_{4h}$  symmetry. The alternative explanation of the extra weak  $\nu(\text{CO})$  frequencies as indicative of the cis isomers of the  $\text{R}_f\text{Fe}(\text{CO})_4\text{C}_4\text{F}_7$  derivatives is less probable because observations on numerous authentic *cis*- $\text{L}_2\text{M}(\text{CO})_4$  isomers (e.g., (diphos) $\text{M}(\text{CO})_4$ )<sup>26</sup> indicate similar relative intensities of at least three of the four  $\nu(\text{CO})$  modes in contrast to the actual observations on the  $\text{R}_f\text{Fe}(\text{CO})_4\text{C}_4\text{F}_7$  derivatives.

(26) For an example of data of this type, see the first three entries of Table I in R. B. King, P. N. Kapoor, and R. N. Kapoor, *Inorg. Chem.*, **10**, 1841 (1971).

The cobalt compounds  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{R}_f)(\text{C}_4\text{F}_7)$  (V,  $\text{R}_f = \text{C}_2\text{F}_5$  or  $\text{CF}_3\text{CF}_2\text{CF}_2$ ) are of interest since few compounds of the type  $\text{C}_5\text{H}_5\text{Co}(\text{CO})\text{R}_2$  are known, even when both alkyl or perfluoroalkyl groups are the same. Previous examples of compounds of the type  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{R}_f)_2$  in the literature include only the perfluoro-tetramethylene derivative  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{CF}_2)_4$  (VI)<sup>27</sup> obtained from  $\text{C}_5\text{H}_5\text{Co}(\text{CO})_2$  and tetrafluoroethylene; compounds such as  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{C}_3\text{F}_7)_2$  were briefly mentioned at a meeting several years ago<sup>28</sup> but have not been studied in detail because their preparation was inefficient and inconvenient. Like  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{CF}_2)_4$  (VI) the compounds  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{R}_f)(\text{C}_4\text{F}_7)$  (V,  $\text{R}_f = \text{C}_2\text{F}_5$  or  $\text{CF}_3\text{CF}_2\text{CF}_2$ ) are volatile yellow solids which exhibit the expected single  $\nu(\text{CO})$  frequency around  $2100\text{ cm}^{-1}$ . However, in the *n*-heptafluoropropyl derivative  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{C}_2\text{F}_5)(\text{C}_4\text{F}_7)$  (V;  $\text{R}_f = \text{C}_2\text{F}_5$ ) this  $\nu(\text{CO})$  frequency is split by  $8\text{ cm}^{-1}$  possibly because of the presence of conformational isomers arising from restricted rotation around one of the metal-fluorocarbon bonds similar to the splitting of the  $\nu(\text{CO})$  frequencies in cyclopentadienyliron dicarbonyl compounds such as  $\text{CH}_3\text{SiCl}_2\text{Fe}(\text{CO})_2\text{C}_5\text{H}_5$  arising from restricted rotation around the iron-silicon bond.<sup>29</sup> This type of isomerism could also be responsible for the observation of two  $\beta$ -CF resonances separated by  $\sim 2\text{ ppm}$  in the fluorine nmr spectra of both  $\text{C}_5\text{H}_5\text{Co}(\text{CO})(\text{R}_f)-(\text{C}_4\text{F}_7)$  derivatives (Table IV and ref 3).

The mass spectra of the perfluoro-1-methylpropenyl transition metal derivatives were investigated. The compounds  $\text{C}_4\text{F}_7\text{Mn}(\text{CO})_5$  and  $\text{R}_f\text{Fe}(\text{CO})_4\text{C}_4\text{F}_7$  failed to exhibit any metal-containing ions because of decomposition even when the chamber temperature was held to about  $100^\circ$ . Instead, only fluorocarbon ions were observed in these mass spectra.

The mass spectrum of the rhenium compound  $\text{C}_4\text{F}_7\text{Re}(\text{CO})_5$  (II,  $\text{M} = \text{Re}$ ) exhibited the series of ions  $\text{C}_4\text{F}_7\text{Re}(\text{CO})_n^+$  ( $n = 5, 4, 3, 2, 1$ , and  $0$ ),  $\text{C}_4\text{F}_6\text{Re}(\text{CO})_n^+$  ( $n = 1$  and  $0$ ),  $\text{Re}(\text{CO})_n\text{F}^+$  ( $n = 4, 3, 2, 1$ , and  $0$ ), and  $\text{Re}(\text{CO})_n^+$  ( $n = 5, 4, 3, 2, 1$ , and  $0$ ) and the dipositive ion series  $\text{C}_4\text{F}_6\text{Re}(\text{CO})_n^{2+}$  ( $n = 3, 1$ , and  $0$ ). The dipositive ions  $\text{C}_4\text{F}_6\text{Re}(\text{CO})_n^{2+}$  can arise by loss of fluoride ( $\text{F}^-$ ) from the relatively abundant monopositive ions  $\text{C}_4\text{F}_7\text{Re}(\text{CO})_n^+$ . In addition, fragments of the types  $\text{C}_3\text{F}_n\text{Re}^+$  ( $n = 3$  and  $2$ ),  $\text{C}_2\text{F}_n\text{Re}^+$  ( $n = 3, 2, 1$ , and  $0$ ), and  $\text{CF}_n\text{Re}^+$  ( $n = 4, 3, 2, 1$ , and  $0$ ) are observed where the rhenium-carbon bond to the perfluoro-1-methylpropenyl group has been maintained but some carbon-carbon bonds of the perfluoro-1-methylpropenyl group have broken.

The mass spectrum of  $\text{C}_4\text{F}_7\text{Fe}(\text{CO})_2\text{C}_5\text{H}_5$  (III,  $\text{M} = \text{Fe}$ ;  $\text{L} = \text{CO}$ ) exhibited features typical of cyclopentadienyliron dicarbonyl derivatives<sup>30</sup> particularly those with fluorocarbon groups bonded to the iron atom.<sup>31</sup> The observed metastable ions indicate that the molecular ion  $\text{C}_4\text{F}_7\text{Fe}(\text{CO})_2\text{C}_5\text{H}_5^+$  can fragment to the  $\text{C}_5\text{H}_4\text{Fe}^+$  ion by the successive losses of its two carbonyl groups followed by elimination of a neutral

(27) T. D. Coyle, R. B. King, E. Pitcher, S. L. Stafford, P. M. Treichel, and F. G. A. Stone, *J. Inorg. Nucl. Chem.*, **20**, 172 (1961).

(28) P. M. Treichel and G. Werber, Abstracts, 150th National Meeting of the American Chemical Society, Atlantic City, N. J., Sept 1965, No. INOR 51.

(29) W. Jetz and W. A. G. Graham, *J. Amer. Chem. Soc.*, **89**, 2773 (1967).

(30) R. B. King, *ibid.*, **90**, 1417 (1968).

(31) R. B. King, *Appl. Spectrosc.*, **23**, 137 (1969).

$C_4F_6$  fragment (presumably hexafluorobutene-2) from  $C_4F_7FeC_5H_5^+$  to give  $C_5H_5FeF^+$  which can then undergo loss of a neutral hydrogen fluoride fragment to give  $C_5H_4Fe^+$ . The mass spectrum of the chromium derivative  $C_4F_7Cr(NO)_2C_5H_5$  (III,  $M = Cr$ ;  $L = NO$ ), as expected, exhibited many features similar to the mass spectrum of the isostructural and isoelectronic  $C_4F_7Fe(CO)_2C_5H_5$  (III,  $M = Fe$ ;  $L = CO$ ).

The mass spectra of the cobalt derivatives  $C_5H_5Co(CO)(R_f)(C_4F_7)$  (V) provide qualitative comparisons of the relative elimination tendencies of perfluoro-1-methylpropenyl and saturated perfluoroalkyl groups. In both cases (V,  $R_f = C_2F_5$  and  $CF_3CF_2CF_2$ ) the ion  $C_5H_5Co(CO)(C_4F_7)^+$  is over twice as abundant as the ion  $C_5H_5Co(CO)R_f^+$  ( $R_f = C_2F_5$  or  $CF_3CF_2CF_2$ ) suggesting that elimination of a saturated perfluoroalkyl group from the molecular ion  $C_5H_5Co(CO)(R_f)(C_4F_7)^+$  occurs significantly more readily than elimination of the unsaturated perfluoro-1-methylpropenyl group. This suggests that the bond of the cobalt atom to a saturated perfluoroalkyl group is weaker than the bond of the cobalt atom to the unsaturated perfluoro-1-methylpropenyl group. This effect can be rationalized by the availability of empty antibonding orbitals in the carbon-carbon double bond of the perfluoro-1-methyl-

propenyl group which can overlap with filled d orbitals of the cobalt atom to provide additional means to strengthen the metal-perfluoro-1-methylpropenyl bond by retrodonative bonding. However, arguments of this type based on mass spectra are necessarily imprecise since they rely on the assumption of similar further fragmentation tendencies of the ion  $C_5H_5Co(CO)(C_4F_7)^+$  and the saturated perfluoroalkyl ion  $C_5H_5Co(CO)R_f^+$ .

The mass spectra of the cobalt derivatives  $C_5H_5Co(CO)(R_f)(C_4F_7)$  (V,  $R_f = C_2F_5$  and  $CF_3CF_2CF_2$ ) also exhibited some features similar to those found in the mass spectra of the iron derivative  $C_4F_7Fe(CO)_2C_5H_5$  (III,  $M = Fe$ ;  $L = CO$ ) discussed above. Thus metastable ions were observed in the mass spectra of both cobalt compounds which correspond to the elimination of a neutral  $C_4F_6$  fragment from  $C_5H_5CoC_4F_7^+$  to give  $C_5H_5CoF^+$  and the elimination of a neutral HF fragment from  $C_5H_5CoF^+$  to give  $C_5H_4Co^+$ .

**Acknowledgment.**—We are indebted to the National Institute of General Medical Sciences for partial support of this work under Grant GM-14664-03. We also acknowledge the assistance of Dr. Alan Bond with some of the fluorine nmr experiments.

CONTRIBUTION FROM THE CHEMISTRY DEPARTMENT, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BRITISH COLUMBIA, CANADA

## Reactions of Fluorocarbon-Bridged Di(tertiary phosphines and arsines) with Manganese and Rhenium Carbonyls

By J. P. CROW, W. R. CULLEN,\* AND F. L. HOU

Received December 27, 1971

Fluorocarbon-bridged di(tertiary arsines and phosphines) react with  $M_2(CO)_{10}$  ( $M = Mn, Re$ ) under a variety of conditions to give complexes of formula  $(L-L)M_2(CO)_8$  which are ligand bridged. The bridged complexes readily react with iodine with cleavage of the M-M bond and yield  $(L-L)[M(CO)_4I]_2$ . In the case where  $(L-L)$  is  $(CH_3)_2AsC=CA_s(CH_3)_2CF_2CF_2$  isomers of the bridged complexes can also be prepared. These have the structure  $(CO)_4M(CH_3)_2AsM(CO)_4(CH_3)_2AsC=CCF_2CF_2$  and are the result of a ligand rearrangement reaction.

### Introduction

Reactions of dirhenium and dimanganese decacarbonyls with various monodentate ligands have resulted in the replacement of one to four carbonyl groups with the formation of dinuclear compounds such as  $M_2(CO)_9L$ ,<sup>1-4</sup>  $[M(CO)_4L]_2$ ,<sup>2-11</sup>  $M_2(CO)_7L_8$ ,<sup>3-5</sup> and

$[M(CO)_3L_2]_2$  in which the metal-metal bond of the parent carbonyl is preserved. Bidentate ligands are reported to yield some chelated products of the types  $M(CO)_3(L-L)$ ,<sup>11,12</sup>  $M(CO)(L-L)_2$ ,<sup>11,12</sup>  $M_2(CO)_8(L-L)$ ,<sup>1,13</sup> and  $[M(CO)_3(L-L)]_2$ ,<sup>11,14</sup> in which the ligand replaces two or four carbonyl groups on a single metal atom.

In the course of a systematic study of the reactions of the versatile fluorocarbon-bridged ligands  $f_4fars$  and  $f_4fos$   $YC=CYCF_2CF_2$  ( $Y = (CH_3)_2As$  or  $(C_6H_5)_2P$ ) with metal carbonyls,<sup>15,16</sup> we have found that these ligands react with dimanganese and dirhenium decacarbonyls to

- (1) M. L. Ziegler, H. Haas, and R. K. Sheline, *Chem. Ber.*, **98**, 2454 (1965).
- (2) H. Wawersik and F. Basolo, *Chem. Commun.*, 368 (1966).
- (3) R. J. Clark, J. P. Hargaden, H. Haas, and R. K. Sheline, *Inorg. Chem.*, **7**, 673 (1968).
- (4) J. T. Moelwyn-Hughes, A. W. B. Garner, and N. Gordon, *J. Organometal. Chem.*, **26**, 373 (1971).
- (5) E. Singleton, J. T. Moelwyn-Hughes, and A. W. B. Garner, *ibid.*, **21**, 449 (1970).
- (6) A. G. Osborne and M. H. B. Stiddard, *J. Chem. Soc.*, 634 (1964).
- (7) W. Hieber and W. Freyer, *Chem. Ber.*, **93**, 462 (1960).
- (8) J. Lewis, R. S. Nyholm, A. G. Osborne, S. S. Sandhu, and M. H. B. Stiddard, *Chem. Ind. (London)*, 1398 (1963).
- (9) P. W. Jolly and F. G. A. Stone, *J. Chem. Soc.*, 5259 (1965).
- (10) W. Hieber and W. Freyer, *Chem. Ber.*, **92**, 1765 (1959).
- (11) M. Prenti, D. Giusto, and P. Romiti, *J. Inorg. Nucl. Chem.*, **29**, 761 (1967).

- (12) A. Sacco, *Gazz. Chim. Ital.*, **93**, 698 (1963).
- (13) W. Hieber and W. Schropp, *Z. Naturforsch. B*, **15**, 271 (1960).
- (14) R. S. Nyholm and D. V. Ramana Rao, *Proc. Chem. Soc., London*, 130 (1959).
- (15) W. R. Cullen, D. A. Harbourne, B. V. Liengme, and J. R. Sams, *Inorg. Chem.*, **9**, 702 (1970).
- (16) W. R. Cullen and D. A. Harbourne, *ibid.*, **9**, 1839 (1970).