

of H, the modes involving the former are essentially not visible in the present case, where only 1 in 38 atoms is D and only small samples (~ 2 g) were available. This sample therefore served as a "blank", and its INS spectrum was subtracted from that of the hydrogenated compound to thereby isolate¹³ those vibrations involving mainly the hydrogen atom. The resulting difference spectrum (Figure 1) obtained by using the Be analyzers of the FDS only shows two broad peaks, at approximately 940 and 1100 cm^{-1} in the region of interest. These data are therefore in qualitative agreement with the IR results on Nujol mulls and on CsI disks before additional grinding and pressing was done on the latter. Small differences in the peak positions in the INS data compared with optical data are not uncommon and are often the result of the relatively large value of momentum transfer used in the neutron studies.

It appears therefore that the hydrogen vibrational spectrum is dependent on the nature of the counterion, since the INS data for the Cs^+ -containing compound shows just one line, whereas, in the K^+ analogue, two well-separated INS bands are observed. The former is thus consistent with the approximately centered position of H determined by the crystal structure study (with $[(\text{Ph}_3\text{P})_2\text{N}]^+$ counterion), while the present data suggest that at least some hydrogen atoms are located in less symmetric sites.

We may attempt a consistent assignment of the principal features of the spectra in the present work, the prior INS study,¹⁰ and the IR absorption work¹¹ by assuming that the band(s) in the region 1050–1100 cm^{-1} are assigned to $\nu(\text{Co-H})$ for $\mu_6\text{-H}$. The reason for this assumption is that all experimental data to date have these bands in common. The additional peak observed by INS and IR¹¹ spectroscopy at ca. 950 cm^{-1} would then have to be associated with a different hydride, most likely doubly or triply bridging two or three Co atoms. In either case, the 950- cm^{-1} line should be accompanied by another excitation in the frequency range of these experiments. For $\mu_3\text{-H}$ a doubly degenerate $\nu_{\text{as}}(\text{Co-H})$ should be observed in addition to $\nu_s(\text{Co-H})$. Vibrational data on such systems³ suggest that the two lines should have frequencies within $\pm 20\%$ of each other. There is, however, no strong line within this range about 950 cm^{-1} that could unambiguously be so assigned, but a weak peak at about 1150 cm^{-1} in the IR¹¹ data does show the expected isotropic shift, and the present INS spectrum appears to have a shoulder in this frequency range. Since the INS intensity of the doubly degenerate ν_{as} would

be about twice that of the nondegenerate ν_s , the former would be assigned at 950 cm^{-1} and the latter at 1150 cm^{-1} .

Extensive compilations of data² on $\mu_2\text{-H}$ show that, for a deformation mode ν_2 at 950 cm^{-1} , the antisymmetric stretch ν_3 would be expected to occur at about 1800 cm^{-1} . Since this frequency is well outside the range of the experimental studies in question, a doubly bridging hydride remains a possibility as the second hydrogen species in the K^+ compound.

From the intensities of the INS bands, one may then conclude that somewhat more than half of the hydrogen in $\text{K}[\text{HCo}_6(\text{CO})_{15}]$ is located in the interstitial site with the rest most likely in either $\mu_2\text{-H}$ or $\mu_3\text{-H}$ form on the inside (or possibly outside) of the Co cluster. In the Cs^+ analogue, however, as well as the $[(\text{Ph}_3\text{P})_2\text{N}]^+$ system studied by neutron diffraction, essentially all the hydrogen is found to be in the interstitial site. One is thus led to the conclusion that the $\mu_6\text{-H}$ coordination in these complexes is highly sensitive to the surroundings of the complex such as the type of counterion used in the preparation of the solid.

The essential result of the present INS study then is that there is substantial agreement with those vibrational features that were unambiguously assigned to $\nu(\text{Co-H})$ in the IR work¹¹ on $\text{K}[\text{HCo}_6(\text{CO})_{15}]$ and that vibrational band(s) in the region 1050–1100 cm^{-1} are most likely to be associated with $\mu_6\text{-H}$ in these octahedral Co clusters.

Further work will be aimed at identifying the hydride species that gives rise to the excitation at ca. 950 cm^{-1} , which, in fact, is found¹¹ to disappear upon repeated grinding and pressing of $\text{K}[\text{HCo}_6(\text{CO})_{15}]$ in alkali-metal halide disks. Diffraction studies of the K^+ or Cs^+ salt (instead of $[(\text{Ph}_3\text{P})_2\text{N}]^+$) will be undertaken to locate the noninterstitial hydrogen. It may also be worthwhile to perform an INS study of the Ru analogue, $[\text{HRu}_6(\text{CO})_{18}]^-$, where a two-peak spectrum associated with H was observed,¹⁴ while neutron diffraction work⁶ showed the hydride to be located on the interstitial site. The suggestion that one of these arises from a Fermi resonance could thereby be tested.

Acknowledgment. We wish to thank T. Koetzle for illuminating discussions. A.A. acknowledges support from NATO Grant No. 0068/85. Work at Los Alamos was performed under the auspices of the U.S. Department of Energy, Office of Basic Energy Sciences.

Registry No. $\text{K}[\text{HCo}_6(\text{CO})_{15}]$, 76723-29-0.

(13) Eckert, J. *Physica* **1986**, *136B*, 150.

(14) Oxtou, I. A.; Kettle, S. F. A.; Jackson, P. F.; Johnson, B. F. G.; Lewis, J. J. *Chem. Soc., Chem. Commun.* **1979**, 687.

Additions and Corrections

1988, Volume 27

Boon K. Teo* and Hong Zhang: Magic Numbers in Atom and Electron Counting of Supraclusters Based on Vertex-Sharing Icosahedra.

Page 416. The G_7 , S_7 , B_7 , T_7 , and N_7 values of $s_7(75)$, the pentagonal-bipyramidal array of seven vertex-sharing icosahedra (cf. Figure 1f), should be 75, 67, 43 or 46, 445 or 448, and 890 or 896, respectively.—
Boon K. Teo