

New Aspect on Di- μ -[bis(diphenylphosphino)methane]dichlorotrigold(I) Chloride, Including Its Crystal and Molecular Structure


Ivan J. B. Lin,^{*,†} J. M. Hwang,[†] Da-Fa Feng,[†] M. C. Cheng,[‡] and Yu Wang^{*,‡}

Department of Chemistry, Fu Jen Catholic University, Hsinchuang, Taipei, 242, Taiwan, and
Department of Chemistry, National Taiwan University, Taipei, 107, Taiwan

Received October 6, 1993[⊙]

The reaction of $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ with dppm ($\text{PPh}_2\text{CH}_2\text{PPh}_2$) in ethanol produces $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{Cl}$, **3**. This compound equilibrates with $\text{Au}_2(\text{dppm})\text{Cl}_2$, **1**, and $\text{Au}_2(\text{dppm})_2\text{Cl}_2$, **2**, in solution. ^{31}P NMR studies at various temperatures suggest that the exchange rate between compounds **1** and **3** is faster than that between compounds **2** and **3**. The free chloride anion in **3** is responsible for these exchanges. The strong emission of **3** in solid is attributed to the short Au...Au distances. Crystal structure data for compound **3**·EtOH: space group $C2/c$, with $a = 14.184(3)$ Å, $b = 18.556(6)$ Å, $c = 20.597(6)$ Å, $\beta = 97.56(2)^\circ$, $Z = 4$, and $V = 5285(3)$ Å³.

In the past few years we have been interested in the use of the phase-transfer catalysis technique to prepare ylide complexes¹ of palladium, platinum, and gold. In the course of our work, we chose $\text{Au}_2(\text{dppm})\text{Cl}_2$ (dppm = $\text{Ph}_2\text{PCH}_2\text{PPh}_2$), **1**, as one of our starting materials.^{1b} This common starting material, reported back in 1977 by Schmidbaur,² was prepared by the reaction of "AuCl₃" with dppm in refluxing EtOH. In an attempt to prepare compound **1** by the reaction of $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ with dppm at room temperature, we have been able to identify and isolate at least three different Au(I)-dppm compounds, depending on the reaction conditions. Besides compound **1**, we have isolated the dinuclear gold compound $\text{Au}_2(\text{dppm})_2\text{Cl}_2$, **2**, which was also reported by Schmidbaur³ by the reaction of compound **1** with dppm. The luminescent property of compound **2** has recently attracted considerable interest.⁴ Another Au(I)-dppm complex, a trinuclear gold compound which has the formulation $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{Cl}$, **3**, was also obtained. A similar trinuclear gold compound $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{X}$ (X = ClO_4 , $\text{Au}(\text{C}_6\text{F}_5)_3\text{Cl}$), was reported by Uson⁵ in 1983. Although **3** and Uson's compound have identical cations, the difference in anions causes very large differences in their chemical properties. While Uson's trinuclear gold compound is stable both as a solid and in solution, compound **3** is not. Once dissolved in solution compound **3** equilibrates with compound **1** and **2** very quickly. In this paper we wish to report our interesting discoveries related to these Au-dppm compounds.



Results and Discussion

Synthesis and Characterization. Schmidbaur² reported the synthesis of the colorless compound **1** in 51% yield by refluxing

"AuCl₃" with dppm (1:1 molar ratio) in EtOH. In our work, compound **1** is obtained in 93% yield by the reaction of $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ with dppm at room temperature in a mixture of EtOH/ CH_2Cl_2 / CHCl_3 (1:1:1 by volume). Such a solvent combination is necessary to synthesize **1** in good yield. A ternary solvent system (EtOH/ CHCl_3 / $(\text{CH}_3)_2\text{CO}$) has also been used for the synthesis of $\text{Au}(\text{PPh}_3)\text{Cl}$.⁶ Alternatively reduction of Au(III) to Au(I) by the reaction of $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ with 2,2'-thiodiethanol (thiodiglycol, $(\text{HOCH}_2\text{CH}_2)_2\text{S}$) followed by the reaction with a 0.5 molar ratio of dppm produces **1** in 85% yield. Although **1** is stable in most solvents, it decomposes in DMSO after 1 day. ^{31}P NMR spectroscopy study suggests the formation of a tetranuclear gold compound, $\text{Au}_4(\text{dppm})(\text{dppm-H})\text{Cl}_3$, which has been prepared by a different route and will be published elsewhere.

Preparation of **2** was reported by Schmidbaur *et al*³ in 92% yield by the reaction of **1** with dppm. When $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ simply reacts with a 1.5 molar ratio of dppm in an EtOH/ CH_2Cl_2 / CHCl_3 solvent system, we are able to obtain **2** in 80% yield after recrystallization. In many cases, we isolated a species $\text{Au}_2(\text{dppm})_3\text{Cl}_2$,⁷ which upon dissolving in solvent was difficult to recover. ^{31}P NMR spectrum of this compound in CDCl_3 shows two broad signals assignable to compound **2** and dppm. Attempts to recrystallize this compound give compound **2**.

Synthesis of compound **3** is achieved simply by dissolving $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ in a vast quantity of EtOH in the presence of a 1.2 molar ratio of dppm. Large crystals of **3** are obtained in 61% yield after recrystallization from ethylacetate or acetone. Alternatively, **3** is obtained in 60% yield by reacting $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ with 2,2'-thiodiethanol in EtOH before treating it with dppm. Uson's trinuclear gold-dppm compound was prepared either by the reaction of $\text{Au}_2(\text{dppm})_2\text{Cl}_2$ with $\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}$ or from the

[†] Fu Jen Catholic University.

[‡] National Taiwan University.

[⊙] Abstract published in *Advance ACS Abstracts*, July 1, 1994.

- (1) (a) Lin, I. J. B.; Shen, H. I. *J. Chin. Chem. Soc.* **1992**, *39*, 583. (b) Lin, I. J. B.; Liu, C. W.; Liu, L.-K.; Wen, Y.-S. *Organometallics* **1992**, *11*, 1447. (c) Lai, J. S.; Wu, R. F.; Lin, I. J. B.; Cheng, M. C.; Wang, Y. *J. Organomet. Chem.* **1990**, *393*, 431. (d) Wu, R. F.; Lin, I. J. B.; Lee, G. H.; Cheng, M. C.; Wang, Y. *Organometallics* **1990**, *9*, 126.
- (2) Schmidbaur, H.; Wohlleben, A.; Wagner, F.; Orama O.; Huttner, G. *Chem. Ber.* **1977**, *110*, 1748.

(3) Schmidbaur, H.; Wohlleben, A.; Schubert, U.; Frank, A.; Hutter, G. *Chem. Ber.* **1977**, *110*, 2751.

(4) (a) Khan, M. N. I.; King, C.; Heinrich, D. D.; Fackler, J. P., Jr.; Porter, L. C. *Inorg. Chem.* **1989**, *28*, 2150. (b) King, C.; Wang, J. C.; Khan, M. N. I.; Fackler, J. P., Jr. *Inorg. Chem.* **1989**, *28*, 2145. (c) Che, C.-M.; Kwong, H.-L.; Yam, V. W.-W.; Cho, K.-C. *J. Chem. Soc., Chem. Commun.* **1989**, 885. (d) Che, C.-M.; Kwong, H.-L.; Poon, C.-K.; Yam, V. W.-W. *J. Chem. Soc., Dalton Trans.* **1990**, 3215.

(5) Uson, R.; Laguna, A.; Fernandez, E.; Villacampa, M.; Jones, P. G.; Sheldrick, G. M. *J. Chem. Soc., Dalton Trans.* **1983**, 1679.

(6) McAuliffe, C. A.; Parish, R. V.; Randall, P. D. *J. Chem. Soc., Dalton Trans.* **1979**, 1730.

(7) Anal. Calcd for $\text{C}_{73}\text{H}_{36}\text{Cl}_2\text{P}_6\text{Au}_2$: C, 53.4; H, 4.1. Found: C, 53.3; H, 4.6. The analogous bromo compound was reported: Al-Baker, S.; Hill, W. E.; McAuliffe, C. A. *J. Chem. Soc., Dalton Trans.* **1985**, 2655. The observation of **2** and free dppm in ^{31}P NMR spectrum while dissolving $\text{Au}_2(\text{dppm})_3\text{Cl}_2$ in CDCl_3 is different from that reported by Berners-Price and Sadler.^{11c}

Table 1. Crystal Data for $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{Cl}\cdot\text{EtOH}$ and $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{Cl}\cdot\text{EtOH}\cdot\text{H}_2\text{O}$

formula	$\text{Au}_3\text{P}_4\text{C}_{52}\text{Cl}_3\text{H}_{44}\text{O}_1$	$\text{Au}_3\text{P}_4\text{C}_{52}\text{Cl}_3\text{H}_{46}\text{O}_2$
fw	1506.06	1522.06
diffractometer used	CAD4	CAD4
space group	$C2/c$	$P2_1/n$
a , Å	14.184(3)	16.557(4)
b , Å	18.556(6)	18.886(4)
c , Å	20.257(6)	18.702(7)
β , deg	97.56(2)	113.19(3)
V , Å ³	5285(3)	5194(3)
Z	4	4
$D(\text{calcd})$, g·cm ⁻³	1.893	1.946
$\lambda(\text{Mo K}\alpha)$, Å	0.71069	0.71069
$F(000)$	2848	2880
unit cell detn: no. 2θ range, deg	25; 19–24	24; 18–22
scan type	θ/θ	$\theta/2\theta$
2θ scan width, deg	$2(0.85 + 0.35 \tan \theta)$	$2(0.75 + 0.35 \tan \theta)$
2θ range, deg	2–45	2–45
h, k, l range	–15 to 15, 0 to 20, 0 to 21	–17 to 16, 0 to 20, 0 to 19
$\mu(\text{Mo K}\alpha)$, cm ⁻¹	58.9	87.4
cryst size, mm	$0.2 \times 0.35 \times 0.40$	$0.25 \times 0.30 \times 0.20$
transm factor	0.709–1.0	0.617–1.0
temp, K	298	298
no. of measd reflns	3622	6835
no. of obsd reflns	2648	3491
$(I > 2\sigma(I)) R, R_w^a$	0.030, 0.026	0.045, 0.043
refinement program	NRCVAX ^b	NRCVAX ^b
no. of refined params	286	558
minimized function	$w F_o - F_d ^2$	$w F_o - F_d ^2$
weighting scheme	$1/\sigma^2(F_o)$	$1/[\sigma^2(F_o) + 0.3 \times 10^{-4}F_o^2]$
$g(\text{second. ext coeff}) \times 10^4$	0.46(2)	0.29(3)
$(\Delta/\sigma)_{\text{max}}$	0.514	0.142
$(\Delta\rho)_{\text{max}}, (\Delta\rho)_{\text{min}}, \text{e}\text{Å}^{-3}$	0.84, –0.77	1.16, –0.78
solvent	EtOH	H ₂ O, EtOH

^a $R = [\sum |F_o - F_d|/F_o]$. $R_w = [\sum w(|F_o - F_d|/\sum w(|F_d|^2))]^{1/2}$; $\sigma^2(F_o)$ from counting statistics. ^b NRCVAX ref 13.

reaction of $\text{Au}_2(\text{dppm})\text{Cl}_2$ with AgClO_4 followed by dppm then $\text{AuCl}(\text{C}_4\text{H}_4\text{S})$. Surprisingly, in spite of the similarity in formula between 3 and Uson's compound, 3 has never been reported. This may be due to the instability of 3 in solvent. When dissolved in CHCl_3 or CH_2Cl_2 , 3 is difficult to recover.

Since the characterization of 3 in solution is difficult, single crystal X-ray diffraction was performed. Two different crystals in different space groups were structurally determined. Both crystals contained a solvent molecule, EtOH. One crystal had an additional H₂O molecule as solvent molecule. The crystal data are shown in Table 1. The molecular structures of the cations are the same for both crystals. The atomic parameters of only one crystal (the one without H₂O) are given in Table 2. The molecular structure of the cation, having a C_2 symmetry, is depicted in Figure 1. Some selected bond distances and angles are given in the caption. The three gold atoms are in a V shape arrangement, with two short (3.076(1) Å) and one long (3.729–(1) Å) Au...Au distances. This is compared with the two short Au...Au distances of 3.067(5) and 3.164(5) Å, and the long distance of 3.676 (5) Å of Uson's compound. The luminescent properties of 3 may be attributed to these gold–gold interactions. The Au2–Au1–Au2a angle (74.61(3)^o) (the atom Au2a is related to Au2 by C_2 symmetry) is larger than the one in the literature (72.3^o).⁵ There is a substantial trans influence in Au2–P1 (2.240 Å, trans to chloride) in comparison with Au1–P2 (2.319 Å, trans to phosphorus). Similar results were observed in Uson's trinuclear gold compound. This trans influence may have some implications on the exchange behavior of 3 in solution and will be discussed in a later section. The two P–Au–Cl limbs are pointing toward opposite directions such that further reaction with dppm to close up the trinuclear gold is unlikely. The free chloride anion has no interaction with any of the three gold atoms (>8 Å), although interaction of the free halide with the cation is expected in solution.

Compound 3 is luminescent in the solid state. Excitation of the solid sample at 357 or 366 nm at room temperature results in an intense greenish blue emission with a maximum at 485 nm and a weak shoulder at 525 nm (Figure 2). Both emissions have

Table 2. Atomic Parameters x , y , z , and B_{iso} of $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{Cl}\cdot\text{EtOH}$, where Esds Refer to the Last Digit Printed

	x	y	z	B_{iso} , Å ²
Au1	0	0.20744(3)	$1/4$	2.54(3)
Au2	0.12994(3)	0.33930(3)	0.280257(23)	3.653(22)
P1	0.21188(20)	0.29073(16)	0.20407(14)	3.10(14)
P2	0.04946(19)	0.19499(14)	0.14592(13)	2.41(13)
Cl	0.05038(24)	0.39652(19)	0.35649(18)	6.72(20)
C	0.1333(7)	0.2647(5)	0.1312(5)	2.6(5)
C11A	0.2803(7)	0.2118(5)	0.2302(5)	3.4(5)
C12A	0.3225(7)	0.1703(6)	0.1862(5)	4.0(6)
C13A	0.3712(8)	0.1089(6)	0.2071(6)	5.3(7)
C14A	0.3776(9)	0.0888(7)	0.2724(6)	6.3(7)
C15A	0.3388(9)	0.1300(7)	0.3169(6)	6.0(8)
C16A	0.2914(8)	0.1909(6)	0.2955(5)	4.3(6)
C11B	0.2916(7)	0.3545(5)	0.1713(5)	3.3(5)
C12B	0.2535(8)	0.4164(6)	0.1422(6)	4.9(7)
C13B	0.3113(10)	0.4672(6)	0.1165(6)	6.3(8)
C14B	0.4066(9)	0.4562(7)	0.1234(6)	6.9(8)
C15B	0.4456(9)	0.3951(7)	0.1528(8)	8.7(10)
C16B	0.3897(8)	0.3439(7)	0.1788(7)	6.5(8)
C21A	0.1077(7)	0.1094(5)	0.1394(5)	2.6(5)
C22A	0.1393(8)	0.0897(6)	0.0787(5)	4.6(6)
C23A	0.1843(9)	0.0258(7)	0.0740(6)	6.1(7)
C24A	0.1966(9)	–0.0201(6)	0.1298(7)	6.4(7)
C25A	0.1669(9)	–0.0008(6)	0.1888(5)	5.0(7)
C26A	0.1213(7)	0.0635(5)	0.1943(5)	3.4(6)
C21B	–0.0419(7)	0.1961(5)	0.0744(5)	2.9(5)
C22B	–0.1268(7)	0.1607(6)	0.0784(5)	3.8(5)
C23B	–0.1940(8)	0.1554(6)	0.0246(6)	5.2(6)
C24B	–0.1795(8)	0.1862(6)	–0.0337(6)	5.1(6)
C25B	–0.0960(9)	0.2209(6)	–0.0399(5)	5.3(7)
C26B	–0.0260(8)	0.2257(6)	0.0134(5)	4.7(7)
Cl1	$1/4$	$1/4$	0	13.0(5)
O31 ^a	0.5978(14)	0.0247(10)	0.0316(10)	9.3(6)
O32 ^a	0.5219(20)	0.0221(14)	0.0170(14)	6.4(7)
O33 ^a	0.4542(16)	0.0673(12)	0.0213(11)	4.9(6)

^a Mean atoms with occupancy = 0.5.

identical lifetimes of 11 μs. The luminescent properties of many gold(I) phosphine complexes have been studied. The emissive




Figure 1. Molecular structure of compound 3. The thermal ellipsoids are in 50% probability. For clarity phenyl carbons are not labeled. Complete labels can be found in the deposit material. Selected bond lengths and bond angles are as follows: Au1–Au2–3.076(1) Å, Au1–P2–2.319(3) Å; Au2–P1–2.240(3) Å, Au2–Cl–2.290(3) Å; Au2–Au1–Au2a–74.61(3)°, P2–Au1–P2a–168.6(1)°.




Figure 2. Solid state excitation (---) and emission (—) spectra of [Au₃(dppm)₂Cl₂]Cl at room temperature.

excited states of these compounds have been suggested to arise from the metal-centered transitions,^{4b,8} the gold to phosphine charge-transfer transitions,⁹ or the metal mediated intraligand transitions.^{9a,10} Since symmetric linear bis(phosphine)gold(I) complexes are not luminous and Au(PPh₃)Cl only shows a slight bluish white emission due to $\pi^* \rightarrow \pi$ transition on the phenyl ring, the strong emission of 3 appears to arise from its gold–gold interactions.^{10b} The strong luminescent property of [Au₂(dppm)₂]²⁺ has also been attributed to the short Au–Au distances (<3.0 Å).^{3,4a,b} For a comparison, the emission spectrum of the dinuclear compound Au₂(dppm)Cl₂ was also taken. When excited at 300 nm, a strong emission appeared at 620 nm in the solid state. Again, this compound has a Au–Au distance of 3.351(2) Å.²

NMR Studies. The ³¹P NMR spectrum of 3 in CDCl₃ at 297 K is shown in Figure 3b. It has two very broad signals and a relatively sharp signal at 39.0, 27.6, and 32.6 ppm respectively. There is no correlation between their relative intensities. The ³¹P




Figure 3. ³¹P{¹H} NMR spectra of compound 3 in CDCl₃ at (a) 283 K, (b) 297 K, (c) 307 K, and (d) 317 K.

NMR spectrum of 3 at 297 K in DMSO-*d*₆ is also shown in Figure 4a. The two smaller broad signals at 40.2 and 34.5 ppm are of equal intensity, and the two large sharper signals at 37.4 and 31.4 ppm are also of equal intensity. Low-temperature spectra in CDCl₃ were taken. At 283 K (Figure 3a) a set of AA'BB' pattern (signal III) appears at 38.0 and 29.5 ppm ($J(\text{AA}') = 309$, $J(\text{AB}) = 4.8$, $J(\text{BB}') = -0.2$, $J(\text{AB}') = -1.9$ Hz), together with two singlets at 32.6 ppm (signal II) and 27.2 ppm (signal I). The relative intensity of signal I to II is 1 to 1, but there is no correlation between the intensities of signal III and II (or I). Increasing the temperature to 297 K (Figure 3b), signals I and III begin to merge while signal II remains relatively sharp. At 307 K, all signals become broader. When the temperature is further increased to 317 K, the signals all merge to a broad band. Variable temperature spectra of 3 in DMSO-*d*₆ in the high temperature range are shown in Figure 4. Signals I and III begin to merge at ~307 K, and all signals merge to a broad signal only at 357 K.

Apparently, in both solvents, there are two different stages of line broadening, and the exchange rate between signal I and III is faster than that of II and III. It is also interesting to note that the exchange rates among signals I, II, and III in DMSO-*d*₆ are slower than those in CDCl₃ at the same temperature. In both solvents, the relative intensities of signals I, II, and III are different at different temperatures. Signal I is assigned to compound 1, signal II to compound 2, and signal III to compound 3. The assignments of signals I, II, and III are done in the following manner.

When a 1:1.20 molar ratio of AgBF₄ in proportions is added to a freshly prepared CDCl₃ solution of 3, the ³¹P NMR spectrum changes dramatically (Figure 5). The broad signals become sharp gradually and eventually show a sharp AA'BB' pattern (at 37.1 and 29.0 ppm), very close to that of [Au₃(dppm)₂Cl₂]X reported

- (8) (a) Gray, H. B.; Harvey, P. D. *J. Am. Chem. Soc.* **1988**, *110*, 2145. (b) Che, C.-M.; Yip, H.-K.; Yam, V. W.-W.; Cheung, P.-Y.; Lai, T.-F.; Shieh, S.-J.; Peng, S.-M. *J. Chem. Soc., Dalton Trans.* **1992**, 427. (c) Yam, V. W.-W.; Lee, W.-K. *J. Chem. Soc., Dalton Trans.* **1993**, 2097.
- (9) (a) Ziolo, R. F.; Lipton, S.; Dori, Z. *J. Chem. Soc., Chem. Commun.* **1970**, 1124. (b) Vogler, A.; Kunkely, H. *Chem. Phys. Lett.* **1988**, *150*, 135.
- (10) (a) Segers, D. P.; DeArmond, M. K.; Grutsch, P. A.; Kutal, C. *Inorg. Chem.* **1984**, *23*, 2874. (b) King, C.; Khan, M. N. I.; Staples, R. J.; Fackler, J. P., Jr. *Inorg. Chem.* **1992**, *31*, 3236. (c) Shieh, S.-J.; Li, D.; Peng, S.-M.; Che, C.-M. *J. Chem. Soc., Dalton Trans.* **1993**, 195. (d) Tzeng, B.-C.; Li, D.; Peng, S.-M.; Che, C.-M. *J. Chem. Soc., Dalton Trans.* **1993**, 2365.




Figure 4. $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of compound **3** in $\text{DMSO}-d_6$ at (a) 297 K, (b) 307 K, (c) 327 K, (d) 337 K, (e) 347 K, and (f) 357 K.

by Uson (36.6 and 29.0 ppm).⁵ At the same time, signal II gradually decreases in intensity and the chemical shift moves downfield from 32.6 to 35.6 ppm. The addition of $[\text{Au}_3(\text{dppm})_2\text{Cl}_2](\text{ClO}_4)$ to the solution does not generate any new signal, but the intensity of signal III increases. The addition of $(n\text{-Bu})_4\text{NCl}$ to the above solution, broadens the signals and moves the chemical shift of signal II from 35.6 ppm back to 32.6 ppm again and the intensity increases. These processes are reversible. Therefore signal III is assigned to compound **3**. The lower field signal of III is assigned to the outer phosphorus atoms, which are trans to chlorides, and the higher field signal of III is assigned to the two trans inner phosphorus atoms of **3**. This assignment is done by comparison with compounds **2** and **1** in a similar environment.

When **2** is added to the CDCl_3 solution of **3**, no additional signal appears, except that the relative intensity of signal II to III increases. The chemical shift of pure **2** in CDCl_3 is also identical to that of signal II. Excess chloride ions has no effect on the chemical shift of **2**. The influence of chlorides and perchlorate anions on the chemical shift of $[\text{Au}_2(\text{dppm})_2]^{2+}$ was reported.^{4c} We, therefore, assign signal II to compound **2**. The slight peak which appears at ~ 27.6 ppm can possibly be derived from compound **1**, but its nature is not certain.

^{31}P NMR spectrum of pure compound **1** in CDCl_3 at room temperature appears at 23.5 ppm, which is 3.1 ppm upfield relative to signal I in Figure 3a. However, when compound **1** is added to the CDCl_3 solution of **3** at 283 K, no additional signal appears, except that the relative intensity of signal I to II increases. We tentatively assign signal I to compound **1**. The large chemical shift difference between pure compound **1** in CDCl_3 and **1** in the above mentioned solution can possibly be due to the influence of




Figure 5. $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of compound **3** in CDCl_3 at 297 K, with various molar ratios of **3** to AgBF_4 : (a) no AgBF_4 ; (b) 1:0.25; (c) 1:0.50; (d) 1:1.00; (e) 1:1.20.




Figure 6. Change of chemical shift of $\text{Au}_2(\text{dppm})\text{Cl}_2$ (8 mg, 9.42×10^{-3} mmol in 400 mL of CDCl_3) with each increment of 0.64×10^{-3} mmol of $n\text{-Bu}_4\text{NCl}$ in 10 mL of CDCl_3 .

the free halide ion. In order to support our proposal, the chemical shifts of compound **1** under various amounts of chloride ions are determined. Indeed, the chemical shift of compound **1** moves gradually downfield upon addition of $(n\text{-Bu})_4\text{NCl}$ (Figure 6). The relationship, however, is not linear. The influence of the chloride ion on the chemical shift of **1** suggests that there is an interaction between **1** and the chloride anion. Interaction of halides with linear gold (I) compounds has been reported.^{4c,11}

To further support our assignment, compound **1** and **2** is mixed (2:1 molar ratio) in CDCl_3 at 297 K. The ^{31}P NMR spectrum






Figure 7. Plot of $\ln K$ versus $1/T$ for the equilibrium between compounds 1, 2, and 3 in CDCl_3 .

shows signals identical to those of 3 dissolved in CDCl_3 . This is not true for a mixture of 1 and $[\text{Au}_2(\text{dppm})_2](\text{ClO}_4)_2$ which has no halide anions. Thus, while 3 in CDCl_3 can be converted to compounds 1 and 2, compound 1 with compound 2 can also be reverted to compound 3 (eq. 1). It appears that compound 3 is in equilibrium with compounds 2 and 1. Equilibrium constants in CDCl_3 at various temperatures are measured ($K_{297} = 1.33 \times 10^{-3}$, $K_{292} = 9.24 \times 10^{-4}$, $K_{287} = 5.46 \times 10^{-4}$, $K_{282} = 3.06 \times 10^{-4}$, $K_{277} = 1.95 \times 10^{-4}$, $K_{272} = 9.10 \times 10^{-5}$). A plot of $\ln K$ versus $1/T$ gave a straight line (Figure 7). Using the Gibbs free energy equation, ΔH° (80.89 ± 2.63 kJ/mol) and ΔS° (222.86 ± 0.68 J/(mol·K)) are obtained. These values are consistent with our observation that, at low temperature, equilibrium favors the formation of 3. The variation of the spectral intensity upon gradual addition of AgBF_4 in Figure 4 can also be explained by the equilibrium. Among the three compounds, 3 will be the first compound to have its Cl^- removed by AgBF_4 . 3 is followed by 2 and then 1, according to the strength of these Au–Cl bonds. The equilibrium will shift the reaction of eq 1 toward the formation of 3 to increase the concentration of Cl^- . Therefore the amount of $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{BF}_4$ will gradually increase, while 2 and 1 will gradually decrease upon addition of AgBF_4 . Addition of chloride ion will shift the reaction of eq 1 toward the formation of 2 and 1.



Apparently, the driving force for the instability of 3 in solution is the reaction of chloride anion with the trinuclear gold compound. The weakly bonded chloride anion in 2 is also responsible for the reaction of 1 and 2 to yield 3. A plausible pathway for these interconversion is shown in Scheme 1. While each of the noncentral gold atom bonds to a neutral phosphine and an anionic chloride ligand, the central gold atom has two neutral phosphine ligands, suggesting that the central gold atom might bear more positive charge and hence be more susceptible to be attacked by a nucleophile. The longer central Au1–P2 bond length and the shorter noncentral Au2–P1 bond length also suggest that the central Au1–P2 bond may be more reactive than the noncentral Au2–P1 bond. An attack of a chloride anion on the central gold atom will produce compound 1 and $\text{Au}(\text{dppm})\text{Cl}$. Combining two molecules of $\text{Au}(\text{dppm})\text{Cl}$ will generate 2 through several steps. The equilibrium between $\text{Au}(\text{dppm})\text{Cl}$ and 2 was proposed.^{11c} Thus, while compound 1 can be produced directly from 3, compound 2 requires at least two more steps to be formed. These reactions are all reversible. Therefore, one should expect that the interconversion rate between compound 3 and 1 is faster

Scheme 1



than that between compound 3 and 2. This argument is consistent with our observation that signals I and III merge before signals II and III, and the exchange rates depend on the concentration of chloride ion. The spectral change of 3 upon addition of AgBF_4 , and the spectral changes of $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{BF}_4$ upon addition of Cl^- are also consistent with the proposed scheme. DMSO, which has a much higher dielectric constant (45) than chloroform (4.8),¹² has better ability to separate ions. This mechanism also explains the slower exchange rates in DMSO.

In conclusion, we have demonstrated that the instability of 3 in solution is due to the presence of the free chloride anion. In solution, 3 equilibrates with 1 and 2; therefore, the characterization of 3 in solution is difficult. This is the reason why that 3 has not been reported. The possible isolation of 3 in this work is due to the proper choice of solvent, such that both the equilibrium and the solubility favor its isolation.

Experimental Section

³¹P{¹H} NMR spectra were recorded on a Bruker AC-F300 spectrometer (at 121.5 MHz). Chemical shifts are reported relative to an external standard 85% H_3PO_4 , with downfield shifts being positive. Spectral simulation of the AA'BB' pattern was done by the PANIC of Bruker. Room-temperature solid emission spectrum was recorded on a Aminco-Bowman luminescence spectrometer. Emission lifetime were measured by using an Edinburgh Analytical Instruments CD900 spectrometer. Microanalyses were performed by Taiwan Instrumentation Center. $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ was purchased from Kanto Chemicals. Dppm was purchased from Strem Chemicals. 2,2'-Thiodiethanol (thiodiglycol, $(\text{HOCH}_2\text{CH}_2)_2\text{S}$) was purchased from Riedel-deHaën Laboratory Chemicals.

The equilibrium constants at different temperatures were obtained as described below. $[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{Cl}$ (32.0 mg, 2.18×10^{-2} mmol) was dissolved in CDCl_3 (250 mL). ³¹P NMR spectra of the sample were taken at 297, 292, 287, 282, 277, and 272 K. The concentrations of compounds 1, 2, and 3 were calculated by the integrations of the individual signal. The equilibrium constants, K , were calculated by the following equation:

$$K = \frac{[\text{Au}_2(\text{dppm})_2\text{Cl}_2][\text{Au}_2(\text{dppm})\text{Cl}_2]^2}{[\text{Au}_3(\text{dppm})_2\text{Cl}_2]^2[\text{Cl}]^2}$$

This equation produced the best linear fit of $\ln K$ versus $1/T$. However, this was not the case for the K values obtained in DMSO; presumably, partial dissociation of chloride in compound 2 occurred.

$\text{Au}_2(\text{dppm})\text{Cl}_2$ (1). This compound can be prepared by the method reported by Schmidbaur² or by the following two different methods: (a)

(11) (a) Jaw, H.-R.; Savas, M. M.; Rogers, R. D.; Mason, W. R. *Inorg. Chem.* 1989, 28, 1028. (b) Shain, J.; Fackler, J. P., Jr. *Inorg. Chim. Acta* 1987, 131, 157. (c) Berners-Price, S. J.; Sadler, P. J. *Inorg. Chem.* 1986, 25, 3822.

(12) Huheey, J. E. *Inorganic Chemistry*, 3rd ed.; Harper & Row: New York, 1983, p 340.

(13) Gabe, E. J.; Le page, Y.; Charland, J.-P.; Lee, F. L.; White, P. S. J. *Appl. Crystallogr.* 1989, 22, 384.

$\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ (313 mg, 0.76 mmol) in EtOH (20 mL) was mixed with dppm (300 mg, 0.78 mmol) in CH_2Cl_2 (20 mL). After the suspended solution was stirred for 30 min., CHCl_3 (20 mL) was added, and was stirred for another hour. The colorless resultant solution was set aside until the volume was reduced to ~20 mL, at which time, large colorless crystals were formed. This was filtered and dried in air. The yield was 93%. (b) 2,2'-Thiodiethanol (0.2 mL, excess) was added to $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ (121 mg, 0.294 mmol) in EtOH (10 mL), and the mixture was stirred at 40 °C for 40 min at which time the yellow solution became colorless. To this resultant solution dppm (56 mg, 0.147 mmol) in CH_2Cl_2 (10 mL) was added. After stirring for an additional 1 h, the white precipitates were filtered and then recrystallized from CH_2Cl_2 . Large colorless crystals were obtained in 85% yield. The identification of this compound was confirmed by EA, FAB/MS, mp, and single-crystal X-ray diffraction.

$\text{Au}_2(\text{dppm})_2\text{Cl}_2$ (2). This compound can be prepared by the method reported by Schmidbauer³ or by the following method: dppm (461 mg, 1.20 mmol) in CH_2Cl_2 (20 mL) was added to $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ (313 mg, 0.76 mmol) in EtOH (20 mL). This solution was stirred for 30 min, and then CHCl_3 (20 mL) was added and the mixture was stirred for another 30 min. The resultant solution was dried under vacuum. The residue was then recrystallized from ethyl ester. Colorless crystals were collected in ~80–90% yield. This compound was identified by mp and EA. The corresponding $[\text{Au}_2(\text{dppm})_2](\text{ClO}_4)_2$ was prepared by the literature method.^{4c,7}

$[\text{Au}_3(\text{dppm})_2\text{Cl}_2]\text{Cl} \cdot \text{EtOH} \cdot \text{H}_2\text{O}$ (3·H₂O). This compound was prepared by two different methods: (a) Solid dppm (384 mg, 1.00 mmol) was added to $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ (330 mg, 0.80 mmol) in EtOH (50 mL). This suspended solution was stirred for 30 min, during which time the color changed from a deep yellow to a light yellow. The solvent was then dried completely with a rotary evaporator, and the solid was washed several times with ethyl ether. The residue was dissolved in ethyl acetate or acetone together with a few drops of ethanol and was allowed to evaporate

slowly. Large colorless crystals were obtained in 61% yield. (b) $\text{HAuCl}_4 \cdot 4\text{H}_2\text{O}$ (93 mg, 0.226 mmol) in EtOH (10 mL) was added to an excess of 2,2'-thiodiethanol (0.2 mL) and was stirred at 40 °C for 40 min. To this resultant colorless solution was added dppm (58 mg, 0.151 mmol) in CH_2Cl_2 (10 mL). After being stirred for another hour, the solution evaporated to an oily liquid. Upon addition of ether (10 mL) to this oily liquid, white crystals precipitated. This white solid was filtered and redissolved in a minimum amount of acetone or ethyl acetate together with a few drops of ethanol. Large colorless crystals were collected after the solvent was allowed to evaporate slowly to half of its volume. The yield was 60%. The purity of this compound can be checked with a portable UV lamp. The compound showed a greenish blue emission. Mp: 205–206 °C. Anal. Calcd for $\text{C}_{52}\text{H}_{50}\text{Cl}_3\text{OP}_4\text{Au}_3$: C, 41.3; H, 3.3. Found: C, 41.7; H, 3.2.

X-ray Diffraction of 3. Suitable crystals were chosen to do the single-crystal structure determinations. A crystal of 3·EtOH was obtained as described in the preparation method. Compound 3·EtOH·H₂O was obtained from the crystallization of 3 in ethanol solution. The X-ray diffraction data were measured on a four-circle diffractometer. Intensities of three standard reflections were monitored every hour throughout the data measurement. The variation was less than 2%. The other essential details of single-crystal data measurement and refinement are given in Table 1.

Acknowledgment. This work was supported by the National Science Council of Taiwan, NSC82-0618-M-030-013.

Supplementary Material Available: Tables giving complete positional and isotropic thermal parameters, anisotropic thermal parameters, and bond distances and angles and an ORTEP drawing of 3·EtOH with complete labels (4 pages). Ordering information is given on any current masthead page.