

Tetramethylammonium Difluorobromate(I), $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$

Rolf Minkwitz* and Raimund Bröchler

Anorganische Chemie, Fachbereich Chemie der Universität Dortmund, 44221 Dortmund, Germany

Ralf Ludwig

Physikalische Chemie, Fachbereich Chemie der Universität Dortmund, 44221 Dortmund, Germany

Received October 24, 1996[®]

The thermolysis of $(\text{CH}_3)_4\text{N}^+\text{Br}(\text{OCF}_3)_2^-$ between -70 and -10 °C, evolving F_2CO during the decomposition, gave $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$. The characterization of $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$ was carried out by IR, Raman, and ^{19}F - and ^{13}C -NMR. For BrF_2^- two infrared absorption bands at 236 and 450 cm^{-1} and one band in the Raman spectrum at 460 cm^{-1} were observed in accord with a centrosymmetric structure. Ab initio calculations are presented for BrF_2^- the isoelectronic KrF_2 , and other related compounds with $D_{\infty h}$ symmetry.

Introduction

Fluorohalogenates $[\text{XF}_2]^-$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) are well-known from the literature. However, a controversy about their vibrational spectroscopy is still going on.^{1–6}

The first triatomic fluorohalogenate, ClF_2^- , was prepared by Christe and Guertin in 1965.¹ It was based on the reaction of ClF with NOF and identified by infrared spectroscopy. Later the full characterization of ClF_2^- was achieved via Raman and IR spectra with cesium, rubidium, and potassium as cations.² The theoretically predicted linear structure with $D_{\infty h}$ symmetry was apparently not observed. The IR spectra showed an intense band attributed to $\nu_s(\text{Cl}-\text{F})$, which is forbidden for a linear centrosymmetric structure. The difference between the theoretically predicted $D_{\infty h}$ symmetry and the experimental results was explained as a result of crystal field effects.

First attempts to prepare IF_2^- were described by Meinert and Klamm.⁷ Unfortunately no spectroscopic data were reported. Further attempts to synthesize IF_2^- failed.^{8,9} Only disproportionation products such as I_2 and Cs_3IF_6 were obtained.⁸

The synthesis and full characterization of $(\text{C}_2\text{H}_5)_4\text{N}^+\text{IF}_2^-$ was recently performed by Naumann and Meurer, by the direct reaction of IF with $(\text{C}_2\text{H}_5)_4\text{N}^+\text{F}^-$.³ Two IR- and one Raman-active frequencies were observed in the vibrational spectra. This is consistent with a linear structure of symmetry $D_{\infty h}$.

The first attempts to prepare BrF_2^- from CsF and BrF were described by Surles et al.⁴ The Raman spectrum of the reaction product showed four frequencies instead of the one frequency expected for $D_{\infty h}$ symmetry. Surles et al. explained their data with symmetry reduction from $D_{\infty h}$ to C_{2v} . The synthesis of $\text{Cs}^+\text{BrF}_2^-$ by codepositing CsBr salt with an argon/fluorine mixture was described by Miller and Andrews.⁶ In the IR spectrum of the deposit, two bands at 533 and 527 cm^{-1} were

attributed to the BrF_2^- anion, while the IR-active deformation frequency was not observed.

Although many attempts to prepare BrF_2^- were made since 1965, the anion could never be isolated and its symmetry is still unclear because of conflicting spectroscopic data.

For the preparation of BrF_2^- , alkali-metal cations should not be used as counterions to avoid distortions of the anion which are well described in the literature.^{2–4}

Recently we reported the preparation of the new bromate(I) salt $(\text{CH}_3)_4\text{N}^+(\text{CF}_3\text{O})_2\text{Br}^-$.¹⁰ The compound was synthesized by the reactions of $(\text{CH}_3)_4\text{N}^+\text{Br}^-$ or $(\text{CH}_3)_4\text{N}^+\text{BrCl}_2^-$ with an excess of CF_3OCl at -78 °C. It was assumed, that the salt might be a suitable precursor for $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$.

Experimental Section

All synthetic work and sample handling were performed by employing standard Schlenk techniques and a standard glass vacuum line. The glass vacuum line and the reaction vessels were dried by vacuum and treated with CF_3OCl . Nonvolatile materials were handled under dry nitrogen. CF_3OCl was prepared, according to the literature method, from F_2CO with ClF and a CsF catalyst.^{22,23} The infrared spectra were recorded on a Bruker IFS 113v FT-IR spectrometer in a cooled cell²⁴ equipped with CsBr plates.

- Minkwitz, R.; Bröchler, R. *Z. Anorg. Allg. Chem.* **1997**, 623, 487.
- Wilson, W. W.; Christe, K. O. *Inorg. Chem.* **1989**, 28, 4172.
- Lustig, M.; Ruff, J. K. *Inorg. Chem.* **1966**, 5, 2124.
- Claassen, H. H.; Goodman, G. L.; Malm, J. G.; Schreiner, F. J. *Chem. Phys.* **1965**, 42, 1229.
- Herzberg, G. *Infrared and Raman Spectra of Polyatomic Molecules*; D. Van Nostrand Co., Inc.: Princeton, NJ, 1945; S. 274, 311.
- Overend, J. *Infrared Spectra and Molecular Structure*; Davies, M., Ed.; Amer. Elsevier: New York, 1963, S. 352.
- Christe, K. O.; Naumann, D. *Inorg. Chem.* **1973**, 12, 59.
- Claassen, H. H.; Chernick, C. L.; Malm, J. G. *J. Am. Chem. Soc.* **1963**, 85, 1927.
- Kalinowski, H.-J.; Berger, S.; Braun, S. *¹³C-NMR Spektroskopie*; Georg Thieme Verlag: Stuttgart, New York, 1984; p 201.
- Simons, W. W., Ed. *The Sadtler Guide to Carbon-13 NMR-Spectra*; Sadtler-Heyden: London, 1983; S. 279.
- Hesse, M.; Meier, H.; Zeeh, B. *Spektroskopische Meth. in der organischen Chemie*; Georg Thieme Verlag: Stuttgart, New York, 1987; Vol. 3, p 179.
- Wilson, W. W.; Christe, K. O. *Inorg. Chem.* **1989**, 28, 4172.
- Gould, D. E.; Anderson, L. R.; Young, D.; Fox, W. B. *J. Am. Chem. Soc.* **1969**, 91, 1310.
- Schack, C. J.; Maya, W. *J. Am. Chem. Soc.* **1969**, 91, 2902.
- Bayersdorfer, L.; Minkwitz, R.; Jander, J. *Z. Anorg. Allg. Chem.* **1976**, 392, 137.

[®] Abstract published in *Advance ACS Abstracts*, August 15, 1997.

- Christe, K. O.; Guertin, J. P. *Inorg. Chem.* **1965**, 4, 905.
- Christe, K. O.; Sawodny, W.; Guertin, J. P., *Inorg. Chem.* **1967**, 6, 1159.
- Naumann, D.; Meurer, A. *J. Fluor. Chem.* **1995**, 70, 83.
- Surles, T.; Quarterman, L. A.; Hyman, H. H. *Z. Anorg. Allg. Chem.* **1973**, 35, 668.
- Baran, E. J. *Z. Naturforsch.* **1973**, 28b, 502.
- Miller, J. H.; Andrews, L. *Inorg. Chem.* **1979**, 18, 988.
- Meinert, H.; Klamm, H. *Z. Chem.* **1965**, 5, 468.
- Meinert, H. *Habilitation*, Humboldt-Universität Berlin, 1968.
- Schmeisser, M.; Sartori, P.; Naumann, D. *Chem. Ber.* **1970**, 103, 880.

The Raman spectra were recorded on a T 64000 Jobin Yvon, using the 514.5 nm exciting line of an Ar⁺ laser tube (Spectra Physics). Owing to their hygroscopic nature, the BrF₂⁻ salts were measured in the dry nitrogen atmosphere of a duran glass cell.²⁵

The NMR spectra of (CH₃)₄N⁺BrF₂⁻ in CH₃CN solution were recorded at -38 °C on a Bruker AM 300 with CFCl₃ as external reference.


Preparation of (CH₃)₄N⁺BrF₂⁻. A dry 35 mL cap glass vessel containing 1 g (3.08 mmol) of tetramethylammonium bis(trifluoromethoxy)bromate(I) (CH₃)₄N⁺(CF₃O)₂Br⁻ was slowly warmed from -78 to -10 °C. Carbonyl difluoride, which is formed during the thermolysis, was removed continuously. The completion of the reaction was checked by weighing the reaction product. (CH₃)₄N⁺BrF₂⁻ is a colorless solid which is extremely sensitive toward hydrolysis and can be stored without decomposition at -70 °C for several weeks. At room temperature in a disproportionation reaction the solid decomposes into Br⁻ and BrF₄⁻, whereas the solution decomposes at -35 °C.

Ab Initio Calculations. Ab initio calculations for BrF₂⁻ were performed with the Gaussian 94 program²⁷ at the Hartree-Fock level using standard 3-21G*, 6-31*, 6-31+G*, and 6-311G** basis sets. Including electron correlation, the anion was also calculated at the MP2 level of theory and with a 6-31+G* basis set. Harmonic vibrational frequencies were computed for the minimum-energy structures and scaled by empirical factors to maximize their fit with the experimentally observed frequencies. The calculated frequencies for all basis sets are corrected throughout by 10% on the Hartree-Fock level^{28,29} and by 5% when electron correlation is taken into account.³⁰

For a consistent comparison with the isoelectronic KrF₂ and other related compounds, only calculations on a small 3-21G* basis set could be performed. However, it is well-known that a 3-21G* basis set is already quite successful in accounting for hypervalent structures. The mean absolute deviations of calculated from experimental bond lengths between heavy atoms are only 1.5 pm for this basis set which is the same as for the larger 6-31G* basis. The same is true for calculated frequencies where the scatter in the 3-21G* data is much less than that resulting from the 3-21G basis.³¹

Results and Discussion

Formation and Stability. The thermolysis of (CH₃)₄N⁺(CF₃O)₂Br⁻ in the temperature range -70 to -10 °C (eq 1)



yields tetramethylammonium difluorobromate(I), (CH₃)₄N⁺BrF₂⁻. (CH₃)₄N⁺BrF₂⁻ can be stored at -70 °C for several weeks without decomposition. The salt disproportionates at room temperature to (CH₃)₄N⁺BrF₄⁻ and Br₂.

Vibrational Spectra. The Raman and infrared spectra of (CH₃)₄N⁺BrF₂⁻ are shown in Figure 1. The observed frequencies are listed in Table 1 along with their assignments. Figure 2




Figure 1. Vibrational spectra of (CH₃)₄N⁺BrF₂⁻: Trace A, spectrum of the solid on CsBr plates at -70 °C; trace B, Raman spectrum of the solid at -78 °C (cation bands marked with an asterisk).

Table 1. Raman and Infrared Frequencies^a of (CH₃)₄N⁺BrF₂⁻

Raman	IR	assignt
470 (sh)		2δ(F-Br-F)
460 (100)		ν _s (Br-F)
	450 (vs, br)	ν _{as} (Br-F)
	236 (m)	δ(F-Br-F)
79 (96)		lattice
66 (70)		lattice
	3048 (w)	ν(CH ₃)
3036 (69)	3035 (w)	ν(CH ₃)
2958 (43)	2964 (w)	ν(CH ₃)
2913 (14)		ν(CH ₃)
2829 (11)		2δ _s (CH ₃)
1474 (27)	1483 (sh)	δ _{as} (CH ₃)
1449 (8)	1455 (vs)	δ _{as} (CH ₃)
1412 (8)	1424 (m)	δ _s (CH ₃)
	1399 (vw)	ρ(CH ₃)
1290 (5)	1288 (w)	ρ(CH ₃)
1178 (6)	1181 (w)	ρ(CH ₃)
	1033 (m)	ρ(CH ₃)
950 (26)	953 (vs)	ν _{as} (C ₄ -N)
759 (32)	750 (w)	ν _s (C ₄ -N)
	709 (m)	
	470 (sh)	δ _{as} (C ₄ -N)
	455 (m)	δ _s (C ₄ -N)
377 (2)		δ(C ₄ -N)

^a Frequencies in cm⁻¹.

shows expanded sections of the Raman spectrum of (CH₃)₄N⁺BrF₂⁻ (A) as well as of the decomposition spectrum of BrF₂⁻ (B). Figure 3 depicts a correlation diagram of the experimental and calculated frequencies.

The assignment for the normal modes of the tetramethylammonium cation is obtained from comparison with tetramethylammonium tetrafluorohalogenates(III) (CH₃)₄N⁺ClF₄⁻

(25) Werner, A. Dissertation, Universität Dortmund, 1988.

(26) Hyman, H. H. *Science* **1964**, *145*, 773.

(27) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T. A.; Peterson, G. A.; Montgomery, J. A.; Raghavachari, K.; Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Ciolowski, J.; Stefanov, B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Steward, J. J. P.; Head-Gordon, M.; Gonzales, C.; Pople, J. A. *G94 Gaussian 94 (revision A.1)* A; Gaussian, Inc.: Pittsburgh, PA, 1995.

(28) Pulay, P.; Forgarasi, G.; Pang, F.; Boggs, J. E. *J. Am. Chem. Soc.* **1979**, *101*, 2550.

(29) Pople, J. A.; Schlegel, H. B.; Krishnan, R.; Freese, D. J.; Binkley, J. S.; Frisch, M. J.; Whiteside, R. A.; Hout, R. F.; Hehre, W. J. *Int. J. Quant. Chem.* **1981**, *15*, 269.

(30) Yamaguchi, Y.; Schaeffer, H. F. *J. Chem. Phys.* **1980**, *73*, 2310.

(31) For an authoritative description of the ab initio methods and basis sets employed here, see: Hehre, W. G.; Radom, L.; Schleyer, P. v. R.; Pople, J. A. *Ab initio Molecular Orbital Theory*; Wiley: New York, 1986.




Figure 2. Trace A: Scale expansion of the Raman spectrum of $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$ and assignment of the resonances. Trace B: Partial decomposition of $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$. Beside $\nu_s(\text{Br-F})$ and the first overtone $\delta(\text{F-Br-F})$ of BrF_2^- , disproportionation product BrF_4^- is seen (cation bands marked with an asterisk).




Figure 3. Correlation diagram of the experimental frequencies of BrF_2^- , KrF_2 , and other related compounds with $D_{\infty h}$ symmetry (A) compared with the frequencies calculated with a HF-3-21G* basis set (B).

and $(\text{CH}_3)_4\text{N}^+\text{BrF}_4^-$,¹¹ as well as $(\text{CH}_3)_4\text{N}^+\text{I}(\text{NO}_3)_2^-$, $(\text{CH}_3)_4\text{N}^+\text{I}(\text{NO}_3)_4^-$, and $(\text{CH}_3)_4\text{N}^+\text{Br}(\text{NO}_3)_2^-$.¹²

For a linear, centrosymmetric triatomic ion such as XY_2^- three normal modes of vibration, classified as $(\Sigma_g^+ + \Pi_u + \Sigma_u^+)$, are expected, where ν_2 (Π_u) and ν_3 (Σ_u^+) are only infrared active and ν_1 (Σ_g^+) is only Raman active.

The recorded Raman spectrum of $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$ shows an intense line at 460 cm^{-1} which is assigned as the symmetric Br-F stretch. A comparison of $\nu_s(\text{Br-F})$ with $\nu_s(\text{Kr-F})$ for the isoelectronic Krypton difluoride¹³ at 462 cm^{-1} yields good agreement.

A shoulder on the high-frequency side of the symmetric BrF_2^- stretching mode is attributed to the first overtone of the

deformation mode whose intensity is enhanced by Fermi resonance (Figure 2A). Under the assumption of Fermi resonance,¹⁴ the unperturbed frequencies of ν_1 and $2\nu_2$ are expected at 463 and 467 cm^{-1} , respectively.¹⁵ In contrast Claassen et al. observed no Fermi resonance for the isoelectronic krypton difluoride and explained their findings by the large difference of about 16 cm^{-1} between ν_1 and $2\nu_2$.¹³ The intense lines at 77 and 66 cm^{-1} in the Raman spectrum are attributed to lattice vibrations.

In the infrared spectrum of $(\text{CH}_3)_4\text{N}^+\text{BrF}_2^-$ the antisymmetric Br-F stretching mode appears at 450 cm^{-1} and the bending mode was observed at 236 cm^{-1} . The bending frequency of 236 cm^{-1} is in good agreement with that of 232.6 cm^{-1} observed for the isoelectronic krypton difluoride, whereas the antisymmetric stretching mode exhibits a significant shift toward smaller wavenumbers.¹³

This effect is well-known for pentatomic fluorohalogenates. A comparison of IF_4^- ¹⁶ with the isoelectronic XeF_4 ¹⁷ shows that the antisymmetric stretching mode of the anion is 138 cm^{-1} lower than that of the isoelectronic, neutral compound. The antisymmetric stretching mode also has a lower frequency than the symmetric stretching mode.

The vibrational spectroscopic data of BrF_2^- clearly suggest a linear anion with $D_{\infty h}$ symmetry. The large tetramethylammonium cations are preventing interanionic contacts and distortion of the linear structure. This observation is in agreement with the results of Naumann and Meurer, who obtained a linear IF_2^- anion with tetraethylammonium as counterion.³

Ab Initio Calculations. Ab initio calculations are of great help for the assignment of the BrF_2^- resonances in the experimental infrared and Raman spectra. In Table 2 and Figure 3, BrF_2^- is compared to isoelectronic KrF_2 and other related species with $D_{\infty h}$ symmetry. Unfortunately structures with atoms larger than Br could be calculated only with a small 3-21G* basis set. However, this basis set is known to yield quite reasonable geometries and frequencies for hypervalent structures. For consistency, all species were calculated with this basis set. For all species only positive force constants were obtained (see Table 3). This result is in contradiction to calculated force constants for BrF_2^- ⁵ and KrF_2 ¹³ using the FG-matrix method, which due to the underdetermined nature of the problem could not yield well-determined interaction constants.

As can be seen from Figure 3, the $\delta(\text{F-X-F})$ ($X = \text{Cl}, \text{Br}, \text{I}$) frequencies are in acceptable agreement with the available experimental data. The trend of increasing frequencies on going from IF_2^- to ClF_2^- allows the prediction of the $\delta(\text{F-Cl-F})$ in ClF_2^- , which could not be observed by Christie et al. because the spectra were recorded only for wavenumbers larger than 400 cm^{-1} . The calculated symmetric and antisymmetric stretching modes are in agreement with the experimental spectra of BrF_2^- and IF_2^- . In particular the calculations indicate that both frequencies lie close together within 30 cm^{-1} and $\nu_{as}(\text{Br-F})$ may be smaller than $\nu_s(\text{Br-F})$. Naumann and Meurer³ found the opposite behavior for IF_2^- , with $\nu_{as}(\text{I-F})$ being observed at higher frequency than $\nu_s(\text{I-F})$. On the basis of our own observations for BrF_2^- and other fluorohalogenates, we should point out that the accurate determination of the experimental frequencies of the antisymmetric stretching mode is difficult because of the large bandwidths in the IR spectra. This may cause uncertainties of about 20 cm^{-1} in the experimental values of ν_3 .

The consistency of the calculated results definitely allows us to rule out the experimental assignment for $\nu_{as}(\text{F-Cl-F})^2$ at 661 cm^{-1} . Our calculations indicate that the intense IR band

Table 2. Experimental and Calculated Raman and Infrared Frequencies^a and Calculated Bond Distances^b and Energies^h of (CH₃)₄N⁺BrF₂⁻, Compared to Those of Isoelectronic KrF₂ and Other Related Compounds with *D_{∞h}* Symmetry

Rb ⁺ ClF ₂ ^{-c}		Me ₄ N ⁺ BrF ₂ ⁻		Et ₄ N ⁺ IF ₂ ^{-d}		KrF ₂ ^e		XeF ₂ ^f		assgnt
IR	Raman	IR	Raman	IR	Raman	IR	Raman	IR	Raman	
661		450		462		588		557		$\nu_{as}/\nu_3 (\Sigma_u^+)$
470	476		470 ⁱ				462		496	$2\delta/2\nu_2$
		236	460 ⁱ	206	445	232		213		$\nu_s/\nu_1 (\Sigma_g^+)$
			79/66		111				108	$\delta/\nu_2 (\Pi_u)$ lattice
ClF ₂ ^{-g}		BrF ₂ ^{-g}		IF ₂ ^{-g}		KrF ₂ ^g		XeF ₂ ^g		assgnt
IR	Raman	IR	Raman	IR	Raman	IR	Raman	IR	Raman	
479		480		494		649		635		$\nu_{as}/\nu_3 (\Sigma_u^+)$
	487		489		501		558		583	$\nu_s/\nu_1 (\Sigma_g^+)$
317		250		219		267		236		$\delta/\nu_2 (\Pi_u)$
Calculated Bond Distances ^b										
ClF ₂ ^{-g}		BrF ₂ ^{-g}		IF ₂ ^{-g}		KrF ₂ ^g		XeF ₂ ^g		
182.1		190.9		202.4		182.6		194.8		
Calculated Energies ^h and Signs Reversed										
ClF ₂ ^{-g}		BrF ₂ ^{-g}		IF ₂ ^{-g}		KrF ₂ ^g		XeF ₂ ^g		
655.11591		2758.03025		7085.72668		2936.97222		7398.44800		

^a Frequencies in cm⁻¹. ^b Bond distances in pm. ^c Data from ref 2. ^d Data from ref 3. ^e Data from ref 13. ^f Data from ref 26. ^g Ab initio calculations with a 3-21G* basis set. ^h Energies in Hartrees. ⁱ Fermi resonance.

Table 3. Calculated Frequencies^a for BrF₂⁻ at the RHF and MP2 Levels Using Different Standard Basis Sets

HF-3-21G		HF-3-21G*		HF-6-31+G*		HF-6-311G**		MP2-6-31+G*		assgnt
IR	Raman	IR	Raman	IR	Raman	IR	Raman	IR	Raman	
391		480		368		393		434		$\nu_{as}/\nu_3 (\Sigma_u^+)$
	446		489		454		453		441	$\nu_s/\nu_1 (\Sigma_g^+)$
239		250		231		231		215		$\delta/\nu_2 (\Pi_u)$

^a Frequencies in cm⁻¹.

at 470 cm⁻¹ observed by Christe et al. must be the antisymmetric stretching mode and not the symmetric one. This assignment is in good agreement with *D_{∞h}* symmetry for ClF₂⁻ and follows the rule of mutual exclusion.

NMR Spectra. The resonance signal at 54.5 ppm in the ¹³C-NMR spectrum can be assigned to the quaternary ammonium cation. Literature data for similar quaternary ammonium salts¹⁸⁻²⁰ are between 54.3 and 54.6 ppm. No signals in the range between 119 and 125 ppm for CF₃ groups remained. Obviously the reaction, as described in eq 1, was complete. Two resonances were observed in the ¹⁹F-NMR spectrum due to some

disproportionation of BrF₂⁻ into Br⁻ and BrF₄⁻. The signal at -35.9 ppm is assigned to BrF₄⁻, in good agreement with literature values of about -37 ppm.²¹ The second signal with a chemical shift of about -41.6 ppm belongs to the BrF₂⁻ anion.

Acknowledgment. The financial support of this research by the Fonds der Chemischen Industrie is gratefully acknowledged.

IC961283S