4372

Additions and Corrections

1999, Volume 38

Shuang Liu,* D. Scott Edwards, Anthony R. Harris, Stuart J. Heminway, and John A. Barrett: Technetium Complexes of a Hydrazinonicotinamide-Conjugated Cyclic Peptide and 2-Hydrazinopyridine: Synthesis and Characterization.

Pages 1326–1335. The authors sincerely apologize to D. J. Rose and co-workers for the oversight of their *Inorganic Chemistry* paper (*Inorg. Chem.* **1998**, *37*, 2701–2716), which was inadvertently omitted as a reference in this paper.

IC9909328

10.1021/ic9909328 Published on Web 08/31/1999

1999, Volume 38

Hiromasa Kurosaki, Kentarou Hayashi, Yoshinobu Ishikawa, Masafumi Goto,* Kazufumi Inada, Isao Taniguchi, Mitsuhiko Shionoya, and Eiichi Kimura*: New Robust Bleomycin Analogues: Synthesis, Spectroscopy, and Crystal Structures of the Copper(II) Complexes.

Pages 2829 and 2830. Titration curves e for $Zn^{II}-L_3$ and $-L_4$ in Figures 1 and 2, respectively, are lacking. The correct Figures 1 and 2 and their captions are provided herein.




Figure 1. pH titration curves of triprotonated ligands H_3L_3 in the absence and the presence of equimolar Cu^{II} , Cu^{I} , Fe^{II} , or Zn^{II} at 25 °C and I = 0.1 M NaNO₃. Key: (a) 1.0 mM L₃·3HCl; (b) (a) + 1.0 mM CuSO₄·6H₂O; (c) (a) + 1.0 mM Cu(CH₃CN)₄·ClO₄; (d) (a) + 1.0 mM FeSO₄·7H₂O; (e) (a) + 1.0 mM Zn(NO₃)₂·6H₂O.




Figure 2. pH titration curves of triprotonated ligands H_4L_4 in the absence and the presence of equimolar Cu^{II} , Cu^{I} , Fe^{II} , or Zn^{II} at 25 °C and I = 0.1 M NaNO₃. Key: (a) 1.0 mM L₄·3HCl; (b) (a) + 1.0 mM CuSO₄·6H₂O; (c) (a) + 1.0 mM Cu(CH₃CN)₄·ClO₄; (d) (a) + 1.0 mM FeSO₄·7H₂O; (e) (a) + 1.0 mM Zn(NO₃)₂·6H₂O.

IC990898X

10.1021/ic990898x Published on Web 09/03/1999