

Protonation of Coordinated N₂ on Tungsten with H₂ Mediated by Sulfido-Bridged Dinuclear Molybdenum Complexes¹

Yoshiaki Nishibayashi,^{†,‡} Issei Wakiji,[†] Kenji Hirata,[†] Mary Rakowski DuBois,^{*,§} and Masanobu Hidai^{*,†,‡}

Department of Chemistry and Biotechnology, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-8656, Japan, and Department of Chemistry and Biochemistry, University of Colorado at Boulder, Boulder, Colorado 80309-0215

Received June 27, 2000

Introduction

During the course of our continuing study of the reactivities of coordinated N₂ in molybdenum and tungsten complexes of the type [M(N₂)₂(L)₄] (M = Mo, W; L = tertiaryphosphine),² we have recently found that the coordinated N₂ on tungsten can be protonated with acidic ruthenium dihydrogen complexes to form NH₃ under mild conditions.³ Furthermore, the coordinated N₂ is transformed into NH₃ by treatment with hydrosulfido-bridged dinuclear iridium and iron complexes, although organic thiols and H₂S are not effective for the reaction.⁴ This indicates that a similar type of protonation might occur in biological nitrogen fixation, which is mediated by the bridging sulfur ligands in the FeMo cofactor.⁵ An extension of our study has now led to the investigation of the reactions of the N₂ complexes with the sulfido-bridged dimolybdenum complexes [Cp'Mo(μ₂-S₂CH₂)(μ-S)(μ-SR)MoCp']OTf (**2a**, Cp' = η⁵-C₅H₅, R = H; **2b**, Cp' = η⁵-C₅H₅, R = Me; **2c**, Cp' = η⁵-C₅H₄Me, R = H; **2d**, Cp' = η⁵-C₅H₄Me, R = Me; OTf = OSO₂CF₃) under H₂ because DuBois and co-workers previously reported that the C≡N triple bond in nitriles can be cleaved in the presence of **2a** under H₂ and complex **2d** induces the heterolytic cleavage of H₂ by reaction with pyridine under H₂.^{6,7} The results are described here.

[†] The University of Tokyo.

[‡] Present address: Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan.


[§] University of Colorado at Boulder.

¹ Present address: Department of Materials Science and Technology, Faculty of Industrial Science and Technology, Science University of Tokyo, Noda, Chiba 278-8510, Japan.


- (1) Preparation and Properties of Molybdenum and Tungsten Dinitrogen Complexes. 68. Part 67: Seino, H.; Nonokawa, D.; Nakamura, G.; Mizobe, Y.; Hidai, M. *Organometallics* **2000**, *19*, 2002.
- (2) (a) Hidai, M.; Mizobe, Y. *Chem. Rev.* **1995**, *95*, 1115. (b) Hidai, M.; Ishii, Y. *Bull. Chem. Soc. Jpn.* **1996**, *69*, 819. (c) Hidai, M.; Mizobe, Y. *Top. Organomet. Chem.* **1999**, *3*, 227.
- (3) (a) Nishibayashi, Y.; Iwai, S.; Hidai, M. *Science* **1998**, *279*, 540. (b) Nishibayashi, Y.; Takemoto, S.; Iwai, S.; Hidai, M. *Inorg. Chem.* **2000**, *39*, 5946.
- (4) Nishibayashi, Y.; Iwai, S.; Hidai, M. *J. Am. Chem. Soc.* **1998**, *120*, 10559.
- (5) (a) Dance, I. G. *Aust. J. Chem.* **1994**, *47*, 979. (b) Dance, I. G. *Transition Metal Sulfur Chemistry*; Stiefel, E. I., Matsumoto, K., Eds.; ACS Symposium Series 653; American Chemical Society: Washington, DC, 1996. (c) Dance, I. *Chem. Commun.* **1997**, 165. (d) Dance, I. *Chem. Commun.* **1998**, 523. (e) Zhong, S.-J.; Liu, C.-W. *Polyhedron* **1997**, *16*, 653.
- (6) Bernatis, P.; Laurie, J. C. V.; DuBois, M. R. *Organometallics* **1990**, *9*, 1607.
- (7) Laurie, J. C. V.; Duncan, L. D.; Haltiwanger, R. C.; Weberg, R. T.; DuBois, M. R. *J. Am. Chem. Soc.* **1986**, *108*, 6234.

Results and Discussion

Treatment of *trans*-[W(N₂)₂(dppe)₂] (**1a**; dppe = 1,2-bis(diphenylphosphino)ethane) with 2 equiv of [(η⁵-C₅H₄Me)Mo(μ₂-S₂CH₂)(μ-S)(μ-SMe)Mo(η⁵-C₅H₄Me)]OTf (**2d**) in THF at 25 °C for 1 h under 1 atm of H₂ produced the hydrazido(2-) complex *trans*-[W(OTf)(NNH₂)(dppe)₂]OTf (**3a**) and [(η⁵-C₅H₄Me)Mo(μ₂-S₂CH₂)(μ-SMe)(μ-SH)Mo(η⁵-C₅H₄Me)] (**4**) in >99% and 69% NMR yields, respectively (eq 1). Under N₂, no reaction



proceeded, and both **1a** and **2d** were recovered quantitatively. Hydrazido(2-) complex **3a** was previously obtained by protonation of **1a** with 2 equiv of trifluoromethanesulfonic acid (HOTf).⁸ In the reaction of **1a** and **2d**, the heterolytic splitting of H₂ occurs on **2d**, where one H atom is used for the N–H bond formation at the coordinated N₂ on the W atom and the other H atom remains at the bridging-sulfido ligand. The formal oxidation state of the starting and product molybdenum complexes is +IV and +III, respectively, and the two electrons required for the reduction come from the H₂. Employment of **2b** in place of **2d** also gives rise to the formation of **3a** in high yield. On the other hand, when **1a** was treated with **2a** under H₂, **3a** and [(η⁵-C₅H₅)Mo(μ₂-S₂CH₂)(μ-S)₂Mo(η⁵-C₅H₅)] (**5**) were obtained in >99% and 84% NMR yields, respectively (eq 2). The neutral molybdenum complex **5** is presumed to



formed from dehydrogenation of the initially formed complex [(η⁵-C₅H₅)Mo(μ₂-S₂CH₂)(μ-SH)₂Mo(η⁵-C₅H₅)].⁹

- (8) Field, L. D.; Jones, N. G.; Turner, P. *Organometallics* **1998**, *17*, 2394.
- (9) Casewit, C. J.; Coons, D. E.; Wright, L. L.; Miller, W. K.; DuBois, M. R. *Organometallics* **1986**, *5*, 951.

Table 1. Reaction of Coordinated N₂ on Tungsten and Sulfido-Bridged Dinuclear Molybdenum Complexes under H₂^a

run	W–N ₂ complex	Mo complex	condition temp (°C)/time (h)	yield of NH ₃ (%) ^b		
				free ^c	in extract ^d	total
1	1b	2a	25/24 → 55/24	3	35	38 ^e
2	1b	2a	25/120	4	30	34
3	3b ^f	2a	25/24 → 55/24	5	94	99
4	1b	2b	25/24 → 55/24	5	28	33
5	1b	2c	25/24 → 55/24	3	43	46
6	1b	5	25/24 → 55/24	5	12	17
7	1b	6	25/24 → 55/24	2	15	17
8	1c	2a	25/24 → 55/24	4	48	52 ^e

^a All of the reactions were carried out in THF using W–N₂ complex and 10 equiv of Mo complex under 1 atm of H₂ at 25 °C for 24 h and then at 55 °C for 24 h unless otherwise stated (see text). ^b Yield of NH₃ was based on the W atom. ^c Free NH₃ was observed in the reaction mixture. ^d Yield of NH₄⁺ in the water extract of the reaction mixture. ^e Variation of ±8% between experiments. ^f **3b** was used in place of W–N₂ complex.

Interestingly, treatment of *cis*-[W(N₂)₂(PMe₂Ph)₄] (**1b**) with 10 equiv of **2a** under 1 atm of H₂ in THF at 25 and then 55 °C for 24 h each resulted in the formation of NH₃ in 38% total yield based on the W atom, whereas reactions of **1a** under similar conditions did not produce NH₃. Free NH₃ in 3% yield was detected in the reaction mixture, and further NH₃ in 35% yield was observed as NH₄⁺ in the water extract when the reaction mixture was extracted with an excess of water. The typical results are shown in Table 1. In the absence of H₂ or **2a**, no NH₃ was formed. When the reaction was performed at 25 °C for a much longer time, NH₃ was produced in moderate yield even at that temperature (Table 1, run 2). However, if the reaction was carried out at 55 °C, the formation of NH₃ was not observed. This is probably because **2a** reacts much more quickly with THF than with **1b** at that temperature to give a cationic complex with a 4-hydroxybutanethiolate ligand.¹⁰ In all cases, only a trace amount of NH₂NH₂ was observed. When **1b** was treated with 2 equiv of **2a** at 25 °C for 1 h under H₂, the formation of the hydrazido(2–) complex *trans*-[W(OTf)(NNH₂)(PMe₂Ph)₄]OTf^{3b} (**3b**) in 39% yield was revealed by NMR analysis of the reaction mixture. Furthermore, the reaction of **3b** with 10 equiv of **2a** under H₂ at 25 and then 55 °C for 24 h each produced NH₃ in 99% total yield (Table 1, run 3). These results indicate that the formation of NH₃ proceeds through the protonation of **3b** as an intermediate.

Reactions of **1b** with 10 equiv of other cationic sulfido-bridged dinuclear molybdenum complexes (**2b** and **2c**) under H₂ led to the formation of NH₃ in similar yields (Table 1, runs 4 and 5). However, in the case of neutral dinuclear molybdenum complexes such as **5** and [(η⁵-C₅H₅)Mo(μ-SH)(μ-S)]₂ (**6**), NH₃ was obtained in lower yields (Table 1, runs 6 and 7).

The reaction of *trans*-[W(N₂)₂(PMePh₂)₄] (**1c**) with 10 equiv of **2a** under the same conditions gave NH₃ in 52% yield (Table 1, run 8), which is almost the same as that from the reaction of **1b** with **2a** (vide supra). This is in sharp contrast to our previous observation that treatment of **1b** with 10 equiv of *trans*-[RuCl(η²-H₂)(dppe)₂]OTf at 55 °C under H₂ gives NH₃ in moderate yield while the similar reaction of **1c** affords NH₃ in very low yield.^{3b}

In summary, we have found that the reactions of tungsten N₂ complexes and sulfido-bridged dinuclear molybdenum complexes under H₂ produce NH₃ under mild conditions. The

reaction mechanism is not yet clear; however, it is noteworthy that sulfido-bridged dinuclear molybdenum complexes are able to induce heterolytic cleavage of H₂, leading to the N–H bond formation of coordinated N₂. Thus, heterolytic H₂ activation at a metal–sulfur site, i.e., M–S + H₂ → M(H[–])–SH⁺, has been proposed to realize catalytic nitrogen fixation.¹¹

Experimental Section

General Procedure. Preparation of complexes was performed under 1 atm of N₂ or Ar dried by passage through silica gel and P₂O₅. Reactions of N₂ complexes with sulfido-bridged dimolybdenum complexes were carried out under 1 atm of H₂ dried by passage through silica gel and P₂O₅. THF was freshly distilled over sodium benzophenone ketyl just before use. Unless otherwise noted, all manipulations were done by use of Schlenk techniques. NMR spectra were run on a JEOL JNM-LA-400 or a JEOL JNM-EX-270 spectrometer. IR spectra were recorded on a Shimadzu FTIR-8100M spectrometer. Absorption spectra were measured on a Shimadzu UV-2400PC. Tungsten N₂ complexes^{12–14} (**1**) and sulfido-bridged dimolybdenum complexes^{15,16} (**2**, **5**, **6**) were prepared according to literature procedures.

Reaction of *trans*-[W(N₂)₂(dppe)₂] (1a**) with Sulfido-Bridged Dinuclear Molybdenum Complexes (**2**) under H₂.** A typical experimental procedure for the reaction of **1a** with 2 equiv of **2d** is as follows. In a 20 mL flask was placed **2d** (28.6 mg, 0.0436 mmol) under 1 atm of N₂. Dry THF (15 mL) was added, and then the solution was magnetically stirred at 25 °C for 5 min. After the N₂ atmosphere was replaced by 1 atm of H₂, **1a** (22.5 mg, 0.0217 mmol) was added portionwise. The reaction mixture was stirred at 25 °C for 1 h under 1 atm of H₂. The solvent was then removed under vacuum, and the residue was dissolved in THF-*d*₈ to measure the ¹H NMR spectrum. Ferrocene (10.1 mg, 0.0554 mmol) was added into the THF-*d*₈ solution as an internal reference. The NMR yield of the produced dimolybdenum complex (**4**) was determined by integration of the ¹H resonances against the ferrocene standard. However, the NMR yield of the produced hydrazido(2–) complex (**3a**) could not be determined in this solvent because of overlap of the ¹H resonances with those of impurities. Thus, the solvent was again evaporated under vacuum, and the residue was dissolved in CDCl₃ to determine the NMR yield of **3a** by integration of the ¹H resonances against the ferrocene standard. ⁴:¹⁷ ¹H NMR (THF-*d*₈) δ –1.35 (br s, 1H, SH), 1.52 (s, 3H, SMe), 2.00 (s, 6H, C₅H₄Me), 5.43 (br, 8H, C₅H₄Me), 5.79 (s, 2H, S₂CH₂); 69% NMR yield. **3a**:⁸ ¹H NMR (CDCl₃) δ 2.87 (br s, 4H, CH₂), 3.00 (br s, 4H, CH₂), 5.01 (br s, 2H, NNH₂), 7.21–7.45 (m, 40H, Ph); >99% NMR yield; ³¹P{¹H} NMR (CDCl₃) δ 37.8 (s with ¹⁸³W satellites, *J*_{PW} = 320 Hz).

Formation of NH₃ in the Reactions of Dinitrogen Tungsten Complexes (1b**, **1c**) with Sulfido-Bridged Dinuclear Molybdenum Complexes (**2**) under H₂.** A typical procedure for the reaction of **1b** with 10 equiv of **2a** under 1 atm of H₂ is as follows. In a 500 mL flask was placed **2a** (307 mg, 0.500 mmol) under 1 atm of N₂. Dry THF (20 mL) was added, and then the mixture was magnetically stirred at 25 °C for 5 min. After the N₂ atmosphere was replaced by 1 atm of H₂,

(10) Godziela, G.; Tonker, T.; DuBois, M. R. *Organometallics* **1989**, *8*, 2220.

- (11) Sellmann, D.; Sutter, J. *Acc. Chem. Res.* **1997**, *30*, 460 and references therein.
 (12) (a) Chatt, J.; Pearman, A. J.; Richards, R. L. *Nature (London)* **1975**, *253*, 39. (b) Chatt, J.; Pearman, A. J.; Richards, R. L. *J. Chem. Soc., Dalton Trans.* **1977**, 1852. (c) Anderson, S. N.; Fakley, M. E.; Richards, R. L.; Chatt, J. *J. Chem. Soc., Dalton Trans.* **1981**, 1973.
 (13) Hussain, W.; Leigh, G. J.; Ali, H. M.; Pickett, C. J.; Rankin, D. A. *J. Chem. Soc., Dalton Trans.* **1984**, 1703.
 (14) Chatt, J.; Pearman, A. J.; Richards, R. L. *J. Chem. Soc., Dalton Trans.* **1977**, 2139.
 (15) (a) Casewit, C. J.; Haltiwanger, R. C.; Noordik, J.; DuBois, M. R. *Organometallics* **1985**, *4*, 119. (b) Birnbaum, J.; Godziela, G.; Maciejewski, M.; Tonker, T. L.; Haltiwanger, R. C.; DuBois, M. R. *Organometallics* **1990**, *9*, 394. (c) Birnbaum, J.; Haltiwanger, R. C.; Bernatis, P.; Teachout, C.; Parker, K.; DuBois, M. R. *Organometallics* **1991**, *10*, 1779.
 (16) DuBois, M. R.; VanDerveer, M. C.; DuBois, D. L.; Haltiwanger, R. C.; Miller, W. K. *J. Am. Chem. Soc.* **1980**, *102*, 7456.
 (17) Casewit, C. J.; DuBois, M. R. *J. Am. Chem. Soc.* **1986**, *108*, 5482.

1b (39.4 mg, 0.0497 mmol) was added portionwise. The reaction mixture was stirred at 25 and then 55 °C for 24 h each under 1 atm of H₂. The reaction mixture was evaporated under reduced pressure, and the distillate was trapped in dilute H₂SO₄ solution (1 N; 10 mL). The residue was extracted with H₂O (150 mL), and the aqueous extract was treated with activated charcoal and filtered through Celite. NH₃

and NH₂NH₂ present in the H₂SO₄ and the aqueous solution were quantitatively analyzed by using indophenol and *p*-(dimethylamino)-benzaldehyde reagents, respectively.^{3b,18,19}

-
- (18) Takahashi, T.; Mizobe, Y.; Sato, M.; Uchida, Y.; Hidai, M. *J. Am. Chem. Soc.* **1980**, *102*, 7461.
(19) Weatherburn, M. W. *Anal. Chem.* **1967**, *39*, 971.

Acknowledgment. This work was supported by a Grant-in-Aid for Specially Promoted Research (09102004) from the Ministry of Education, Science, Sports, and Culture of Japan.

IC000716V