

Improved Synthetic Route to *n*-B₁₈H₂₂

Yuqi Li and Larry G. Sneddon*



Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323

Received October 4, 2005


Simple iodine oxidation of the B₉H₁₂[−] anion in toluene at room temperature reliably gives excellent yields (~80%) of *n*-B₁₈H₂₂ (*anti*-B₁₈H₂₂) and thus provides a convenient, large-scale, safe route to this important polyborane cluster.

Octadecaborane has been isolated in two isomeric forms, *n*-B₁₈H₂₂ (also known as *anti*-B₁₈H₂₂) and *i*-B₁₈H₂₂ (*sym*-B₁₈H₂₂). Crystallographic determinations¹ have shown that both compounds have fused-cage structures in which two 10-vertex frameworks share a common edge. The *n*-B₁₈H₂₂ isomer has a structure (Figure 1) containing a center of symmetry, while *i*-B₁₈H₂₂ has the two fragments related by C₂ symmetry. Their high boron content and air stability make these compounds particularly attractive for many potential medicinal^{2,3} and materials^{2,4} applications.


The *n*-B₁₈H₂₂ isomer was originally prepared by degradation of the B₂₀H₁₈^{2−} anion,⁵ but better yield syntheses have been developed starting with 4-*L*-*arachno*-B₉H₁₃ (L = Lewis base) adducts. As shown in eq 1, pyrolysis of 4-Me₂S-*arachno*-B₉H₁₃ either in vacuo or in xylene solution produced *n*-B₁₈H₂₂ in ~20–28% yields along with a mixture of other polyboranes,⁶ while pyrolysis of in situ generated 4-Bu₂O-*arachno*-B₉H₁₃ (eq 2) was reported to give a 34% yield.⁷



More recently, we have found⁸ (eq 3) that when the 4-Me₂S-*arachno*-B₉H₁₃ pyrolysis is carried out in the presence of the inert ionic liquid solvent 1-butyl-3-methylimidazolium chloride (BmimCl) under biphasic conditions, the yield of *n*-B₁₈H₂₂ increased to ~50%.



The best previously reported route⁹ to *n*-B₁₈H₂₂ has been via oxidation of the B₉H₁₂[−] anion with mercuric ion, which gave a 68% yield according to eq 4.



Although this method has proven useful for laboratory-scale syntheses of *n*-B₁₈H₂₂, the cost, safety concerns, and waste disposal problems associated with the use of the HgBr₂ oxidant limit the usefulness of this route for the large-scale production of *n*-B₁₈H₂₂.

The oxidation of polyborane anions with iodine has been previously employed in a number of reactions to effect *arachno* to *nido* or *nido* to *closo* polyhedral conversions. Examples include the synthesis of *nido*-SB₉H₁₁ from *arachno*-SB₉H₁₂^{−10} and the synthesis of [*closo*-1-CB₈H₉[−]] from *nido*-1-CB₈H₁₂ (in the presence of Et₃N).¹¹ In other cases, such as in the formation of B₃H₈[−] from BH₄[−],¹² iodine oxidation

* To whom correspondence should be addressed. E-mail: lsneddon@sas.upenn.edu.

- (1) (a) Simpson, P. G.; Lipscomb, W. N. *Proc. Natl. Acad. Sci.* **1962**, *48*, 1490–1491. (b) Simpson, P. G.; Lipscomb, W. N. *J. Chem. Phys.* **1963**, *39*, 26–34. (c) Simpson, P. G.; Foltling, K.; Dobrott, R. D.; Lipscomb, W. N. *J. Chem. Phys.* **1963**, *39*, 2339–2348.
- (2) Plešek, J. *Chem. Rev.* **1992**, *92*, 269–278.
- (3) (a) Hawthorne, M. F. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 950–984. (b) Soloway, A. H.; Tjarks, W.; Barnum, B. A.; Rong, F.-G.; Barth, R. F.; Codogni, I. M.; Wilson, J. G. *Chem. Rev.* **1998**, *98*, 1515–1562.
- (4) (a) Lu, X.; Shao, L.; Wang, X.; Chen, Q.; Liu, J.; Chu, W.-K.; Bennett, J.; Larson, L.; Ling, P. *Proc. Electrochem. Soc.* **2001**, 2001–2009, 337–344. (b) Kawasaki, Y.; Kuroi, T.; Yamashita, T.; Horita, K.; Hayashi, T.; Ishibashi, M.; Togawa, M.; Ohno, Y.; Yoneda, M.; Horsky, T.; Jacobson, D.; Krull, W. *Nucl. Instrum. Methods Phys. Res., Sect. B* **2005**, *237*, 25–29.
- (5) Pitochelli, A. R.; Hawthorne, M. F. *J. Am. Chem. Soc.* **1962**, *84*, 3218.
- (6) (a) Plešek, J.; Hermanek, S.; Stibr, B.; Hanousek, F. *Collect. Czech. Chem. Commun.* **1967**, *32*, 1095–1103. (b) Plešek, J.; Hermanek, S.; Hanousek, F. *Collect. Czech. Chem. Commun.* **1968**, *33*, 699–705.
- (7) Dobson, J.; Keller, P. C.; Schaeffer, R. *Inorg. Chem.* **1968**, *7*, 399–402.

- (8) Li, Y.; Carroll, P. J.; Sneddon, L. G. *Abstracts of Papers of the 230th National Meeting of the American Chemical Society*, Washington, DC, Aug 2005; American Chemical Society: Washington, DC, 2005; INOR-063.
- (9) Gaines, D. F.; Nelson, C. K.; Steehler, G. A. *J. Am. Chem. Soc.* **1984**, *106*, 7266–7267.
- (10) Pretzer, W. R.; Rudolph, R. W. *J. Am. Chem. Soc.* **1976**, *98*, 1441–1447.
- (11) Brellachs, B.; Backovsky, J.; Stibr, B.; Jelinek, T.; Holub, J.; Bakardijev, M.; Hnyk, D.; Hofmann, M.; Cisarova, I.; Wrackmeyer, B. *Eur. J. Inorg. Chem.* **2004**, 3605–3611.





Figure 1. Structure of $n\text{-B}_{18}\text{H}_{22}$.

has been reported to result in oxidative dehydrocondensation to produce fused polyborane fragments. We report here that simple iodine oxidation of the $\text{B}_9\text{H}_{12}^-$ anion in toluene at room temperature (eq 5) likewise results in an intermolecular dehydrocondensation reaction that reliably gives excellent yields ($\sim 80\%$) of $n\text{-B}_{18}\text{H}_{22}$. This reaction thus provides a convenient, large-scale, safe route to this important polyborane cluster.



In a typical reaction, a 250-mL two-necked, round-bottomed flask equipped with a stir bar was charged, under an argon atmosphere, with 10.11 g (55.0 mmol) of $[\text{Me}_4\text{N}^+][\text{B}_9\text{H}_{12}^-]^{13}$ and ~ 60 mL of dry toluene. After the mixture was sonicated to form a slurry, 7.39 g (29.1 mmol) of I_2 and an additional ~ 85 mL of toluene were added. This mixture was then vigorously stirred at room temperature until (~ 40 min) the iodine color disappeared and a precipitate had formed. The precipitate was removed from the light-yellow solution by filtration. Rotovaporization of the filtrate solvent gave a yellow solid, which $^{11}\text{B}\{^1\text{H}\}$ NMR analysis (Figure 2a) showed to be predominantly $n\text{-B}_{18}\text{H}_{22}$ along with minor amounts of $\text{B}_{10}\text{H}_{14}$ and another unidentified product. The impurities were then easily removed by pumping the material overnight at room temperature on the high vacuum line to leave behind 4.87 g (22.5 mmol, 81.6% yield) of $n\text{-B}_{18}\text{H}_{22}$ as a light-yellow solid. Only trace impurities in the final product could be detected by thin-layer chromatography, $^{11}\text{B}\{^1\text{H}\}$ NMR (Figure 2b), $^1\text{H}\{^{11}\text{B}\}$ NMR (Figure 3), or mass spectrometry. The spectroscopic data and observed melting point ($178\text{--}181$ °C) of this powdery solid agree with the literature values.¹⁴ If necessary, further purification can




Figure 2. (a) $^{11}\text{B}\{^1\text{H}\}$ NMR spectrum of the crude reaction product: (*) $n\text{-B}_{18}\text{H}_{22}$; (●) $\text{B}_{10}\text{H}_{14}$; (#) unidentified product. (b) $^{11}\text{B}\{^1\text{H}\}$ NMR spectrum of the final product obtained after pumping in vacuo.




Figure 3. $^1\text{H}\{^{11}\text{B}\}$ NMR spectrum of the final product obtained after pumping in vacuo.

be readily achieved by sublimation in vacuo at 115 °C onto a -78 °C coldfinger.

Because of its combination of mild conditions, high yields, simple workup, innocuous side products, and the use of an economical and safe oxidation reagent (iodine), the method described above has clear advantages over the other reported procedures. This method will now allow the production of $n\text{-B}_{18}\text{H}_{22}$ on much larger scales than previously possible. These results also suggest that analogous iodine oxidations of other polyborane anions may provide simple routes to other fused-cage polyboranes.

Acknowledgment. We thank the National Science Foundation for support of this research.

IC051712Z

- (12) (a) Nainan, K. C.; Ryschkewitsch, G. E. *Inorg. Nucl. Chem. Lett.* **1974**, *6*, 765. (b) Ryschkewitsch, G. E.; Nainan, K. C. *Inorg. Synth.* **1970**, *15*, 113–114.
- (13) Graybill, B. M.; Ruff, J. K.; Hawthorne, M. F. *J. Am. Chem. Soc.* **1961**, *83*, 2669–2670.
- (14) (a) Olsen, F. P.; Vasavada, R. C.; Hawthorne, M. F. *J. Am. Chem. Soc.* **1968**, *90*, 3946–3951. (b) Fontaine, X. L. R.; Greenwood, N. N.; Kennedy, J. D.; Mackinnon, P. *J. Chem. Soc., Dalton Trans.* **1988**, 1785–1793.