

Gas-Phase Generation and Electronic Structure Investigation of Chlorosulfanyl Thiocyanate, CISSCN

Li Yao,^{†,‡} Maofa Ge,^{*,†} Weigang Wang,^{†,‡} Xiaoqing Zeng,^{†,‡} Zheng Sun,[†] and Dianxun Wang[†]

Beijing National Laboratory for Molecular Sciences (BNLMS), Institute of Chemistry, Chinese Academy of Sciences, Beijing, 100080, China, and Graduate University of Chinese Academy of Sciences, Beijing, 100039, China

Received March 27, 2006

The chlorosulfanyl thiocyanate molecule, CISSCN, was generated in the gas phase through heterogeneous reaction of SCl_2 on the surface of finely powdered AgSCN for the first time. The reaction products were detected and characterized in situ by ultraviolet photoelectron and photoionization mass spectrometry. The molecular geometry and electronic structures of CISSCN were investigated by a combination of PES experiment and theoretical calculations with the density functional theory and ab initio methods. It was found that the outermost electrons of CISSCN reside in the Cl–S antibonding π orbital, predominantly localized on the sulfur atom, and the experimental first vertical ionization potential of CISSCN is 10.20 eV. The dominant fragment SSCN^+ in the mass spectrum indicates that the CISSCN cation prefers the dissociation of the Cl–S bond.

1. Introduction

The sulfur–sulfur bond is a structural feature of widespread occurrence in both organic and inorganic sulfur chemistry. Because of the frequency of disulfide bridges in the structure of many proteins, enzymes, and antibiotics, it plays an important role in stabilizing the structure and in determining the biological activity of the molecules.^{1–3} Recently, considerable attention has been paid to the structure and conformational preferences of simple disulfides. Geometric gas-phase structures of noncyclic disulfides XSSX are characterized by a gauche conformation around the S–S bond. Structural studies of nonsymmetrically substituted disulfides of the type XSSY are less common. Recently, experimental studies on the structure of FC(O)SSCF_3 , FC(O)SSCH_3 , and FC(O)SSC(O)CF_3 have been performed in the gas phase.⁴

Thiocyanates with a SCN group attached directly to sulfur, XSSCN , which possess a sulfur–sulfur bond, have long been known. The first unstable thiocyanate containing a single

S–S bond, NCSSCN , was synthesized by Söderbäck in 1918.⁵ Various polysulfides NCS_nCN ($n < 9$) have been studied by Raman spectroscopy.⁶ Thiacyl thiocyanide NS-SCN was generated by an on-line process using NSCl passed over heated AgSCN and studied by gas-phase FTIR spectroscopy.⁷

We have a long-standing interest in the generation, spectroscopy, and electronic structure of small stable and unstable species. In this work, we present the generation of chlorosulfanyl thiocyanate CISSCN and its subsequent detection and characterization by ultraviolet photoelectron and photoionization mass spectrometry with the help of density functional theory (DFT) and ab initio calculations. The joint spectroscopic and theoretical study provided useful information about the molecular structures, stability, ionization, reaction, and dissociation properties of the interesting unstable CISSCN molecule.

2. Experimental Section

2.1. Generation. Silver thiocyanate was found to be an ideal precursor for the preparation of thiocyanates.^{8,9} In this work,

- (4) (a) Hermann, A.; Ulic, S. E.; Della Védova, C. O.; Mack, H.-G.; Oberhammer, H. *J. Fluorine Chem.* **2001**, *112*, 297–305. (b) Fantoni, A. C.; Della Védova, C. O. *J. Mol. Spectrosc.* **1992**, *154*, 240–245. (c) Erben, M. F.; Della Védova, C. O.; Willner, H.; Trautner, F.; Oberhammer, H.; Boese, R. *Inorg. Chem.* **2005**, *44*, 7070–7077.
- (5) Söderbäck, E. *Liebigs Ann. Chem.* **1919**, *419*, 217–223.
- (6) Fehér, F.; Weber, H. *Chem. Ber.* **1958**, *91*, 642–650.
- (7) Ajji, Z.; Naima, D. Y.; Odeh, M. N.; Allaf, A. W. *Spectrochim. Acta, Part A* **1999**, *55*, 1753–1756.

* To whom correspondence should be addressed. Phone: 86-10-62554518. Fax: 86-10-62559373. E-mail: gemaofa@iccas.ac.cn.

[†] Beijing National Laboratory for Molecular Sciences, Chinese Academy of Sciences.

[‡] Graduate University of Chinese Academy of Sciences.

(1) Wedemeyer, W. J.; Welker, E.; Narayan, M.; Scheraga, H. A. *Biochemistry* **2000**, *39*, 4207–4216.

(2) Kadokura, H.; Katzen, F.; Beckwith, J. *Annu. Rev. Biochem.* **2003**, *72*, 111–135.

(3) Rietsch, A.; Beckwith, J. *Annu. Rev. Genet.* **1998**, *32*, 163–184.

CISSCN was generated by passing SCl_2 vapor over finely powdered AgSCN at 40 °C, and in situ photoelectron and mass spectra were recorded. The precursor SCl_2 was prepared according to a previous report,¹⁰ and its PE spectrum is identical to that recorded previously.¹¹ AgSCN was purchased from the ACROS Company, and its purity was better than 99%. Before reaction, silver thiocyanate was dried in a vacuum (1×10^{-4} Torr) for 2 h at 60 °C.

2.2. Instrumentation. The experimental apparatus used in this work has been described previously.^{12,13} Briefly, the photoelectron and photoionization mass spectrometer consists of two parts; one part is the double-chamber UPS-II machine,¹⁴ the other is a time-of-flight mass spectrometer. The photoelectron (PE) spectrum was recorded on the double-chamber UPS-II machine¹⁴ which was built specifically to detect transient species at a resolution of about 30 meV as indicated by the $\text{Ar}^+(\text{P}_{3/2})$ photoelectron band. Experimental vertical ionization energies are calibrated by the simultaneous addition of a small amount of argon and methyl iodide to the sample. Mass analysis of ions is achieved with the time-of-flight mass analyzer mounted directly to the photoionization point. The relatively soft ionization is provided by single-wavelength HeI radiation. The PE and PIMS spectra can be recorded within seconds of each other under identical conditions.


2.3. Computation. Electronic structure calculations were carried out using the Gaussian series of programs.¹⁵ The geometries of the CISSCN molecule were optimized with B3LYP and MP2 methods at the 6-311+G* and 6-311+G(3df) basis set levels. To assign the PE spectrum of CISSCN molecule, we applied the outer-valence Green's function (OVGF) calculations with a 6-311+G* basis set, which include sophisticated correlation effects of the self-energy, to the molecules to give accurate results of the vertical ionization energies.¹⁶ The adiabatic ionization energy was calculated by taking the difference between the energy of the neutral molecule and the ion at the CBS-QB3 level of theory.¹⁷ Three-dimensional MO plots were obtained with the GaussView program. Each orbital displayed with the 0.08 isodensity value was oriented in a way that displayed for the best view.

Table 1. Total Molecular Energy (au) and Geometric Constants (Å, deg) for the Equilibrium Structure of CISSCN at Different Theoretical Levels

	B3LYP/ 6-311+G*	B3LYP/ 6-311+G(3df)	MP2/ 6-311+G*	MP2/ 6-311+G(3df)
<i>E</i>	-1349.498367	-1349.536166	-1347.560489	-1347.777085
Cl-S	2.102	2.062	2.064	2.030
S-S	2.080	2.039	2.056	2.018
S-C	1.709	1.702	1.701	1.697
C-N	1.158	1.155	1.181	1.176
Cl-S-S	106.2	106.4	104.7	104.9
S-S-C	103.1	103.9	101.0	102.1
S-C-N	175.7	176.5	175.2	177.2
Cl-S-S-C	87.37	86.92	85.05	84.60
S-S-C-N	179.7	178.8	176.3	172.7

3. Results and Discussion

3.1. Molecular Structure. Geometry optimizations were performed for CISSCN using the DFT and ab initio methods.



The structural parameters are depicted in Table 1 (for further geometric constraints, see the Supporting Information). It is well-known that the B3LYP method and the MP2 approximation with small or intermediate basis sets predict S-S, S-Cl, S-F, S-O, etc., bonds longer than they actually are. As demonstrated in test calculations for S_2Cl_2 , the values derived with the MP2 method using a large basis set are expected to be close to the experimental values. Comparing the predicted Cl-S bond with the experimentally determined bond length of some similar molecules, we can see that the predicted Cl-S bond length of CISSCN is close to that of the SCl_2 ¹⁸ (2.015 ± 0.002 Å) and S_2Cl_2 ¹⁹ (2.057 ± 0.002 Å) molecules. In several molecules containing triatomic pseudohalide groups, e.g., SCN, NCO, NNN, etc., it has been established that the atoms are not collinear.^{20,21} When combining with CIS, the SCN angle shows about 5° deviation from linearity, which is similar to those of XCSCN ($X = \text{CN},^{22}$ $\text{SCN},^8$ Cl^{23}). The calculated S-C and C-N bond lengths are also close to the previous reported values. Particularly important parameters are the S-S bond length and Cl-S-S-C dihedral angle as the joint between the CIS and SCN moieties. Comparison of the S-S bond length in S_2X_2 ($X = \text{Cl},^{24}$ $\text{H},^{25}$ CSN_3^{26}) shows that the S-S bond in CISSCN is a typical $\text{S}^{\text{II}}-\text{S}^{\text{II}}$ bond ($2.04-2.06$ Å). The dihedral angle Cl-S-S-C that determines the molecular symmetry is in

- (8) Frost, D. C.; Kirby, C.; Lau, W. M.; MacDonald, C. B.; McDowell, C. A.; Westwood, N. P. C. *Chem. Phys. Lett.* **1980**, *69*, 1-5.
- (9) Yimin, L.; Zhimin, Q.; Qiao, S.; Jincui, Z.; Haiyang, L.; Dianxun, W. *Inorg. Chem.* **2003**, *42*, 8446-8448.
- (10) Brauer, G., Eds.; *Handbook of Preparative Inorganic Chemistry*, 2nd ed.; Academic Press: New York, 1963; pp 186-187.
- (11) Solouki, B.; Rosmus, P.; Bock, H. *Chem. Phys. Lett.* **1974**, *26*, 20-24.
- (12) Yao, L.; Zeng, X. Q.; Ge, M. F.; Wang, W. G.; Sun, Z.; Du, L.; Wang, D. X. *Eur. J. Inorg. Chem.* **2006**, *12*, 2469-2475.
- (13) Yao, L.; Wang, W. G.; Zeng, X. Q.; Ge, M. F.; Du, L.; Sun, Z.; Wang, D. X.; Liang, F.; Qu, H. B.; Li, H. Y. *Anal. Instrum.* **2006**, *1*, 5-9.
- (14) Zhu, X. J.; Ge, M. F.; Wang, J.; Sun, Z.; Wang, D. X.; *Angew. Chem., Int. Ed.* **2000**, *112*, 2016-2019.
- (15) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, J. A.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, Ö.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Andres, J. L.; Gonzalez, C.; Replogle, E. S.; Pople, J. A. *Gaussian 03*, revision B.04; Gaussian, Inc.: Pittsburgh, PA, 2003.
- (16) Ortiz, J. V. *J. Chem. Phys.* **1988**, *89*, 6348-6352.

- (17) Montgomery, J. A.; Frisch, M. J.; Ochterski, J. W.; Petersson, G. A. *J. Chem. Phys.* **1999**, *110*, 2822-2827.
- (18) Davis, R. W.; Gerry, M. C. L. *J. Mol. Spectrosc.* **1977**, *65*, 455-473.
- (19) (a) Beagley, B.; Echersley, G. H.; Brown, D. P.; Tomlinson, D. *Trans. Faraday Soc.* **1969**, *65*, 2300-2307. (b) Marsden, C. J.; Brown, R. D.; Godfrey, P. D. *J. Chem. Soc., Chem. Commun.* **1979**, *9*, 399-401.
- (20) Schulz, A.; Tornieporth-Oetting, I. C.; Klapötke, T. M. *Inorg. Chem.* **1995**, *34*, 4343-4346.
- (21) Klapötke, T. M.; Schulz, A. *Inorg. Chem.* **1997**, *36*, 1929-1933.
- (22) Rosmus, P.; Stafast, H.; Bock, H. *Chem. Phys. Lett.* **1975**, *34*, 275-280.
- (23) Frost, D. C.; MacDonald, C. B.; McDowell, C. A.; Westwood, N. P. C. *J. Am. Chem. Soc.* **1981**, *103*, 4423-4427.




Figure 1. Photoionization mass spectra of (a) CISSCN and (b) the gaseous products of the reaction at 70 °C.

close agreement with the experimental values of 84.8–85.2°, 83.5 ± 1.1°, 90.6 ± 0.5° found for S₂Cl₂,¹⁹ S₂Br₂,²⁷ and S₂H₂, respectively.²⁸

3.2. Photoionization Mass Spectrometry. The reaction process for generating CISSCN was monitored by photoionization mass spectrometry. The ultraviolet photoionization mass spectra of the reaction products of gaseous SCl₂ and AgSCN at different temperatures are shown in Figure 1. Figure 1a depicts the mass spectrum of the reaction at 40 °C, which shows a typical fragmentation distribution of CISSCN cation. The peaks in the spectrum are SCN⁺, SCI⁺, SSCN⁺, S₂Cl⁺, and CISSCN⁺, respectively, with the dominant features being the SSCN⁺ and parent CISSCN⁺ peaks. The appearance of the CISSCN molecular ion confirms its generation and identification. Normally, HeI photoionization produces a fragmentation distribution that is very similar to the distribution observed using electron-impact ionization. This may be rationalized by a similar energy deposition mechanism for the two methods, because the interaction energies (approximately tens of electronvolts) are comparable. As the strongest peak in the HeI PIMS of CISSCN, the SX⁺ (X = SCN) fragment is analogous to that of the electron-impact spectra of other CISX molecules (X = Cl, SCl, and CH₂CH₃),²⁹ which mostly result from the direct dissociation of the Cl–S bond in the CISX⁺ parent ions.

Figure 1b illustrates the photoionization mass spectrum of the gaseous products of the reaction at 70 °C, which reveals features for SCN⁺, SSCN⁺, (SCN)₂⁺, and S(SCN)₂⁺, respectively. Other fragments came from the dissociation of molecular ions (SCN)₂⁺ and S(SCN)₂⁺. Several attempts





Figure 2. HeI photoelectron spectrum of CISSCN.

Table 2. Experimental and Calculated Ionization Energies^a and MO Characters for CISSCN^a

band	exp ^b	calcd	MO	character
1	10.20(V) 9.86(A) ^c	10.19(V) ^d 10.24(V) ^e	31	$\pi_{\text{nb}}(\text{ClS})$
2	10.83	10.84	30	$\pi_{\text{nb}}(\text{SCN})$
3	11.91	11.93	29	$\pi_{\text{nb}}(\text{ClS}) + \pi_{\text{nb}}(\text{SCN})$
4	13.01	12.92	28	$n_{\text{Cl}}, \pi_{\text{b}}(\text{CN})$
		13.07	27	$n_{\text{Cl}}, \pi_{\text{b}}(\text{CN})$
5	13.65	13.66	26	$\pi_{\text{b}}(\text{ClS})$
6	14.07	14.05	25	$\pi_{\text{b}}(\text{ClS}) + \pi_{\text{b}}(\text{SCN})$
7	14.76	14.60	24	σ_{CN}

^a All energies are in eV units; (V) = vertical ionization energy; (A) = adiabatic ionization energy; if no symbol is listed the vertical ionization energy is implied. ^b ±0.01 eV. ^c CBS-QB3. ^d ROVGF from the geometries of MP2/6-311+G*. ^e ROVGF from the geometries of B3LYP/6-311+G*.

were made to maximize the intensities of S(SCN)₂⁺ by raising the reaction temperature, but they all failed until CS₂ appeared, the product of pyrolysis for AgSCN. A possible reaction route is illustrated by the following sequence



In a word, by analyzing the results of ultraviolet photoionization mass spectrometry, we confirmed the generation of CISSCN under our experimental conditions. And the dominant fragment SSCN⁺ found in the mass spectrum indicates that the CISSCN cation prefers the dissociation of Cl–S bond.

3.3. Photoelectron Spectroscopy. The HeI photoelectron spectrum of CISSCN is shown in Figure 2 (for that of the gaseous products of the reaction between SCl₂ and AgSCN, see the Supporting Information). It exhibits seven ionization bands in the low-energy region. We attempted to obtain the PE spectrum of S(SCN)₂, of which the crystal structure has been determined by X-ray methods,³⁰ but found that gaseous S(SCN)₂ is highly unstable and very easily decomposes into (SCN)₂ and S. The PE spectrum of the gaseous products of the reaction at 70 °C, corresponding to the mass spectrum in Figure 1b, shows the typical bands of (SCN)₂.⁸

The experimental and calculated ionization energies as well as the molecular orbital character of CISSCN are listed in Table 2. It is noted that the ROVGF results are in good

(24) Palmer, K. J. *J. Am. Chem. Soc.* **1938**, *60*, 2360–2369.

(25) Stevenson, D. P.; Beach, J. Y. *J. Am. Chem. Soc.* **1938**, *60*, 2872–2876.

(26) Crawford, M.-J.; Klapötke, T. M.; Klüfers, P.; Mayer, P.; White, P. S. *J. Am. Chem. Soc.* **2000**, *122*, 9052–9053.

(27) Hirota, E. *Bull. Chem. Soc. Jpn.* **1958**, *31*, 130–138.

(28) Winniewisser, G. J. *Mol. Spectrosc.* **1972**, *41*, 534–547.

(29) *NIST Chemistry WebBook*; Linstrom, P. J., Mallard, W. G., Eds.; National Institute of Standards and Technology: Gaithersburg, MD, 2001.

(30) Fehér, F.; Linke, K.-H. *Z. Anorg. Allg. Chem.* **1964**, *327*, 151–158.

agreement with the experimental data.^{31,32} Considering the PE spectra of S_2Cl_2 ³³ and $(SCN)_2$,⁸ we can characterize the chlorosulfanyl thiocyanate by its bands that are attributed to the π systems of the CIS and SCN moieties, the nonbonding orbitals of chlorine atom, and the C–N bond. Therefore, the study of these bands may reflect the extent and character of the intramolecular interactions. The MO calculation predicts that the first three transitions for CISSCN belong to an antibonding electron from the CIS and SCN π systems.

The first band, at 10.20 eV with an ionization onset of 9.88 eV, corresponds to the ionization process of CISSCN (X^1A) \rightarrow CISSCN⁺ (X^2A). The experimental result is in good agreement with the calculated vertical IPs. The first band is due to ionization of an electron from the π type of Cl–S antibonding orbital, predominantly localized on the sulfur atom. It is similar to the first ionization band of the analogous molecule SCl_2 , CISCN.^{23,33} Because there is more S 3p involvement in the first orbital of CISSCN, its first IP is lower than that of CISCN (10.52 eV).

The second band is localized at 10.83 eV, which is in agreement with the ROVGF results. It can be described as a π_{SCN} antibonding orbital primarily residing on the sulfur atom. This orbital can also be regarded as being the antisymmetric combination of the sulfur lone-pair and π_{CN} orbital. In most cases, the spectra of pseudohalides are somewhat more complicated. The electronic structure of the SCN group can be explained in two ways. One can assume a single S–C bond and a triple C \equiv N bond. In this case, the appearance of two new bands resulting from the C \equiv N π bonds is expected in the low-energy region of the spectrum. Another interpretation is that the SCN unit is considered to be two perpendicular four-electron, three-center π systems. Consistent with analogous XSCN (X = CN,²³ SCN⁸) molecules, there are ionization events from two π_{CN} orbitals following the first band in CISSCN (Figure 3). Corresponding to the second π_{CN} orbitals, the third PE band at 11.91 eV is the ionization result of the combination of π_{CIS} and π_{SCN} antibonding character.

The 3p nonbonding orbitals of the chlorine atoms are expected to locate in the energy range 11–14 eV. The first ionization potential (IP) of the Cl atom is 12.97 eV,³⁴ and that of chlorine-containing molecules exhibits an IP very close to this value (e.g., $Cl_2 = 11.59$,³⁵ $HCl = 12.75$,³⁵ $ClF = 12.77$ eV³⁶). On this basis, the fourth and fifth bands at 13.01 and 13.65 eV are expected to be the results of electron




Figure 3. Characters of the first eight highest occupied molecular orbitals for CISSCN.

ejection from orbitals containing considerable amounts of chlorine 3p character. The ionization of nonbonding electrons should raise a typical sharp intense band. However, the unusually large width of the fourth band suggests the possibility that this band is actually a composite of two overlapping transitions. And the calculated IPs of orbitals 28 and 27 indicate that both transitions are too close to being resolved. Because of the extensive incorporation of the SCN group, both have primary MO characters that are a mixture of nonbonding orbitals of the chlorine atom and bonding π_{CN} . The remaining chlorine orbitals form MO 26, which is strongly S–Cl bonding. It raises a typical band at 13.65 eV for ionization of electrons from bonding orbitals.

The bands at higher energy 14.07 and 14.76 eV are due to the ionization of bonding orbitals, corresponding to a combination of π_{SCL} and π_{SCN} bonding and a σ_{CN} bonding, respectively (Figure 3).

In a word, the outer molecular orbital structure of CISSCN was studied through photoelectron spectroscopy and quantum chemical calculations for the first time. The outermost electrons reside in the Cl–S antibonding π orbital, predominantly localized on the sulfur atom. The halogen nonbonding electrons are more tightly bound than the antibonding electrons from the CIS and SCN π systems.

4. Conclusions

The novel transient species CISSCN was generated in the gas phase for the first time through in situ heterogeneous reaction of gaseous SCl_2 with $AgSCN$. The gaseous reaction products were detected and characterized by ultraviolet photoelectron and photoionization mass spectrometry. The geometric and electronic structures of CISSCN were investigated by a combination of PES experimental and theoretical studies with the help of DFT and ab initio calculations. The assignment of the PES bands is reasonably supported by previous studies on analogous molecules, as well as the ROVGF and CBS-QB3 calculations. Both the spectroscopic and theoretical investigations have proved the formation of the chlorosulfanyl thiocyanate molecule and suggest that the outermost electrons reside in the Cl–S antibonding π orbital,

(31) Zeng, X. Q.; Ge, M. F.; Sun, Z.; Wang, D. X. *Inorg. Chem.* **2005**, *44*, 9283–9287.

(32) Zeng, X. Q.; Liu, F. Y.; Sun, Q.; Ge, M. F.; Zhang, J. P.; Ai, X. C.; Meng, L. P.; Zheng, S. J.; Wang, D. X. *Inorg. Chem.* **2004**, *43*, 4799–4801.

(33) Colton, R. J.; Rabalais, J. W. *J. Electron Spectrosc. Relat. Phenom.* **1974**, *3*, 345–357.

(34) Kimura, K.; Yamazaki, T.; Achiba, Y. *Chem. Phys. Lett.* **1978**, *58*, 104–107.

(35) Kimura, K.; Katsumata, S.; Achiba, Y.; Yamazaki, T.; Iwata, S. *Ionization Energies, Ab initio Assignments, and Valence Electronic Structure for 200 molecules in the Handbook of HeI Photoelectron Spectra of Fundamental Organic Compounds*; Japan Scientific Society Press: Tokyo, 1981.

(36) Dyke, J. M.; Josland, G. D.; Snijders, J. G.; Boerrigter, P. M. *Chem. Phys.* **1984**, *91*, 419–424.

predominantly localized on the sulfur atom. The halogen nonbonding electrons are more tightly bound than the antibonding electrons from the ClS and SCN π systems. The dominant fragment SSCN^+ in the mass spectrum indicates that the ClSSCN cation prefers the dissociation of the Cl–S bond.

Acknowledgment. This project was supported by the Chinese Academy of Sciences (Hundred Talents Fund) and the National Natural Science Foundation of China (Contract 20577052, 20473094, 50372071, 20503035). L.Y., W.W.,

and X.Z. thank the Chinese Academy of Sciences for receipt of scholarships during the period of this work.

Supporting Information Available: HeI photoelectron spectrum of the gaseous products $(\text{SCN})_2$ of the reaction between gaseous SCl_2 and AgSCN at 70 °C; calculated geometric constants for the equilibrium structure of the neutral molecule and cation; orbital energies with the first eight highest occupied molecular orbitals; and energies of the dissociated fragment of ClSSCN^+ . This material is available free of charge via the Internet at <http://pubs.acs.org>.

IC060513S