

Tetragonal to Triclinic—A Phase Change for [Fe(TPP)(NO)]

Nathan J. Silvernail,[†] Marilyn M. Olmstead,^{‡,*} Bruce C. Noll,[†] and W. Robert Scheidt^{*†}*Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, Indiana 46556, and Department of Chemistry, University of California at Davis, Davis, California 95616*

Received August 25, 2008

The temperature dependence of the crystalline phase of (nitrosyl)(tetraphenylporphinato)iron(II), [Fe(TPP)(NO)], has been explored over the temperature range of 33–293 K. The crystalline complex is found in the tetragonal crystal system at higher temperatures and in the triclinic crystal system at lower temperatures. In the tetragonal system, the axial ligand is strongly disordered, with the molecule having crystallographically required $4/m$ symmetry, leading to eight distinct positions of the single nitrosyl oxygen atom. The phase transition to the triclinic crystal system leads to a partial ordering with the molecule now having inversion symmetry and disorder of the axial nitrosyl ligand over only two positions. The increase in ordering allows subtle molecular geometry features to be observed; in particular, an off-axis tilt of the Fe–N_{NO} bond from the heme normal is apparent. The transition of the reversible phase change begins at about 250 K. This transition has been confirmed by both X-ray diffraction studies and a differential scanning calorimetry study.

Introduction

The perception and understanding of nitric oxide has changed from that of merely a toxic gas to that of an essential cellular signaling agent.¹ NO coordination to heme proteins has many physiological consequences. It has been implicated in a number of fundamental processes including smooth muscle relaxation,^{2,3} platelet deaggregation,⁴ neuronal communication,⁵ aspects of myocardial function, and numerous other physiological functions.⁶ NO is synthesized by nitric oxide synthase through a heme-mediated oxidation of L-arginine^{7–9} or from NO₃⁻/NO₂⁻.¹⁰ In many of these bio-

chemical processes, the binding of NO to a heme protein and the labilization or loss of the ligand trans to the NO or another rearrangement is the significant signaling event, resulting in a five-coordinate heme–NO active complex.

Five-coordinate [Fe(TPP)(NO)]¹¹ was originally characterized in this laboratory in 1975.¹² Although this was more than a decade before the known biological importance of NO, this investigation provided the first insight into the coordination chemistry of NO to heme. Some of the structural features were obscured as a result of the $4/m$ symmetry of the molecular center in the $I4/m$ space group. The molecule had an 8-fold disorder of the NO oxygen, an Fe–N_{NO} bond exactly perpendicular to the porphyrin plane, and phenyl rings rotated exactly 90° from the porphyrin plane. Subsequent structural investigations of well-resolved heme–NO complexes utilizing octaethylporphyrin more precisely detailed the geometrical parameters of the heme–NO com-

* Authors to whom correspondence should be addressed. E-mail mmolmstead@ucdavis.edu (M.M.O.), scheidt.1@nd.edu (W.R.S.).

[†] University of Notre Dame.

[‡] University of California at Davis.

(1) Culotta, E.; Koshland, D. E. *Science* **1992**, *258*, 1862.

(2) Rapoport, R. M.; Murad, F. *Circ. Res.* **1983**, *52*, 352.

(3) Ignarro, L. J.; Adams, J. B.; Horwitz, P. M.; Wood, K. S. *J. Biol. Chem.* **1986**, *261*, 4997.

(4) (a) Azuma, H.; Ishikawa, M.; Sekizaki, S. *Br. J. Pharmacol.* **1986**, *88*, 411. (b) Furlong, B.; Henderson, A. H.; Lewis, M. J.; Smith, J. A. *Br. J. Pharmacol.* **1987**, *90*, 687. (c) Radomski, M. W.; Palmer, R. M. J.; Moncada, S. *Br. J. Pharmacol.* **1987**, *92*, 639.

(5) Garthwaite, J. *Trends Neurosci.* **1991**, *14*, 60.

(6) (a) Koshland, D. E., Jr. *Science* **1992**, *258*, 1861. (b) Butler, A. R.; Williams, D. L. H. *Chem. Soc. Rev.* **1993**, *233*. (c) Moncada, S.; Palmer, R. M. J.; Higgs, E. A. *Pharmacol. Rev.* **1991**, *43*, 109.

(7) Marletta, M. A. *J. Biol. Chem.* **1993**, *17*, 12231.

(8) Griffith, O. W.; Stuehr, D. J. *Annu. Rev. Physiol.* **1995**, *57*, 707.

(9) (a) Alderton, W. K.; Cooper, C. E.; Knowles, R. G. *Biochem. J.* **2001**, *357*, 593. (b) Griffith, O. W.; Stuehr, D. J. *Annu. Rev. Physiol.* **1995**, *57*, 707.

(10) (a) Averill, B. A. *Chem. Rev.* **1996**, *96*, 2951. (b) Eich, R. F.; Li, T.; Lemon, D. D.; Doherty, D. H.; Curry, S. R.; Aitken, J. F.; Mathews, A. J.; Johnson, K. A.; Smith, R. D.; Phillips, G. N., Jr.; Olson, J. S. *Biochemistry* **1996**, *35*, 6976.

(11) The following abbreviations are used in this paper. OEP, dianion of octaethylporphyrin; oxoOEC, dianion of oxooctaethylchlorin, (3,3,7,8,12,13,17,18-octaethyl-(3H)-porphin-2-onato(2-)); OETAP, dianion of octaethyltetraphenylporphyrin; TPP, dianion of *meso*-tetraphenylporphyrin; TP-PBr₄, dianion of 2,3,12,13-tetrabromotetraphenylporphyrin; N_p, porphinato nitrogen atom.

(12) Scheidt, W. R.; Frisse, M. E. *J. Am. Chem. Soc.* **1975**, *97*, 17.

plex.^{13,14} Some of the more salient observations were the off-axis tilting of the Fe–N_{NO} vector from the heme normal and an equatorial asymmetry observed in the Fe–N_p distances with a short pair of bonds bracketing the projection of the tilted and bent NO ligand onto the porphyrin core. These observations were further noted, and the electronic basis for the tilt and the equatorial asymmetry was subsequently recognized in several DFT calculations.^{15–18}

In solid-state vibrational studies utilizing the NRVS technique,¹⁹ a substantial temperature-dependent variation in the Fe–N_{NO} stretch was noted.²⁰ A low-temperature crystallographic study of [Fe(TPP)(NO)] showed that there was an apparent phase change. This led to the detailed temperature-dependent study of the phase change and the resulting phenomenon that accompanies the phase change. The resulting order–disorder phase change, which is completely reversible, is reported herein. The phase change also leads to the observation that, at low temperatures, the structure of [Fe(TPP)(NO)] shows the off-axis tilting of the Fe–N_{NO} bond seen in a number of other iron nitrosyl species.

Experimental Section

General Information. All reactions were carried out under anaerobic conditions using standard Schlenk techniques under an argon atmosphere. All solvents were freeze/pump/thaw degassed ($\times 3$) prior to use. Methanol (Acros) and pyridine (Fisher) were used as received. Nitric oxide (Mittler Specialty Gases) was purified by passing it through a trap containing 4 Å molecular sieves immersed in an ethanol/dry ice slurry.²¹ Free base porphyrin [H₂TPP] was prepared according to the method of Adler et al.²² [Fe(TPP)(Cl)] was prepared according to the metalation procedure of Adler et al.²³ IR measurements were taken on a Nicolet Nexus 670 FT-IR spectrometer. Selected single crystals were ground/mixed minimally with KBr. An approximately 50:1 (KBr/[Fe(TPP)(NO)]) mixture was formed into a pellet using a hydraulic press.

Synthesis of [Fe(TPP)(NO)]. [Fe(TPP)(NO)] was prepared using a modification of the previously reported synthesis.¹² A solution of 100 mg of [Fe(TPP)(Cl)], 3 mL of chloroform, 0.2 mL of methanol, and 0.05 mL of pyridine was placed in a Schlenk flask and freeze/pump/thaw degassed ($\times 3$). The solution was then purged with argon for 5 min, followed by bubbling of NO for approximately 5 min. Upon the addition of NO, the solution turned

from brown to brown-orange, and the final solution may contain some undissolved material. [Fe(TPP)(NO)] was then precipitated by the addition of 100 mL of methanol with thorough stirring. The [Fe(TPP)(NO)] precipitate was isolated by filtration onto sintered glass. Crystals of [Fe(TPP)(NO)] were prepared by dissolving 25 mg (0.04 mmol) of the isolated solid in 7 mL of chloroform and carefully layering the solution in 8 mm glass tubes with degassed methanol followed by flame sealing. X-ray quality crystals of [Fe(TPP)(NO)] were isolated after 10 days at room temperature. IR ν_{NO} in KBr: (1700 cm⁻¹). Note that, if KBr is not thoroughly dried, to > 130 °C, an additional ν_{NO} band is present at 1677 cm⁻¹. This band is irreversibly converted on heating of the KBr pellet (to the 1700 peak).

X-Ray Crystallographic Studies. Four distinct single crystals were indexed at eight temperatures in ascending and descending order using a Bruker D8 Apex II system, with graphite-monochromated Mo K α ($\lambda = 0.71073$ Å) radiation from 33 to 293 K (CRYO Industries or 700 Series Oxford Cryostream).²⁴ Crystals at 273, 290, and 293 K could be indexed as tetragonal (*I*-centered), whereas crystals at 33, 90, 100, 130, and 180 K could be indexed in the triclinic crystal system in the space group *P* $\bar{1}$. Cell parameters and crystal domains were further investigated using cell_now.²⁶ All examples of the triclinic phase are composed of four domains related by ~ 90 or ~ 180 ° rotations about the primitive triclinic [1, 0, 0] axis.²⁷

A total of 11 data sets were collected, and the programs SADABS²⁸ and TWINABS²⁹ were applied for absorption correction. All structures were solved using direct methods as implemented in XS³⁰ and refined using XL.³⁰ All atoms were found after successive full-matrix least-squares refinement cycles on F^2 and refined with anisotropic thermal parameters. Hydrogen atoms were placed at calculated geometries and allowed to ride on the position of the parent atom. Hydrogen thermal parameters were set to 1.2 times the equivalent isotropic U of the parent atom.

Crystal structure data collected at 273, 290, and 293 K were refined in the *I4/m* space group, where the molecule has required *4/m* symmetry (see Table 1). The iron and nitrosyl nitrogen atoms are disordered over two positions equally occupied as required by the mirror plane that lies in the porphyrin plane. The nitrosyl oxygen atom is disordered over eight positions, four positions above and four positions below, the porphyrin plane, equally occupied as required by the 4-fold axis along the Fe–N_{NO} vector. The porphyrin phenyl groups are rotated 90° from the porphyrin plane with required mirror symmetry.

Crystal structure data collected at 33, 90, 100, 130, and 180 K were refined in *P* $\bar{1}$, $Z = 1$. However, alternatively they may be described in a nonstandard setting, *I* $\bar{1}$, $Z = 2$,²⁷ to more conveniently compare with the structures described in *I4/m*. The Fe–N–O unit

(13) Ellison, M. K.; Scheidt, W. R. *J. Am. Chem. Soc.* **1997**, *119*, 7404.

(14) Scheidt, W. R.; Duval, H. F.; Neal, T. J.; Ellison, M. K. *J. Am. Chem. Soc.* **2000**, *122*, 4651.

(15) Leu, B. M.; Zgierski, M. Z.; Wyllie, G. R. A.; Scheidt, W. R.; Sturhahn, W.; Alp, E. E.; Durbin, S. M.; Sage, J. T. *J. Am. Chem. Soc.* **2004**, *126*, 4211.

(16) Praneeth, V. K. K.; Näther, C.; Peters, G.; Lehnert, N. *Inorg. Chem.* **2006**, *45*, 2795.

(17) (a) Ghosh, A.; Wondimagegn, T. *J. Am. Chem. Soc.* **2000**, *122*, 8101. (b) Ghosh, A. *Acc. Chem. Res.* **2005**, *38*, 943.

(18) Cheng, L.; Novozhilova, I.; Kim, C.; Kovalevsky, A.; Bagley, K. A.; Coppens, P.; Richter-Addo, G. B. *J. Am. Chem. Soc.* **2000**, *122*, 7142.

(19) Scheidt, W. R.; Durbin, S. M.; Sage, J. T. *J. Inorg. Biochem.* **2005**, *99*, 60.

(20) Silvernail, N. J.; Sage, J. T.; Alp, E. E.; Sturhahn, W.; Zhang, Z.; Scheidt, W. R. Work in progress. There does not appear to be a relationship in these phenomena.

(21) Dodd, R. E.; Robinson, P. L. *Experimental Inorganic Chemistry*; Elsevier: New York, 1957; p 253.

(22) Adler, A. D.; Longo, F. R.; Finarelli, J. D.; Goldmacher, J.; Assour, J.; Korsakoff, L. *J. Org. Chem.* **1967**, *32*, 476.

(23) Adler, A. D.; Longo, F. R.; Kampus, F.; Kim, J. *J. Inorg. Nucl. Chem.* **1970**, *32*, 2443.

(24) The Oxford control unit is calibrated at the factory using the phase change of Rochelle's salt at 109 K.²⁵ The factory calibration was checked in our laboratory with a well-insulated iron-constantan thermocouple and an Omega 199 temperature meter. All reported temperatures are believed accurate to within 2 K.

(25) Tomaszewski, P. E. *Phase Trans.* **1992**, *38*, 127.

(26) Sheldrick, G. M. *cell_now*; Bruker-Nonius AXS: Madison, WI, 2001.

(27) The transformation matrix (*M*) converts *P* $\bar{1}$, $Z = 1$ to *I* $\bar{1}$, $Z = 2$.

$$M = \begin{pmatrix} 0 & 1 & -1 \\ -1 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

The rotations in the triclinic *I*-centered cell are about the *c* axis.

(28) Sheldrick, G. M. *SADABS*; Universität Göttingen: Göttingen, Germany, 2006.

(29) Sheldrick, G. M. *TWINABS*; Universität Göttingen, Göttingen, Germany, 2006.

(30) Sheldrick, G. M. *Acta Crystallogr.* **2008**, *A64*, 112.

Table 1. Summary of Crystallographic Data for [Fe(TPP)(NO)] at Several Temperatures

temp, K; cryst. #	<i>a</i> , Å	<i>b</i> , Å	<i>c</i> , Å	α , deg	β , deg	γ , deg	<i>V</i> / <i>Z</i> , Å ³	unique data	obsd data ^a	final <i>R</i>	final <i>wR</i> ₂
33, 1 ^b	9.745(2)	9.867(2)	10.375(3)	82.66(2)	66.01(2)	70.05(2)	856.7(4)	4221	2476	0.0486	0.0884
90, 1 ^b	9.754(2)	9.943(2)	10.394(3)	82.25(2)	65.78(2)	69.44(2)	855.81(4)	4337	2843	0.0476	0.1084
100, 2 ^b	9.7310(8)	9.9238(8)	10.3902(6)	82.195(1)	65.693(4)	69.384(5)	855.8(1)	6497	4766	0.0449	0.1213
100, 3 ^b	9.734(3)	9.960(3)	10.391(3)	82.100(3)	65.63(2)	69.25(2)	858.0(4)	5092	4085	0.0788	0.1979
100, 4 ^b	9.7201(7)	9.9105(6)	10.3715(6)	82.304(4)	65.802(4)	69.546(4)	853.81(9)	6741	4480	0.0638	0.1543
130, 4 ^b	9.7394(10)	9.9634(10)	10.3998(10)	82.078(4)	65.661(4)	69.225(4)	859.65(15)	4103	3204	0.0537	0.1354
180, 1 ^b	9.765(2)	10.120(2)	10.466(2)	81.35(2)	65.23(2)	68.12(2)	871.4(4)	4628	2579	0.0481	0.0981
273, 4 ^c	13.4152(2)	13.4152(2)	9.7151(3)	90	90	90	874.2(7)	1101	918	0.0391	0.1132
290, 2 ^c	13.4466(3)	13.4466(3)	9.7498(5)	90	90	90	881.43(5)	1155	973	0.0340	0.0996
293, 1 ^c	13.4748(12)	13.4748(12)	9.7686(18)	90	90	90	886.9(2)	1081	842	0.0351	0.0989
293, 2 ^c	13.4502(3)	13.4502(3)	9.7533(4)	90	90	90	882.23(5)	1108	881	0.0372	0.1024
293, 0 ^{c,d}	13.468(4)	13.468(4)	9.755(8)	90	90	90			951	0.044	

^a [*I* > 2 σ (*I*)]. ^b Triclinic, *P* $\bar{1}$. ^c Tetragonal, *I4/m*. ^d See ref 12.

Table 2. Notable Bonding Parameters for [Fe(TPP)(NO)] and Related Compounds


complex, temp., crystal	Fe–XY ^a	X–Y ^a	Fe–X–Y ^b	<Fe–N _p > ^a	Δ Fe ^c	Fe–N, tilt ^b	ν_{X-Y} ^d	ref
[Fe(TPP)(NO)], 33 K, 1	1.739(6)	1.163(5)	144.4(5)	1.999(4)	0.20	6.3		this work
[Fe(TPP)(NO)], 90 K, 1	1.740(5)	1.153(4)	145.6(4)	1.997(13)	0.20	5.8		this work
[Fe(TPP)(NO)], 180 K, 1	1.737(5)	1.131(4)	147.3(4)	2.002(14)	0.21	4.7		this work
[Fe(TPP)(NO)], 293 K, 1	1.720(6)	1.107(11)	149.5(7)	2.0005	0.23	<i>e</i>	1670/1700 ^f	this work
[Fe(TPP)(NO)]	1.722(2)	1.167(3)	144.4(2)	2.004(15)	0.29	<i>e</i>	1670 ^f	12
[Fe(TPPBr ₄)(NO)](A')	1.734(8)	1.119(11)	147.9(8)	2.01(3)	0.37	5.6	1678 ^g	14
[Fe(TPPBr ₄)(NO)](A'')	1.726(9)	1.144(12)	146.9(9)	2.00(2)	0.32	7.1	1678 ^g	14
[Fe(TPPBr ₄)(NO)](B)	1.691(11)	1.145(16)	145(1)	1.95(3)	0.29		1681 ^g	14
[Fe(OEP)(NO)](A)	1.722(2)	1.167(3)	144.4(2)	2.004(15)	0.29	6.5	1666 ^g	13, 14
[Fe(OEP)(NO)](B)	1.7307(7)	1.1677(11)	142.74(8)	2.009(12)	0.27	8.2	1673 ^g	13, 14
[Fe(oxoOEC)(NO)]	1.7320(13)	1.1696(19)	143.11(15)	2.009(9)	0.26	7.1	1690 ^g	14
[Fe(OETAP)(NO)]	1.721(4)	1.155(5)	143.7(4)	1.932(9)	0.31	7.6	1666 ^g	34
[Fe(OEP)(NO)] ⁺	1.6528(13)	1.140(2)	173.19(13)	1.994(5)	0.32	4.6	1838 ^g	35
[Fe(OEP)(NO)] ⁺	1.644(3)	1.112(4)	176.9(3)	1.994(1)	0.29	0.6	1868 ^g	36
[Fe(OEP)(CO)]	1.7140(11)	1.1463(12)	177.20(8)	1.988(2)	0.20	3.8	1944 ^g	37
[Fe(OEP)(CO)]·C ₆ H ₆	1.7077(13)	1.1259(16)	177.20(11)	1.984(3)	0.22	2.4	1948 ^g	37
[Fe(OEP)(CO) ₂]	1.8558(10)	1.1216(13)	173.95(9)	2.0133(7)	0.00	5.9	2021 ^g	37

^a Measured in angstroms. ^b Measured in degrees. ^c Displacement from 24-atom mean plane. ^d Measured in cm⁻¹. ^e Value not determined, obscured by crystallographic symmetry. ^f KBr pellet. ^g Nujol mull.

is disordered over two positions equally occupied as required by the inversion symmetry. The four domains of the triclinic crystals are related by rotations about the [1, 0, 0] axis, which is coincident with the original Fe–N_{NO} bond.

Powder Diffraction Studies. Powder X-ray diffraction (XRD) patterns were obtained on a Bruker Smart Apex I CCD diffractometer with graphite monochromated Cu K α ($\lambda = 1.54178$ Å) radiation from 100 to 300 K (700 Series Oxford Cryostream). Samples were milled in Apiezon M vacuum grease, mounted on a glass fiber, and transferred to the cold gas stream of the diffractometer. Samples were equilibrated for 5 min at each temperature before data were collected. Data were collected with the 2D Apex detector fixed at 100 mm, $2\theta = 12^\circ$, $\phi = 0^\circ$, and ω scanned between 0 and 20° over a 5 min interval. Data were collected with both increasing and decreasing temperature. Data were integrated using the XRD² plugin in the APEX 2 suite of programs³¹ and further analyzed using the program EVA.³²

DSC Measurements. The differential scanning calorimetry (DSC) measurements were performed at 150 to 400 K with a 10 K/min heating rate using a Netzsch STA 404 DSC. Cooling was provided by liquid nitrogen boiloff. Temperature calibration was performed at the melting points of *n*-hexane, cyclohexane, octane, triply deionized water, gallium, naphthalene, indium, tin, and zinc. The scans were carried out in an atmosphere of flowing He (40 mL/min). A standard 40 μ L aluminum crucible with a lid, which could be sealed hermetically with a crucible sealing press, was used for all experiments. Crucibles and lids were heated before experi-

**Figure 1.** Thermal ellipsoid plot of [Fe(TPP)(NO)] at 293 K (50% probability ellipsoids). Hydrogen atoms are omitted for clarity.

ments at about 570 K for 15 min to convert any hydroxides of aluminum that had been formed on the crucible surface to aluminum oxide. All DSC measurements were performed on 8–12 mg samples sealed under atmospheric conditions. Onset temperatures and enthalpies were determined from two individual measurements using a software package supplied by the manufacturer. Reproducibility was confirmed using Setaram DSC 111.

Results

Crystal and molecular structures of [Fe(TPP)(NO)] have been obtained at 33, 90, 100, 130, 180, 273, 290, and 293 K on four different crystalline specimens. A reversible phase change from the tetragonal crystal system to the triclinic crystal system was identified. Crystallographic details for all 11 structure determinations are summarized in Table 1. Thermal ellipsoid plots of [Fe(TPP)(NO)] at 33 and 293 K are illustrated in Figures 1 and 2. At 293 K, [Fe(TPP)(NO)] has eight equally occupied NO orientations consistent with the previously reported structure at 298 K.¹² At 33 K, [Fe(TPP)(NO)] has two equally occupied NO orientations consistent with the triclinic system. A number of diffraction

(31) APEX 2; Bruker-Nonius AXS: Madison, WI, 2004.

(32) EVA; Bruker-Nonius AXS: Madison, WI, 2002.




Figure 2. Thermal ellipsoid plot of [Fe(TPP)(NO)] at 33 K (50% probability ellipsoids). Hydrogen atoms are omitted for clarity.




Figure 3. Diagrams of the tetragonal and triclinic *I*-centered cells showing the effects of the partial order–disorder phase transition. At 33 K, the Fe–NO group retains the disorder with respect to crystallographic inversion, but the disorder of the NO group due to 4-fold symmetry has been removed. In the tetragonal phase, the porphyrin plane is perpendicular to the *c* axis but is tipped by 9.3° from the perpendicular to the *c* axis in the triclinic phase.

experiments confirm the phase change that is also characterized by differential scanning calorimetry. Figure 3 gives representations of the *I*-centered cells at 33 and 293 K.

Discussion

Crystalline [Fe(TPP)(NO)] displays temperature-dependent phases that change from a tetragonal *I*-centered room-temperature phase to a triclinic phase at reduced temperatures. The triclinic phase can be described in terms of an *I*-centered triclinic cell that is a nonstandard crystallographic choice or as a standard primitive triclinic cell. The transition from room temperature to reduced temperature has been followed by differential scanning calorimetry. The DSC scan is illustrated in Figure 4. Two endotherms are observed that are centered at 280.4 K and 249.2 ± 0.2 K. The lower-temperature transition is the one associated with the crystallographic phase change. The basis of the higher temperature transition is unclear; perhaps it is associated with a modest change in the porphyrin ring conformation. However, the




Figure 4. DSC scan for [Fe(TPP)(NO)]. The endotherms are centered at 280.4 and 249.2 ± 0.2 K. The lower-temperature transition corresponds to the observed crystallographic phase change.




Figure 5. X-ray powder diffraction patterns of [Fe(TPP)(NO)] from 100 to 300 K at 20° increments.

structure determinations (at 273 and 293 K) that bracket this endotherm do not show any distinct, different features. We are thus unable to offer a conclusive explanation for this higher-temperature transition.

We have also followed the phase change by examining the diffraction pattern over a range of temperatures. Figure S1 (Supporting Information) displays the predicted powder diffraction patterns at 100 and 290 K (Cu K α radiation) that are based on the cell constants obtained from single crystals at those temperatures. The tetragonal reflection 101 (at 11.25° 2 θ) cleanly splits into four resolved reflections upon the decrease in crystal system symmetry. Accordingly, we focused on that region in a temperature-dependent powder diffraction measurement. Figure 5 displays the actual powder diffraction traces observed every 20° from 300 to 100 K. Figure S2 (Supporting Information) displays the change in intensity at the 11.25° 2 θ position for both ascending and descending temperatures, and Figure 6 shows the intensity change for two triclinic reflections that are Laue equivalents in the tetragonal phase. Although the transition begins sharply




Figure 6. A plot of measured intensities on a single crystal of Fe(TPP)(NO). Above the transition point, the reflections (0 1 1) and (1 0 1) are Laue equivalents in the tetragonal cell, and the quality of the data sets for the two phases appears similar. They become inequivalent at the phase change, which can be ascribed to the effects of thermal motion.

at about 249 K, there does appear to be an asymmetry in the transition as the temperature is decreased, which approaches the final complete change asymptotically. This behavior is also seen in two other regions where peaks in the two phases are sufficiently resolved to allow a measurement. These are the tetragonal reflections 231 ($2\theta = 25.57^\circ$) and 121 ($2\theta = 17.36^\circ$). These are also illustrated in Figure S2 of the Supporting Information. The calculations of the inflection point, also shown in the figure, are all at essentially the same temperature (243–248 K). A single-crystal data collection taken at 242 K could not be indexed solely in terms of either the triclinic cell or the tetragonal cell but rather only in terms of both lattices. There may also be a slight asymmetry in the phase change between ascending and descending temperatures, but the complete reversibility is to be emphasized.

We have also measured the reflection profile of the (3 1 0) reflection on a point detector diffractometer (ω scans). As the temperature is lowered, the peak begins to broaden between 252 and 248 K. The peak begins to split at 245 K and at 235 K is well resolved into two peaks. Little further change occurs below 212 K. These ω scan plots are given in Figure S2 of the Supporting Information. Single-crystal data collected at 180 K or lower are readily indexed in the triclinic cell and that at 273 K or higher in the tetragonal cell, and the quality of the data sets appears to be similar in both crystal systems. Before describing the differences in the crystal structure between the two phases, we first describe the structure of [Fe(TPP)(NO)] in the two phases.

Molecular Structure of [Fe(TPP)(NO)]— $I4/m$ Phase. The molecular structure of the room-temperature phase of [Fe(TPP)(NO)] has been previously reported.¹² The complex's molecular structure agrees well with that previously reported. Complete details of all structure determinations in the tetragonal phase are given in the Supporting Information.

As displayed in Figure 1, the NO ligand is disordered over eight positions. The $4/m$ symmetry at the molecular center necessitates equal occupation of each of the eight positions. One common feature of iron(II) heme nitrosyls is the tilting of the Fe–N_{NO} vector.^{13,14} However, the nitrogen atom of NO is located on the 4-fold axis, making any evaluation of ligand tilt impossible, and the angle is nominally equal to 0° . Past experience suggests that, if the off-axis tilts were similar to those observed previously,¹⁴ the displacement of the NO nitrogen atom is just at the limit of leaving a trace in the thermal ellipsoid of the N atom. Any small changes in the position of nitrogen caused by the eight orientations make it very difficult to resolve a nitrogen position where the Fe–N vector is tilted off the heme normal. The uncertainties in the N–O bond length and Fe–N–O bond angle are further accentuated by the uncertainty in the nitrogen position. The porphyrin ring is required to be precisely planar with the peripheral phenyl group, making a dihedral angle of 90° with the central plane.

Molecular Structure of [Fe(TPP)(NO)]— $P\bar{1}$ Phase. The low-temperature phase of [Fe(TPP)(NO)] was refined in the $P\bar{1}$ setting. The 4-fold symmetry observed in the tetragonal phase is reduced to inversion symmetry in the triclinic phase, and hence, the NO oxygen is disordered only over two positions related by inversion symmetry. An ORTEP diagram of [Fe(TPP)(NO)] at 33 K is given in Figure 2. The ordering allows for a better description of the iron coordination parameters. The elimination of the NO rotational disordering reveals one hallmark feature of Fe(II) nitrosyl porphyrinates.³³ This feature is the off-axis tilting of the Fe–N_{NO} from the heme normal in the direction of the NO oxygen atom. At 33 K, the Fe–N_{NO} vector is tilted 6.3° off the heme normal in the direction of the NO oxygen atom. The Fe–N_{NO} off-axis tilts and several additional bonding parameters for [Fe(Por)(NO)] and related complexes are given in Table 2.^{34–37} The tilt is also usually associated with an asymmetry in the equatorial Fe–N_p bonds. The two Fe–N_p bonds that bracket the NO ligand are shorter than the average, whereas the other two are longer. This asymmetry is clearly a bonding effect. There is only a hint of this asymmetry for triclinic [Fe(TPP)(NO)]. At 33 K, the two short Fe–N_p bonds are 1.995(9) and 1.997(9) Å, whereas the long Fe–N_p bonds are 2.000(9) and 2.004(9) Å. The presence of the inversion center is likely to have led to obscuring of any in-plane asymmetry.

The porphyrin plane is required to be precisely planar in the tetragonal phase. In the triclinic phase, there is no such requirement, but the core remains effectively planar, with the largest core atom displacement from the best 24-atom plane only 0.03 Å (at 33 K). The out-of-plane displacement of iron is 0.20 Å, slightly less than seen in the tetragonal

(33) Wyllie, G. R. A.; Scheidt, W. R. *Chem. Rev.* **2002**, *102*, 1067.

(34) Bohle, D. S.; Debrunner, P.; Fitzgerald, J.; Hansert, B.; Hung, C.-H.; Thompson, A. J. *J. Chem. Soc., Chem. Commun.* **1997**, 91.

(35) Ellison, M. K.; Schulz, C. E.; Scheidt, W. R. *Inorg. Chem.* **2000**, *39*, 5102.

(36) Scheidt, W. R.; Lee, Y. J.; Hatano, K. *J. Am. Chem. Soc.* **1984**, *106*, 3191.

(37) Silvernail, N. J.; Noll, B. C.; Scheidt, W. R. *Inorg. Chem.* **2006**, *45*, 7050.

Table 3. Molecule Center-to-Center Distances and Symmetry Operators for [Fe(TPP)(NO)] in the $\bar{I}\bar{1}$ (33 K) and $I4/m$ (293 K) Phases from Crystal 1

symmetry position	x, y, z	$\bar{I}\bar{1}$, Ct...Ct (Å)	$I4/m$, Ct...Ct (Å)	difference (Å) ^a
x, y, z	1/2, 1/2, 0	0	0	0
$x, y, z + 1$	1/2, 1/2, 1	9.745	9.769	-0.024
$x, y, z - 1$	1/2, 1/2, -1	9.745	9.769	-0.024
$-x, -y, -z$	1, 1, 1/2	9.867	10.707	-0.840
$-x - 1, -y - 1, -z - 1$	0, 0, -1/1	9.867	10.707	-0.840
$-x, -y - 1, -z$	1, 0, 1/2	10.973	10.707	0.266
$-x - 1, -y, -z - 1$	0, 1, -1/2	10.973	10.707	0.266
$-x - 1, -y, -z$	0, 1, 1/2	10.375	10.707	-0.332
$-x, -y - 1, -z - 1$	1, 0, -1/2	10.375	10.707	-0.332
$-x - 1, -y - 1, -z$	0, 0, 1/2	11.256	10.707	0.549
$-x, -y, -z - 1$	1, 1, -1/2	11.256	10.707	0.549
$x + 1, y, z$	3/2, 1/2, 0	13.373	13.475	-0.102
$x - 1, y, z$	-1/2, 1/2, 0	13.373	13.475	-0.102
$x, y + 1, z$	1/2, 3/2, 0	13.353	13.475	-0.122
$x, y - 1, z$	1/2, -1/2, 0	13.353	13.475	-0.122

^a (Ct...Ct in $\bar{I}\bar{1}$) - (Ct...Ct in $I4/m$).

phase. The peripheral phenyl group, required to have exactly orthogonal dihedral angles in the tetragonal phase, have rotated slightly with dihedral angles between the phenyl plane and the core plane of 83.1 and 81.4°.

Phase Change. Why does the phase change occur? The phase change does not appear to lead to a substantial increase in the solid-state packing efficiency. The decrease in cell volume of ~3.5% between 90 and 293 K is similar to those seen in a number of other complexes over a similar temperature change and for which no phase change is observed.³⁸ Rather, the phase change appears to result from the fact that the [Fe(TPP)(NO)] molecule is intrinsically not as symmetric as it appears (and is required) in the tetragonal phase. This is partly, if not entirely, the consequence of the intrinsic off-axis tilt of the Fe-N_{NO} vector that has been observed in all well-ordered (nitrosyl)iron porphyrinates. One explanation of the phase change is that it leads to an order-disorder transition that is sometimes observed upon crystal cooling.³⁹ In this view, the intrinsically lower symmetry of the molecule is favored even though its solid-state structure does not pack with significantly larger efficiency. Nonetheless, the solid-state interaction of the porphyrin molecules displays significant changes, as shown below.

An alternative idea is that the axial ligand distortion plays a fundamental role. As the temperature is decreased, the Fe-NO off-axis tilt becomes increasingly favored. When the distortion occurs, the entire lattice will make small adjustments in the intermolecular packing that leads to the descent to triclinic symmetry and a single Fe-NO orientation above the porphyrin plane. That the NO tilting is driving the phase change is consistent with the observation of four crystal domains, with relative rotations of either 90 or 180° with respect to the major domain. The rotations are along the original Fe-NO axis, consis-

tent with the four possible tilt direction choices. Of course, a similar change must occur on the other side of the porphyrin plane since that side of the porphyrin plane is also populated in the tetragonal phase. The reason for the preservation of an inversion center at the porphyrin ring center is that it is the most symmetric of the possible orientations of the Fe-N-O group and leads to the preservation of the intrinsic NO-induced asymmetry of the Fe interaction with the porphyrin nitrogen atoms (core asymmetry).

The phase change leads to asymmetry in the environment of an individual [Fe(TPP)(NO)] molecule. The *I*-centered triclinic cell constants are $a = 13.373(3)$, $b = 13.353(4)$, $c = 9.745(2)$ Å, $\alpha = 99.310(17)$, $\beta = 93.651(15)$, and $\gamma = 90.64(2)^\circ$ and are clearly similar to those of the tetragonal phase. This has been shown in Figure 3. In the triclinic system, the porphyrin planes are in the xy plane and form parallel layers of porphyrin molecules comparable to the planes parallel to the xy plane in the tetragonal phase. However, in the tetragonal phase, the porphyrin layers are exactly superimposed, but in the triclinic phase, the molecules are no longer directly above each other as a result of the 9.3° tilt of the porphyrin planes from the c axis. Distances between molecular centers reflect this change. The 14 closest intermolecular center...center distances in both crystal systems are given in Table 3. The closest pair of contacts to the target molecule remains the pair related by a translation perpendicular to the molecular array direction. The next eight closest contacts are all equal in the tetragonal system. In the triclinic system, the analogously related intermolecular distances are either shorter or longer and fall along the body diagonals in pairs and are summarized in Table 3. One of the consequences of the resulting asymmetry is that the intermolecular contacts now constrain the Fe-N-O group to a single allowed orientation on each side of the porphyrin plane in the triclinic phase. This also allows the observation of the off-axis tilt of the Fe-NO vector in the triclinic phase that, if present in the tetragonal system, will be obscured by the crystallographically required symmetry.

Summary

Crystalline [Fe(TPP)(NO)] is found to undergo a reversible temperature-dependent phase change from tetragonal to triclinic. In the lower-symmetry crystal system, the intrinsic off-axis tilt of the Fe-NO vector from the porphyrin plane becomes apparent. Two suggestions for the origin of the phase change are given.

Acknowledgment. We thank the National Institutes of Health for support of this research under Grant GM-38401 (W.R.S) and the NSF for X-ray instrumentation support through Grant CHE-0443233. M.M.O. thanks the University of California, Davis, for the He diffraction setup. We acknowledge Olga Trofymuk, Department of Chemistry, Thermochemistry Facility, University of California, Davis, for providing assistance with the DSC results. The thoughtful remarks of the reviewers helped significantly in the final form of this paper.

(38) Silvernail, N. J.; Pavlik, J. W.; Noll, B. C.; Schulz, C. E.; Scheidt, W. R. *Inorg. Chem.* **2008**, *47*, 912.

(39) Kubicki, M *Acta Crystallogr., Sect. B* **2004**, *B60*, 333, and references therein.

Supporting Information Available: Figure S1, simulated powder diffraction data based on single-crystal diffraction experiments; Figure S2, single-reflection intensity/temperature-dependence plots and the first derivative of temperature-dependence plots; Figure S3, reflection profiles; Figures S4–S14, thermal ellipsoid plots; Figure S15, formal diagrams displaying the perpendicular displacement of core atoms from the 24-atom mean planes at all measured

temperatures; Figure S16, stereo diagrams of the two cells; Tables S1–S66, complete crystallographic details, atomic coordinates, bond distances and angles, anisotropic temperature factors, and hydrogen positions for [Fe(TPP)(NO)] at all measured temperatures. Crystallographic data are available as CIF files. This material is available free of charge via the Internet at <http://pubs.acs.org>.

IC801617Q