

Comparative Study of Uranyl(VI) and -(V) Carbonato Complexes in an Aqueous Solution [*Inorg. Chem.* **2007**, *46*, 4212. DOI: 10.1021/ic070051y]. Atsushi Ikeda,* Christoph Hennig, Satoru Tsushima, Koichiro Takao, Yasuhisa Ikeda, Andreas C. Scheinost, and Gert Bernhard

Page 4219. In this paper, we presented uranium L_I- and L_{III}-edges XANES spectra for uranyl(VI) and -(V) tricarbonato complexes in an aqueous solution, together with the energy values of their absorption edges and main oscillation peaks. However, we found that some of the energy values for the L_{III}-edge spectra were incorrect because the wrong procedure was used for the energy correction and definition of $E_{k=0}$ values. The correct energy values and graph are given below. Note that this change has no influence on the conclusion of the paper.




Figure 6. Uranium L_{III}-edge XANES spectra for uranyl(VI) and -(V) tricarbonato complexes.

Table 4. Summary of Uranium L_{III}-Edge XANES Spectra for Uranyl(VI) and -(V) Tricarbonato Complexes

| | absorption edge/eV | P ₁ /eV | P ₂ /eV | P ₃ /eV |
|------------|--------------------|--------------------|--------------------|--------------------|
| uranyl(V) | 17171.2 | 17175.7 | 17184.0 | 17207.7 |
| uranyl(VI) | 17173.3 | 17177.5 | 17193.2 | 17211.1 |

DOI: 10.1021/ic900917v
Published on Web 05/28/2009