

Papers

B10

Spectroscopic Properties of Nd^{3+} in $\text{Cs}_2\text{Na}(\text{La},\text{Nd})\text{Cl}_6$

W. STRĘK, C. SZAFRAŃSKI, J. HANUZA, Z. MAZURAK
and B. JEŻOWSKA-TRZEBIATOWSKA*

*Institute for Low Temperature and Structure Research,
Polish Academy of Sciences, P.O. Box 937, Wrocław 50950,
Poland*

The results of the absorption, infrared, Raman and fluorescence spectra are reported. Owing the analysis of vibrational levels based upon the IR and Raman spectra the assignment of vibronic peaks of fluorescence spectra is proposed. The anomalous concentration dependence of fluorescence decay rates is found. A discussion of the possible mechanism responsible for concentration quenching of Nd^{3+} fluorescence in the $\text{Cs}_2\text{Na}(\text{La},\text{Nd})\text{Cl}_6$ crystal indicates that at low temperature the non-resonant cross relaxation occurs whereas at room temperature both the non-resonant cross relaxation and the anti-Stokes resonant cross relaxation contribute in the Process.

B11

Magnetic Circular Dichroism of Rare Earths and UO_2^+ Complexes

C. GÖRLLER-WALRAND

*Lab. voor anal. en anorg. scheikunde-Celestijnenlaan 200F,
3030 Heverlee, Belgium*

B12

Quasi-relativistic SCF- $X\alpha$ Scattered-wave Calculation of Uranocene, Thorocene and Cerocene

N. RÖSCH

Lehrstuhl für Theoretische Chemie der Technischen, Universität München, Munich, F.R.G.

B13

Evidence for 1D Coupling in the Neutron Spectroscopy Study of Uranium Trihalides

A. MURASIK

Institute of Atomic Energy Swierk, 05-400 Otwock, Poland

Published data of magnetic susceptibilities for UBr_3 and UCl_3 show apparent but rather broad maxima occurring at 15 and 22 K respectively. They were interpreted as due to cooperative magnetic transitions [1]. However, recent neutron scattering experiments performed at low temperatures on polycrystalline samples of UBr_3 and UCl_3 have proved unambiguously that the long range 3D antiferromagnetic ordering occurs at temperatures (UBr_3 : $T_N < 4.2$ K, UCl_3 : $T_N = 6.5$ K) which lie substantially below the values anticipated from magnetic susceptibility measurements [2].

This highly unusual behaviour can be explained by the suggestion that the magnetic exchange interaction is strong within linear chains of uranium ions lying parallel to the c axis of the crystal, and much weaker between uranium ions lying in different chains. The above hypothesis has interesting consequences in the case when intrachain coupling between nearest neighbours is antiferromagnetic, since the dominance of the nearest-neighbour magnetic interaction will result in the ions ordering in antiferromagnetic linear chains along c axis. This by itself does not lead directly to a long-range-ordered structure, and by symmetry argument appears to preclude any simple magnetic ordering.

Consequently, we may expect these compounds to behave as antiferromagnetic linear chains over a considerable range of temperature. The magnetic susceptibility maxima at 15 K and 22 K observed for UBr_3 and UCl_3 may be interpreted as due to exchange interactions within the chains, whereas at sufficiently low temperature, interchain interactions become important. They are expected to be largely, if not entirely dipolar in character, and tend to align the ions which lie in planes perpendicular to the c axis, into what is equivalent to the highly degenerate antiferromagnetic triangular net arrangement [3]. In fact, neutron experiments performed at $T < T_N$ revealed in both compounds a rather complex non collinear magnetic structure. However, the theoretical interpretation of the ordered state is found to be complicated by the unusual symmetry of the lattice.