



ELSEVIER

Inorganica Chimica Acta 228 (1995) 11–61

**Inorganica  
Chimica Acta**

## Cumulative Subject Index of Volumes 215–227

### Acenaphthene complexes

synthesis and X-ray structure of [acenaphthene–chromium(0)-dicarbonyl-tri(2,3,5-trimethylphenyl)phosphite], **219**, 209

### Acetonitrile complexes

synthesis and characterization of uranium triiodide tetrakis(acetonitrile), **218**, 203

### Acetylacetonate complexes

alkylation of 2-vinylpyridine and crystal structure of bis((acetylacetonate)(1-pyridylbutyl)nickel(II)) complex, **227**, 149

### Acetyl complexes

the equilibrium between acetyl- and methylcobalt tetracarbonyl, **227**, 211

kinetics and mechanism of the methanolysis of acetylcobalt tetracarbonyl, **227**, 297

### Acetylide complexes

electron transfer in *trans*-[Pt(PPh<sub>3</sub>)<sub>2</sub>(–C≡C–Fc)<sub>2</sub>] and related compounds, **225**, 35

### Acidity

controlling the acidity of the carboxylic group by a ferrocene based redox switch, **225**, 239

### Acrylonitrile complexes

synthesis, crystal structure and characterization of nickel(0) acrylonitrile complexes with a bidentate N-donor ligand, **227**, 137

### Activation

activation of molecular oxygen by iron nitrosyls in the presence of the bidentate phosphines 1,2-bis(diphenylphosphino)ethane and ethene, **221**, 133

the role of H<sub>2</sub>Ru(CO)<sub>2</sub>(PBU<sub>3</sub>)<sub>2</sub> in the activation of alkynes and alkenes, **227**, 247

### Acylate complexes

the effects of counterion and solvent on the reactivity of nickel acylate complexes, **222**, 165

### Aldol addition reactions

stereoselective aldol addition reactions of Fischer carbene complexes via electronic tuning of the metal center for enolate reactivity, **220**, 215

### Aldolisation reactions

uranium compounds as catalysts for aldolisation reactions, **222**, 91

### Alkali cation complexes

synthesis and electrochemical properties of oxo-bridged manganese dimers incorporating alkali and alkaline earth cations, **225**, 191

### Alkali metal complexes

structures of layered metal sulfonate salts: trends in coordination behavior of alkali, alkaline earth and transition metals, **216**, 101

solution studies of monovalent metal complex ions of the bi-bracchial lariat ether 7,16-bis(2-methoxyethyl)-1,4,10,13-tetraoxa-7,16-diazacyclooctadecane, **219**, 217

a novel sodium template approach for preparing tetraimino macrocycles of 2,6-diformyl-4-methylphenol and diamino de-

rivatives, **223**, 169

### Alkaline earth cation complexes

synthesis and electrochemical properties of oxo-bridged manganese dimers incorporating alkali and alkaline earth cations, **225**, 191

### Alkaline earth metal complexes

structures of layered metal sulfonate salts: trends in coordination behavior of alkali, alkaline earth and transition metals, **216**, 101

influence of pendant cyanoalkyl side-arms on the stabilities of metal ion complexes of monoaza- and diaza-18-crown-6 ethers in methanol, **219**, 31

a novel route to the second-generation alkaline-earth metal precursors for metal-organic chemical vapour deposition: one-step synthesis of M(hfa)<sub>2</sub>·tetraglyme (M=Ba, Sr, Ca and Hhfa=1,1,1,5,5,5-hexafluoro-2,4-pentanedione), **224**, 203

### Alkaline metal complexes

influence of pendant cyanoalkyl side-arms on the stabilities of metal ion complexes of monoaza- and diaza-18-crown-6 ethers in methanol, **219**, 31

### Alkanes

oxidation of *cis*- and *trans*-pinane with RuO<sub>4</sub> generated in situ, **222**, 115

### Alkene complexes

arylation of alkenes by the palladium(II) acetate–dialkyl sulfide system, **216**, 253

$\pi$ -bonded alkene and arene complexes of silver(I): an electropray mass spectrometric study, **220**, 99

substitution kinetics of some tetracarbonyl( $\eta^2$ -alkene)osmium complexes, **227**, 215

### Alkenes

hydroformylation of alkenes and alkynes using a heterobinuclear Rh–W catalyst, **220**, 187

### Alkoxide complexes

uranium compounds as catalysts for aldolisation reactions, **222**, 91

### Alkoxyallyl complexes

chemistry of  $\pi$ -complexes derived from [(Ph<sub>3</sub>P)<sub>2</sub>Pt( $\eta^3$ -CH<sub>2</sub>CCPh)](O<sub>3</sub>SCF<sub>3</sub>). Interconversion between  $\eta^3$ -alkoxyallyl and  $\eta^3$ -oxoallyl complexes and between  $\eta^3$ -aminoallyl and  $\eta^3$ -iminoallyl complexes, **223**, 1

### Alkoxo complexes

synthesis of some dinuclear niobium chloro alkoxy carboxylates [Nb<sub>2</sub>Cl<sub>4</sub>(OEt)<sub>4</sub>(O<sub>2</sub>CR)<sub>2</sub>], and the X-ray crystal and molecular structure of the compound with R=Ph, **227**, 99

### Alkylation

transition metal complexes with sulfur ligands CIV. Protonation and alkylation of thiolate donors of [Fe(CO)<sub>2</sub>(‘S<sub>4</sub>’)] and [Fe(CO)(‘S<sub>5</sub>’)]: syntheses and properties of Fe(II) complexes with open-chain thioether ligands (‘S<sub>4</sub><sup>2-</sup>’=1,2-bis(2-mercaptophenylthio)ethane(2-); ‘S<sub>5</sub><sup>2-</sup>’=2,2’-bis(2-mercapto-phenylthio)diethylsulfide(2-)), **224**, 45

- alkylation of 2-vinylpyridine and crystal structure of bis[(acetylacetonate)(1-pyridyl)butyl]nickel(II) complex, **227**, 149
- Alkyl complexes
- kinetics and mechanism of the phosphate-catalyzed heterolysis of the metal–carbon  $\sigma$  bond of  $\alpha$ -hydroxyalkyl–chromium(III) complexes. Effects of the non-participating ligands, temperature and pressure, **215**, 173
- an electro spray mass spectrometric study of methylmercury(II) nitrogen base derivatives and their ligand exchange reactions, **215**, 179
- trimethylgallium based phosphinogallanes. Synthesis and molecular structure of  $[\text{Me}_2\text{Ga-P}(\text{SiMe}_3)_2]_2$ , **216**, 267
- alkyl for hydride exchange between alane-trimethylamine and Group IVB metal alkyls, **218**, 47
- a mechanistic study of the homogeneous catalytic hydroformylation of formaldehyde: synthesis and characterization of model intermediates, **218**, 89
- bite angle dependence of the rate of reductive elimination from diphosphine palladium complexes, **220**, 249
- non-isothermal spectrophotometric kinetics applied to inorganic reactions, **221**, 31
- an ethyl–cobalt complex with a buckled equatorial ligand, **221**, 35
- positive and negative ion electro spray mass spectrometric studies of some amino acids and glutathione, and their interactions with alkali metal ions and methylmercury(II), **223**, 103
- free radical-mediated electron transfer in organometallic complexes: homolysis and alkyl group crossover reactions for alkyliron(II) porphyrins, **226**, 231
- volumes of activation for  $\beta$ -elimination and related reactions of chromium(III)–alkyl complexes in aqueous solutions, **227**, 57
- Alkylidene complexes
- investigation of the dithiocarbamate-induced coupling of allylidene and carbonyl ligands on a tungsten center, **220**, 335
- Alkyl sulfide complexes
- new methods for synthesis of platinum(II) dimethyl sulfide complexes, **219**, 155
- Alkyl sulfonium complexes
- the structure of  $(\text{CH}_3)_3\text{SI}_3$ . Comparison between the structure in the solid and liquid state, **215**, 33
- Alkyl-transition metal reagents
- chele- and anticheleselectivity IV. Reaction-accelerating neighboring-group effects in the cross-coupling of vinylbromides with alkyl-transition metal reagents, **220**, 85
- Alkyne
- synthesis of symmetric and unsymmetric 1,4-bis(*p*-R-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289
- Alkyne complexes
- insertion reaction of dimethylacetylenedicarboxylate with the head-to-tail isomer of  $\text{Pt}_2\text{I}_2(\mu\text{-Ppy}_3)_2$ ;  $\text{Ppy}_3 = \text{tris}(2\text{-pyridyl})\text{phosphine}$ , **217**, 209
- the study of isolobal displacement reactions. Synthesis and structures of the heterobimetallic  $\mu$ -alkyne cluster complexes  $\eta^5\text{-RC}_3\text{H}_4(\text{CO})_2\text{M}(\mu\text{-R}^1\text{C}_2\text{R}^2)\text{Co}(\text{CO})_3$  ( $\text{M} = \text{Mo}, \text{W}$ ;  $\text{R} = \text{CH}_3\text{C}(\text{O}), \text{CH}_3\text{O}_2\text{C}$ ), **219**, 93
- some new but-2-yne phosphine complexes of tungsten(II), **219**, 99
- regio- and stereoselective dimerization of 1-alkynes catalyzed by an Os(II) complex, **220**, 5
- rhodium-catalyzed synthesis of 2(5*H*)-furanones from terminal alkynes and non-substituted alkynes under water-gas shift reaction conditions, **220**, 45
- Alkyne hydration
- alkyne hydration using Pt(II) catalysts, **222**, 317
- Alkynes
- hydroformylation of alkenes and alkynes using a heterobimetallic Rh–W catalyst, **220**, 187
- Alkynols
- selective transformations of alkynols catalyzed by ruthenium complexes, **222**, 155
- Allene
- preparation of  $\eta^1$ -1,3-dienyl,  $\eta^1$ -1,2-dienyl and  $\eta^3$ -cyclobutene complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305
- Allene complexes
- reactivity of allene at phosphine-bridged di-iron centres: X-ray crystal structures of  $[\text{Fe}_2(\text{CO})_5\{\mu\text{-}\sigma, \eta^3\text{-C}(\text{O})\text{C}(\text{CH}_2)_2\}(\mu\text{-dppm})]$  and  $[\text{Fe}_2(\text{CO})_4\{\mu\text{-}\eta^3, \eta^3\text{-}(\text{CH}_2)_2\text{C}_2(\text{CH}_2)_2\}(\mu\text{-dppm})] \cdot \text{Et}_2\text{O}$ , **220**, 201
- Allylation
- asymmetric catalysis 87\*. Enantioselective allylation of 1,5-dimethylbarbituric acid with Pd catalysts and new optically active PN ligands, **220**, 63
- ultrasound-promoted palladium-catalyzed carbonyl allylation by allylic alcohols with tin(II) chloride in non-polar solvents, **220**, 155
- Allyl complexes
- synthesis of  $\eta^3$ -allyl–Pd complexes and subsequent formation of N-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- formation of an allyl–tin bond in  $\eta^3$ -allyl(trichlorostannyl)palladium(II) complexes, **222**, 21
- strain, silyl and steric effects on the regioselectivity of palladium(0) catalyzed allyl esters reduction as alternative to the Wittig reaction, **222**, 37
- the preparation of chiral titanium reagents: a comparison of new and known ligands, **222**, 95
- carbon dioxide fixation by  $\text{Cp}_2(\eta^3\text{-allyl})\text{Ti}$  complexes generated from various dienes, **222**, 145
- Allylic ligand complexes
- palladium-assisted formation of carbon–carbon bonds Part 2. Stoichiometric synthesis of spirocyclic compounds. X-ray structure of a  $\pi$ -allylic palladium intermediate, **222**, 1
- Allylidene complexes
- bicyclic carbo- and heterocycles from an allylidene complex and cyclic 1,3-dienes by tandem cyclopropanation/Cope rearrangement, **220**, 327
- Aluminium complexes
- alkyl for hydride exchange between alane-trimethylamine and Group IVB metal alkyls, **218**, 47
- the influence of aluminum trichloride on a configuratively labile lactone-bridged biaryl: quantum chemical calculations and optical spectroscopy, **222**, 247
- use of aluminium as an electron pool. Reductive coupling of iodobenzenes promoted by electron transfer in an Ni/Pb/Al multi redox system, **222**, 323
- Amide complexes
- synthesis of  $\alpha$ -hydroxy acid derivatives by the photochemical reaction of *O*-silylchromium(0) carbene complexes and nucleophiles, **222**, 261
- Amide compounds
- benzotriazole stabilized lithium intermediates as a route to the elaboration of N-substituents in amides, **220**, 67
- Amide coordination
- hydrolytic cleavage of peptides by palladium(II) complexes is enhanced as coordination of peptide nitrogen to palladium(II) is suppressed, **217**, 21
- Amidine complexes

- reactions of aziridine with platinum(II) nitriles. Formation of (aziridino)amidines and 2-imidazolines and X-ray structure of *trans*-[PtCl<sub>2</sub>{N(H)=C(Ph)NCH<sub>2</sub>CH<sub>2</sub>}]<sub>2</sub>, **222**, 327
- Amidoxime complexes**  
the synthesis and crystal structure of ethylenedinitrilotetraacetamidoxime nickel(II) sulfate trihydrate, **227**, 171
- Amine complexes**  
binuclear oxo-bridged iron(III) complexes of *N,N'*-bis(2-pyridylmethyl)ethane-1,2-diamine (bispicen) and related ligands, **216**, 21  
unexpected five-coordination in di(1,3-diaminopropane)-3,5-dinitrobenzoatocopper(II) 3,5-dinitrobenzoate. Comparison between the coordinated and non-coordinated anions, **216**, 57  
structure and spectroscopic properties of [di(3-aminopropyl)-amine-*N,N',N''*][iminodiacetato-*N,O,O'*]chromium(III) perchlorate, **216**, 155  
nucleophilic addition of water to coordinated di-(2-pyridyl)-ketone (DPK) in rhenium(V) complexes. Synthesis and crystal structure of the adduct (DPK·H)<sup>+</sup>[ReOCl<sub>2</sub>-(DPK·OH)<sub>2</sub>Cl], **217**, 141  
bromo derivatives of amine and phosphine complexes of cyanodihydroborane. Synthesis and reactivity, **218**, 21  
transition metal complexes with sulfur ligands Part CII.  
Changes in configuration and reactivity caused by alkyl substituents R in [Fe(L)(<sup>n</sup>R<sub>4</sub>S<sub>4</sub>)] complexes (L=CO, NO<sup>+</sup>, NO, PMe<sub>3</sub>, N<sub>2</sub>H<sub>4</sub>; R=CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub><sup>-</sup>), **219**, 75  
equilibrium and kinetic study of piperidine binding to phthalocyaninatoiron(II) in dimethyl sulfoxide, **219**, 199  
exchange of azine ligands in *fac*-Re(CO)<sub>3</sub>(4-cyanopyridine)<sub>2</sub>Cl and *fac*-Re(CO)<sub>3</sub>(quinoline)<sub>2</sub>Cl, **221**, 161  
synthesis and properties of bis(amine)carboxyhydroboron(1+) cations, **223**, 155  
stability of metal ion complexes of *N,N'*-bis(2-hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid, **224**, 151  
charge type effects on the temperature dependence of intervalence transfer in ferrocyanide substituted ruthenium(III) amine and aminopolycarboxylate compounds, **224**, 189  
the synthesis, X-ray structure and hydrolysis kinetics of chloro(2-aminomethylpyridine)(triamine)chromium(III) salts, **227**, 11  
synthesis, crystal structure and characterization of nickel(0) acrylonitrile complexes with a bidentate N-donor ligand, **227**, 137
- Amine cryptand complexes**  
azacryptand host for lead: NMR and structural studies, **218**, 5
- Amine N-oxide complexes**  
bis(2-mercaptopyridine-*N*-oxide)oxovanadium(IV): synthesis, structure and its conversion to the dichloro complex, **219**, 213
- Amino acid complexes**  
mixed-ligand copper(II) complexes with iminodiacetato and 2- or 4-methylimidazole: molecular and crystal structure of the 'remote' isomer iminodiacetato(5-methylimidazole)-copper(II) monohydrate, [Cu(IDA)(5MeImH)]·H<sub>2</sub>O, **215**, 73  
a novel amino acid-binding receptor based on weak non-covalent ligand-ligand interactions, **216**, 249  
'insulation' to lattice effects on quadrupole splitting in the Mössbauer spectra of salts of [Fe<sub>4</sub>S<sub>4</sub>(Cys)<sub>4</sub>]<sup>2-</sup>, a ferredoxin model, **217**, 3  
interaction of the lysine side chain amino group with Cu<sup>II</sup> in (glycyl-L-lysine<sup>2-</sup>)Cu, **217**, 33  
bis(2-diphenylphosphinoethyl)phenylphosphineruthenium(II) complexes of amino acids and dipeptides, **217**, 51  
nitrogen-15 NMR coordination shifts of β-alanine and glycine in cobalt(III) complexes, **217**, 213  
ternary Pt(II)-amino acid-nucleotide complexes: kinetics of formation, **218**, 11  
substituent effect on the coordination ability of the amide group of N-protected amino acids, **218**, 53  
lithium coordination to amino acids and peptides. Synthesis, spectroscopic characterization and structure determination of lithium complexes of neutral and anionic glycine and diglycine, **218**, 121  
a study of the reactivity of (methyl 2-acetamidoacrylate)tricarbonyliron(0) leading to a novel synthesis of β,β,β-trialkyl α-amino acids, **220**, 137  
positive and negative ion electrospray mass spectrometric studies of some amino acids and glutathione, and their interactions with alkali metal ions and methylmercury(II), **223**, 103  
ternary complexes of *cis*- and *trans*-Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (*cis*-, *trans*-DDP) with 9-methylguanine (9-MeG), the dipeptides glycylglycine (glygly), glycyl-L-alanine (glyala), glycyl-L-2-aminobutyric acid (gly-2-aba), glycyl-L-norvaline (glynval), glycyl-L-norleucine (glynleu), and of *trans*-Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> with Na-L-acetylhistidine, **227**, 17
- Aminoallyl complexes**  
chemistry of π-complexes derived from [(Ph<sub>3</sub>P)<sub>2</sub>Pt(η<sup>3</sup>-CH<sub>2</sub>CCPh)](O<sub>3</sub>SCF<sub>3</sub>). Interconversion between η<sup>3</sup>-alkoxyallyl and η<sup>3</sup>-oxoallyl complexes and between η<sup>3</sup>-aminoallyl and η<sup>3</sup>-iminoallyl complexes, **223**, 1
- Aminocarbyne complexes**  
electron-transfer activation of the aminocarbyne and the hydrogen isocyanide complexes *trans*-[ReCl(CNH<sub>n</sub>)-(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>][BF<sub>4</sub>]<sub>n-1</sub> (n=2 or 1). Interconversion of coordinated CNH<sub>2</sub> and CNH, **226**, 9
- Amino complexes**  
cyclic bis(amino)organotin cations, stabilized by π-coordination and new spirocyclic stannole derivatives, **216**, 51  
conformational flexibility of 1,3-diaminopropane in chelate rings. Redetermination of the structure of *trans*-dinitratobis(1,3-diaminopropane)copper(II), **218**, 151
- Aminopolycarboxylate complexes**  
charge type effects on the temperature dependence of intervalence transfer in ferrocyanide substituted ruthenium(III) amine and aminopolycarboxylate compounds, **224**, 189
- Ammine complexes**  
interaction of cyanopentaamminechromium(III) with mercuric ion. Electrochemical, spectral and preparative study of adduct formation and linkage isomerism, **216**, 113  
molecular mechanical and quantum chemical study on the species involved in the hydrolysis of *cis*-diamminedichloroplatinum(II) and substituted bis(ethylenediamine)dichloroplatinum(II) complexes Part I. Reactants and products, **217**, 159  
synthesis and characterisation of cyanide bridged heterobinuclear complexes based on Group VI octacyanometalate(IV) and pentaammineosmium(III) ions: a comparison with Group VIII hexacyano analogues, **219**, 121  
crystal structure and ligand field states of [Cr(NH<sub>3</sub>)<sub>5</sub>NCO](NO<sub>3</sub>)<sub>2</sub>, **221**, 93  
properties of the mixed-valence binuclear complex ion, [(NH<sub>3</sub>)<sub>5</sub>Ru<sup>III</sup>-NC-Os<sup>III</sup>(CN)<sub>5</sub>], **223**, 37  
the synthesis, and UV-Vis, electrochemical, photochemical and intervalence characterization of tetraammine and tris chelated ruthenium(II) complexes bound to the bridging ligand dipyrido(2,3-*a*;2',3'-*h*)phenazine (dpop), **224**, 27  
solvent trapped valency in the [bis(pentaammineruthenium)-(μ-1,4-dicyanamidobenzene)] trication, **225**, 123

- comparison of the rates of substitution in  $[\text{Ru}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  and  $[\text{Os}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  by hexacyano complexes: substitution coupled to electron transfer, **225**, 173
- electrocatalysts for the four-electron reduction of dioxygen based on adsorbed cobalt tetrapyrrolylporphyrin molecules linked by aquaammine complexes of ruthenium(II), **225**, 215
- the reduction of pentaammine(phenanthroline-2-carboxamido-*N*)cobalt(III) by titanium(III) and vanadium(II), **225**, 245
- the electronic structure of pentaammine(pyrazine) ruthenium(II) and (III): the metal–ligand  $\pi$  conjugation and its implications in electron delocalization, **226**, 43
- solvent, electrolyte and solute shape effects on optical electron transfer in mixed-valence ruthenium ammine dimers, **226**, 137
- Analytical detection**  
estrogen derivatives of transition metal complexes for analytical detection enhancement. Part II, **218**, 207
- Anchored complexes**  
formation of an anchored copper(II) complex on a thiourea-functionalized silica gel by *in situ* modification of the tethered ligating function, **221**, 183
- Angular overlap model**  
crystal structure and ligand field states of  $[\text{Cr}(\text{NH}_3)_5\text{NCO}](\text{NO}_3)_2$ , **221**, 93
- Anion complexes**  
preparation of  $\eta^1$ -1,3-dienyl,  $\eta^1$ -1,2-dienyl and  $\eta^3$ -cyclobutene complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305
- outer-sphere coordination of polycyanometallate anions with polyammonium macrocycles: a spectrophotometric study, **227**, 71
- NQR studies of polyhalocuprate(I) anions, **227**, 153
- Anionic complexes**  
simultaneous binding of soft and hard metals to a pyrimidine nucleobase: *trans*- $\text{K}_2[\text{PtI}_2(1\text{-MeU})_2] \cdot 6\text{H}_2\text{O}$ , an anionic Pt-1-methyluracil (1-MeU) complex arranged in circles, **218**, 117
- Anodic oxidation**  
synthesis, properties and reactions of sila-functional  $\alpha$ -heteroatom-substituted organosilicon compounds, **220**, 129
- Anthracene complexes**  
binuclear complexes from 1,8-substituted anthracene derivatives. The crystal structure of  $\{[\mu\text{-}[1,8\text{-anthracenedicarboxaldehydebis}(\text{dimethylhydrazone})]\}\text{tetrachlorobis}(\eta^2\text{-ethylene})\text{di-platinum}\}$ , **221**, 151
- Anticancer drug complexes**  
synthesis and hydrolysis of an ester of the ‘one ring open’ hydrolysis product of the anticancer drug 1,2-bis(3,5-dioxopiperazin-1-yl)propane, ‘Razoxane’, **215**, 55
- Antiferromagnetism**  
pyrazine-mediated antiferromagnetic intermolecular exchange in mixed-valent diruthenium tetracarboxylates, **215**, 203
- antiferromagnetism tuned by the topology of the intervening bridging ligand. Crystal structure and magnetic properties of the  $(\mu\text{-phthalato})\{[\text{bis}(\text{piperidine})]\text{Cu}(\text{II})\}$  chain complex, **218**, 167
- Antitumor agents**  
synthesis and X-ray crystal structure of *cis*-1,4-diaminocyclohexanetetrachloroplatinum(IV): a new antitumor agent, **219**, 193
- thiopyridyl triazine derivatives and their platinum complexes: a new class of potential antitumor agents, **227**, 175
- Antitumour activity**  
the first isolated antineoplastic Ru(IV) complex: synthesis and structure of  $[\text{Cl}_2(1,2\text{-cyclohexanediaminotetraacetate})\text{Ru}] \cdot 2\text{H}_2\text{O}$ , **224**, 15
- Aqua complexes**  
synthesis, spectral and redox properties of a new series of aqua complexes of ruthenium(II), **221**, 125
- synthesis, structure and solid state polymerization of di-aquabis(propynoato)magnesium, **222**, 339
- spectroscopic studies of the *trans*- $[\text{VOCl}_2(\text{H}_2\text{O})_2]$  chromophore in its hydrogen bonding adduct with 18-crown-6, **227**, 91
- Aquation**  
the steric effect on the ring opening process of the decarboxylation of *cis*-carbonato-bis(diamine)cobalt(III) ions, **221**, 9
- Aqueous micelles**  
redox behavior of the iron–sulfur cluster  $[\text{Fe}_4\text{Cp}_4\text{S}_5][\text{PF}_6]_2$  in protic organic solvents and aqueous micellar solutions, **226**, 341
- Aquo complexes**  
preparation and structure of  $[\text{PhPt}(\text{OCO}_2\text{H})(\text{PEt}_3)_2]$  and  $[\text{PhPt}(\text{OH}_2)(\text{PEt}_3)_2]\text{BF}_4$ , **218**, 199
- Arene complexes**  
synthesis and structure of  $\text{ZrCl}_4(\eta^6\text{-C}_6\text{H}_5\text{PMe}_2)_2(\text{PMe}_2\text{Ph})_2$ , an arene complex of zirconium(II), **217**, 187
- organometallic chelating ligands. Synthesis of a new anionic tripod like ligand with an  $\text{N,N',O}$ -donor set. X-ray crystal structure of  $[(\eta^6\text{-C}_6\text{Me}_6)\text{Ru}\{\mu\text{-P}(\text{O})(\text{OMe})_2\}(\mu\text{-Pz})_2\text{Re}(\text{CO})_3]$  ( $\text{Me}$  = methyl,  $\text{Pz}$  = pyrazolate), **219**, 67
- $^1\text{H}$  NMR analysis of some  $\eta^6$ -(substituted aryl phenylsulfonyl-acetonitrile)- $\eta^5$ -cyclopentadienyliron cations, **219**, 221
- $\pi$ -bonded alkene and arene complexes of silver(I): an electro-spray mass spectrometric study, **220**, 99
- $\pi$ -facial selectivities in the coordination of an optically pure fused cyclopentadienyl ligand to arene ruthenium moieties. NMR spectral and structural characterization of a single diastereomer of  $[(\text{PCp})\text{Ru}(\text{Me}_6\text{C}_6)]\text{PF}_6$ , **221**, 41
- arene migration in ruthenium clusters: a kinetic study of the isomerisation of  $\text{Ru}_5\text{C}(\text{CO})_{12}(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_6)$  to  $\text{Ru}_5\text{C}(\text{CO})_{12}(\eta^6\text{-C}_6\text{H}_6)$ , **222**, 299
- Aromatic heterocycle complexes**  
charge and electron transfer from metal-to-carbon bonds of main group organometallics  $\text{MR}_n$  ( $\text{M}$  = Al, Ga, Zn) to aromatic N-heterocycles: colored precursor compounds and radical complex formation, **225**, 23
- Aromatic interaction**  
effect of aromatic–aromatic interaction on ligand binding to zinc porphyrins, **224**, 157
- Arsenate complexes**  
hydrothermal synthesis and crystal and molecular structure of a three-dimensional vanado-arsenate  $(\text{NH}_4)_3(\text{VO})(\text{V}_2\text{O}_3)(\text{AsO}_4)_2(\text{HAsO}_4)$ , exhibiting both  $\{\text{V-O-V}\}_\infty$  chains and isolated  $\{\text{V}_2\text{O}_3\}$  units, **218**, 59
- Artificial peptidase**  
hydrolytic cleavage of peptides by palladium(II) complexes is enhanced as coordination of peptide nitrogen to palladium(II) is suppressed, **217**, 21
- Arylation**  
arylation of alkenes by the palladium(II) acetate–dialkyl sulfide system, **216**, 253
- Aryl complexes**  
NMR parameters as steric probes for arylcyclopentadienyl iron complexes, **215**, 139
- direct nitration of aryl groups  $\sigma$ -bound to ruthenium(II), **220**, 261
- Aryl halide**  
synthesis of symmetric and unsymmetric 1,4-bis(*p*-R-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289
- Asymmetric catalysis**

- asymmetric catalysis 87. Enantioselective allylation of 1,5-dimethylbarbituric acid with Pd catalysts and new optically active PN ligands, **220**, 63
- designed binaphthyl-derived titanium complexes: a new type of asymmetric catalyst for the carbonyl-ene reaction with glyoxylate, **222**, 71
- synthesis and structure of new chiral ferrocenylphosphines for asymmetric catalysis, **222**, 213
- Asymmetric synthesis**
- optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220**, 283
- Atom–atom interaction model**
- a unified view of inorganic stereochemistry and stereochemical changes, **219**, 187
- Atomic weights**
- 1993 Table of Standard Atomic Weights abridged to Five Significant Figures, **217**, 217
- Atropisomerism**
- synthesis, structure, dynamics and first atropisomer-selective cleavage of the chromium tricarbonyl complex of a lactone-bridged biaryl, **222**, 255
- Azacryptand complexes**
- azacryptand host for lead: NMR and structural studies, **218**, 5
- Azamacrocyclic ligand complexes**
- stability constants of the lanthanide(III)-1,4,7,10-tetraazacyclododecane-*N,N',N'',N'''*-tetraacetate complexes, **221**, 165
- Azaplatinadecaborane complexes**
- ten-vertex polyhedral azametallaborane chemistry. An *arachno*-6,9-azaplatinadecaborane and a *nido*-6,8-azaruthenadecaborane, **227**, 163
- Azarhenacycle complexes**
- insertion of NH into the rhenium–carbon bonds of a Fischer-type rhenacyclobutadiene. Preparation and characterization of azarhenacyclic complexes, **222**, 199
- Azaruthenadecaborane complexes**
- ten-vertex polyhedral azametallaborane chemistry. An *arachno*-6,9-azaplatinadecaborane and a *nido*-6,8-azaruthenadecaborane, **227**, 163
- Azauracil complexes**
- analysis of [(dien)Pd]<sup>2+</sup> binding to uracil and azauracils by proton NMR spectroscopy, **223**, 131
- Azido complexes**
- transition metal complexes with sulfur ligands CVI. Phosphine and azido [Fe(L)(‘N<sub>H</sub>S<sub>4</sub>’)] complexes; structural characterization of the key complex [Fe(‘N<sub>H</sub>S<sub>4</sub>’)]·THF (L = PMe<sub>3</sub>, PBU<sub>3</sub>, PMe<sub>2</sub>Ph, PMePh<sub>2</sub>, N<sub>3</sub><sup>−</sup>; ‘N<sub>H</sub>S<sub>4</sub>’<sup>2−</sup> = dianion of 2,2'-bis(2-mercaptophenylthio)diethylamine), **224**, 61
- Azine complexes**
- exchange of azine ligands in *fac*-Re(CO)<sub>3</sub>(4-cyanopyridine)<sub>2</sub>Cl and *fac*-Re(CO)<sub>3</sub>(quinoline)<sub>2</sub>Cl, **221**, 161
- reactions of [Au(C<sub>6</sub>F<sub>5</sub>)(SC<sub>4</sub>H<sub>8</sub>)] with diazoalkanes. Synthesis and molecular structures of [Au(C<sub>6</sub>F<sub>5</sub>)(Ph<sub>2</sub>C=N–N=CPh<sub>2</sub>)], **222**, 267
- Aziridine reactions**
- reactions of aziridine with platinum(II) nitriles. Formation of (aziridino)amidines and 2-imidazolines and X-ray structure of *trans*-[PtCl<sub>2</sub>{N(H)=C(Ph)NCH<sub>2</sub>CH<sub>2</sub>}]<sub>2</sub>, **222**, 327
- Azoxybenzene complexes**
- cyclopalladated *p*-azoxyanisole complexes. Synthesis and structure of {[bis(2-phenylpyridine)][4'-(methoxyphenyl)-*NNO*-azoxy-*N*'<sup>2</sup>]-4-methoxyphenyl-2-ato}palladium(II)}-(tetrafluoroborate), **218**, 211
- Azulene complexes**
- preparation and crystal structure of *anti*-( $\eta^3$ : $\eta^5$ -azulene)-pentakis(trimethylphosphine)dnicobalt. Reactions with carbon monoxide and solution dynamics, **218**, 143
- Barbituric acid**
- asymmetric catalysis 87. Enantioselective allylation of 1,5-dimethylbarbituric acid with Pd catalysts and new optically active PN ligands, **220**, 63
- Benzocyclobutenedione complexes**
- reactions of benzocyclobutene chromium complexes with carbon, nitrogen and oxygen nucleophiles: nucleophilic addition and unexpected proximal ring opening reactions, **220**, 175
- Benzocyclobutenone complexes**
- reactions of benzocyclobutene chromium complexes with carbon, nitrogen and oxygen nucleophiles: nucleophilic addition and unexpected proximal ring opening reactions, **220**, 175
- Benzonaphthopyranone**
- the influence of aluminum trichloride on a configuratively labile lactone-bridged biaryl: quantum chemical calculations and optical spectroscopy, **222**, 247
- Benzosemiquinone complexes**
- indium(III) halide–3,5-di-*tert*-butyl-*o*-benzosemiquinone systems, **225**, 207
- Benzotriazole compounds**
- benzotriazole stabilized lithium intermediates as a route to the elaboration of N-substituents in amides, **220**, 67
- Benzylviologen salts**
- crystal structure and solid-state fluorescence of an oxomolybdenum(V)–benzylviologen electron donor–acceptor salt, **226**, 25
- Bicarbonato complexes**
- preparation and structure of [PhPt(OCO<sub>2</sub>H)(PEt<sub>3</sub>)<sub>2</sub>] and [PhPt(OH<sub>2</sub>)(PEt<sub>3</sub>)<sub>2</sub>]BF<sub>4</sub>, **218**, 199
- Bicyclo[3.2.1]octadiene**
- bicyclic carbo- and heterocycles from an allylidene complex and cyclic 1,3-dienes by tandem cyclopropanation/Cope rearrangement, **220**, 327
- Bicyclo[3.2.1]oxaheptadiene**
- bicyclic carbo- and heterocycles from an allylidene complex and cyclic 1,3-dienes by tandem cyclopropanation/Cope rearrangement, **220**, 327
- Bidentate amine complexes**
- synthesis, spectral and redox properties of a new series of aqua complexes of ruthenium(II), **221**, 125
- synthesis, characterization and X-ray crystal structure of ethylenediamine(oxotato)palladium(II), **221**, 173
- cage escape yields for photoinduced bimolecular electron transfer reactions of Re(I) complexes, **225**, 41
- Bidentate amine ligand complexes**
- synthesis and X-ray crystal structure of *cis*-1,4-diaminocyclohexanetetrachloroplatinum(IV): a new antitumor agent, **219**, 193
- Bidentate amino ligand complexes**
- three-coordinate Pt(0)  $\eta^2$ -complexes: electrophilic hydrogen attack through oxidative-addition of protic acids, **219**, 169
- Bidentate carboxylate complexes**
- structural characterization of bidentate carboxylate derivatives of copper(I) bistrifluorophosphine, **227**, 223
- Bidentate chelate ligand complexes**
- stabilities of 1,2-diethyl-3-hydroxy-4-pyridinone chelates of divalent and trivalent metal ions, **223**, 21
- Bidentate isonitrile complexes**
- synthesis and characterization of a bidentate chelating diisonitrile ligand with two thioetheral sulfurs in a bridging group and its copper(I), rhodium(I) and palladium(II) complexes, **227**, 113
- Bidentate ligand complexes**
- mixed-ligand complexes of rhenium II. Synthesis and characterization of [ReN(X)(Me<sub>2</sub>PhP)<sub>2</sub>(R<sub>2</sub>tcb)] complexes (X = I,

- $N_3$ , SCN, CN;  $R_2tcb^- = N-(N,N\text{-dialkyl- thiocarbamoylbenzamidate})$ ). X-ray crystal structure of  $[ReN(N_3)(Me_2PhP)_2(Et_2tcb)]$ , **215**, 159
- solvent effects on acetylacetonato iron complexes, **215**, 185
- principal component analysis of the structures of the 1,2-diaminobenzene derivatives chelated to metal ions: correlation between structural parameters and redox behavior, **215**, 219
- synthesis of bis- or tris(pentafluorophenyl) thallium(III) complexes with bidentate ligands. X-ray crystal structure of  $[Tl(C_6F_5)_2(acac)\{(O)Ph_2PCH_2PPh_2(O)\}]$ , **215**, 229
- unexpected five-coordination in di(1,3-diaminopropane)-3,5-dinitrobenzoatocopper(II) 3,5-dinitrobenzoate. Comparison between the coordinated and non-coordinated anions, **216**, 57
- magnetic properties and crystal structure of  $\{\mu\text{-pyrazine}\}\text{-bis}\{(aqua)(2,6\text{-xylylaminiodiacetato})\text{copper(II)}\}$ . A pyrazine-bridged dinuclear copper(II) complex, **217**, 39
- spectroscopic and structural studies on volatile gallium  $\beta$ -diketonates as potential precursors for MOCVD, **217**, 71
- unusually stable peroxocopper complexes. Stoichiometry, products and kinetics of oxidation of the dimeric copper(I) complex  $[LCuBr]_2$  ( $L = N,N'\text{-diethylethylenediamine}$ ) by dioxygen in methylene chloride from  $-51$  to  $30^\circ\text{C}$ , **217**, 109
- the synthesis and crystal structure of  $[Cr(acacen)py_2][ZnCl_3py]$ , **217**, 195
- organometallic complexes of ruthenium(IV) with bidentate oxygen and nitrogen donor ligands, **218**, 81
- conformational flexibility of 1,3-diaminopropane in chelate rings. Redetermination of the structure of *trans*-dinitratobis(1,3-diaminopropane)copper(II), **218**, 151
- dimolybdenum(II) and diruthenium(II) complexes with 2-substituted 7-methyl-1,8-naphthyridines as bridging ligands, **219**, 23
- synthesis, crystal structure and spectroscopic properties of bis(diphenyldithiophosphinato)cadmium(II), **219**, 115
- variation of the exchange interaction in oximato-bridged  $Cu^{II}M^{II}$  dimers ( $M = Cu, Ni, Mn$ ). Crystal structure of  $[Cu(pdmg)Cu(bipy)(H_2O)_2](ClO_4)_2 \cdot H_2O$ , **219**, 179
- synthesis, structure and stability of complexes of lanthanide nitrates with  $[(MeO)_2P(O)]_2C(OH)Ph$  and the isomeric  $(MeO)_2P(O)OCHPhP(O)(OMe)_2$ , **223**, 43
- the synthesis, and UV-Vis, electrochemical, photochemical and intervalence characterization of tetraammine and tris chelated ruthenium(II) complexes bound to the bridging ligand dipyrido(2,3-*a*;2',3'-*h*)phenazine (dpop), **224**, 27
- di-manganese carbonyl compounds containing bis(dicyclohexylphosphino)methane (dcpm) ligation, **224**, 105
- multinuclear ( $^{31}P$ ,  $^{77}Se$ ,  $^{113}Cd$ ,  $^{199}Hg$ ) magnetic resonance, electrospray mass spectrometric and electrochemical studies on the reactions of  $M(S_2P(OEt)_2)_2$  ( $M = Cd, Hg$ ) with potentially bidentate mixed Group 15/Group 16 and Group 16 donor ligands, **224**, 137
- electrochemical, photophysical and photochemical properties of a hexanuclear ruthenium(II) bis(2,2'-bipyridyl) complex containing a tris(pyrazyltriazole)benzene ligand, **225**, 67
- dynamical investigations of the catalytic mechanisms of water oxidation by the  $[(bpy)_2Ru(OH_2)]_2O^{4+}$  ion, **226**, 179
- preparation, structural and physical studies of mercury(II) halide complexes with the  $N,N'$ -chelating biheteroaromatic ligand 2-(2'-pyridyl)quinoxaline, **227**, 129
- synthesis, crystal structure and characterization of nickel(0) acrylonitrile complexes with a bidentate N-donor ligand, **227**, 137
- Bidentate phosphine complexes
- activation of molecular oxygen by iron nitrosyls in the presence of the bidentate phosphines 1,2-bis(diphenylphosphino)ethane and ethene, **221**, 133
- solid state and solution studies of a T-shaped gold(I) complex:  $Au[(C_6H_5)_2PCH_2N(C_6H_5)_2]_2Cl$ , **227**, 85
- Bidentate phosphine ligand complexes
- molybdenum carbonyl derivatives containing eight- and nine-membered diphosphine chelating rings: the synthesis of  $[Mo(CO)_4(dpppe)]$  ( $dpppe = Ph_2P(CH_2)_5PPh_2$ ) and  $[Mo(CO)_4(dpph)]$  ( $dpph = Ph_2P(CH_2)_6PPh_2$ ) and the crystal structure of the latter, **218**, 9
- Bimetallic complexes
- photoinduced intramolecular electron transfer in bimetallic donor-acceptor assemblies, **226**, 79
- early/late bimetallic complexes:  $MCo_2$  ( $M = Ti, Zr$ ) derivatives, **227**, 201
- Bimolecular electron transfer
- cage escape yields for photoinduced bimolecular electron transfer reactions of Re(I) complexes, **225**, 41
- bimolecular electron transfer at high driving forces. Photoreactions of gold(I) and iridium(I) complexes with pyridinium derivatives, **225**, 319
- BINAP complexes
- an improved synthesis of Ru(BINAP) type complexes, **223**, 165
- Binaphthyl complexes
- designed binaphthyl-derived titanium complexes: a new type of asymmetric catalyst for the carbonyl-ene reaction with glyoxylate, **222**, 71
- Binding constants
- outer-sphere coordination of polycyanometallate anions with polyammonium macrocycles: a spectrophotometric study, **227**, 71
- Binuclear complexes
- binuclear complexes from 1,8-substituted anthracene derivatives. The crystal structure of  $[\{\mu\text{-}[1,8\text{-anthracenedicarboxaldehydebis(dimethylhydrazone)}]\}\text{tetrachlorobis}(\eta^2\text{-ethylene})\text{-diplatinum}]$ , **221**, 151
- binuclear platinum(II) complexes of 1,1,4,7,10,10-hexaphenyl-1,4,7,10-tetraphosphadecane (P4) containing various bridging ligands as spacers between the P4 coordination units, **223**, 13
- Bioctahedral complexes
- a new type of face-sharing bioctahedral triply bonded dinobium cluster  $(PyH)[Nb_2(\mu\text{-Cl})_2(\mu\text{-THT})Cl_3(Py)_3](THF)(C_6H_6)_{0.5}$ , made by a designed synthesis, **227**, 191
- Biogenic amine complexes
- spectroscopic studies on complex formation and non-covalent interactions in ternary palladium(II) systems involving spermidine and 2,3-diaminopropionate or 2,4-diaminobutyrate, **219**, 107
- Biphenyl complexes
- synthesis and structure of  $Cr(CO)_3$  complexes of biphenyl compounds with axial chirality, **222**, 63
- Bipyridine complexes
- simultaneous photooxidation of  $[Ru(bpy)_2Cl_2]$  and photoreduction of  $[Ru(bpy)_2Cl_2]^+$  in several solvent systems, **226**, 247
- Bite angle
- bite angle dependence of the rate of reductive elimination from diphosphine palladium complexes, **220**, 249
- Bond homolysis
- free radical-mediated electron transfer in organometallic complexes: homolysis and alkyl group crossover reactions for alkyliron(II) porphyrins, **226**, 231
- Borane complexes
- ten-vertex polyhedral mixed diheteromonometalaborane chemistry. Directed synthesis of *arachno*-9,6,8-platinathiacarba-

- decaboranes, **218**, 1
- bromo derivatives of amine and phosphine complexes of cyanodihydroborane. Synthesis and reactivity, **218**, 21
- Borate complexes**
- coordination study of D-xylo 5-hexulosonic acid with borate, molybdate and tungstate in aqueous solution by nuclear magnetic resonance spectroscopy, **221**, 69
- Boron complexes**
- reactions of phosphine–boron trihalide adducts with carbonyl-bis(triethylphosphine)rhodium(I) halides, **215**, 67
- bromo derivatives of amine and phosphine complexes of cyanodihydroborane. Synthesis and reactivity, **218**, 21
- ten-vertex polyhedral dicarbaborane chemistry: new positional isomers of cluster constituents in the ten-vertex *arachno* dicarbaborane family: the parent [*arachno*-5,10-C<sub>2</sub>B<sub>8</sub>H<sub>13</sub>]<sup>−</sup> anion and the isomeric ligand derivatives *exo*-9-L-*arachno*-5,6-C<sub>2</sub>B<sub>8</sub>H<sub>12</sub> and *exo*-6-L-*arachno*-5,10-C<sub>2</sub>B<sub>8</sub>H<sub>12</sub>, **221**, 5
- synthesis and properties of bis(amine)carboxyhydroboron(1+) cations, **223**, 155
- Bridged biaryls**
- synthesis, structure, dynamics and first atropisomer-selective cleavage of the chromium tricarbonyl complex of a lactone-bridged biaryl, **222**, 255
- Bridgehead reactivity**
- oxidation of *cis*- and *trans*-pinane with RuO<sub>4</sub> generated in situ, **222**, 115
- Bridging carbonyl**
- the vibrations of bridging carbonyl groups as more sensitive indicators of molecular geometry than terminal carbonyl groups in transition metal carbonyl cluster compounds, **227**, 241
- Bromazepam complexes**
- metal complexes of benzodiazepines. Kinetic properties of bromazepam toward square planar dimethyl sulfoxide complexes of platinum(II) and palladium(II), **215**, 41
- Bromide complexes**
- the preparation, properties and crystal structure of a salt containing the hexabromochromate(III) anion, **227**, 167
- Bromide compounds**
- magnesium bromide in Grignard reagent formation, **222**, 365
- Cadmium complexes**
- synthesis and characterization of Cd(II) complexes of cinoxacin. Crystal structure of a dicadmium complex containing two monoatomic-bridging carboxylate oxygen atoms, **217**, 149
- zinc(II) and cadmium(II) complexes of the potentially tridentate ligands 2-[[2-pyridinylmethyl]amino]methyl]phenol (HSALAMP) and 2-[[[2-(2-pyridinyl)ethyl]amino]methyl]phenol (HSALAEP). X-ray crystal structures of the [Zn(SALAMP)(NO<sub>3</sub>)<sub>2</sub>] and the [Cd(SALAEP)(NO<sub>3</sub>)<sub>2</sub>] dimers, **219**, 55
- synthesis, crystal structure and spectroscopic properties of bis(diphenyldithiophosphinato)cadmium(II), **219**, 115
- multinuclear (<sup>31</sup>P, <sup>77</sup>Se, <sup>113</sup>Cd, <sup>199</sup>Hg) magnetic resonance, electrospray mass spectrometric and electrochemical studies on the reactions of M(S<sub>2</sub>P{OEt}<sub>2</sub>)<sub>2</sub> (M = Cd, Hg) with potentially bidentate mixed Group 15/Group 16 and Group 16 donor ligands, **224**, 137
- Calculation**
- basis set effects on the direct calculation of intermolecular coupling elements for electron transfer reactions, **226**, 237
- Calix[4]arene**
- a 'calix[4]arene' porphyrin as a new host and an oxygen carrier model, **224**, 1
- Calixarene complexes**
- luminescent europium(III) and terbium(III) ions encapsulated in a 2-aminoethoxy or carbamoyloxy substituted calixarene host, **227**, 293
- Calorimetry**
- thermometric determination of electrochemical Peltier heat (thermal effect associated with electron transfer) of some redox couples, **226**, 69
- direct solution calorimetric measurements of enthalpies of proton and electron transfer reactions for transition metal complexes. Thermochemical study of metal–hydride and metal–metal bond energies, **227**, 285
- Carbamate complexes**
- the anomalous EPR spectra of the iron(III) dichalcogenocarbamates, **217**, 101
- Carbanion complexes**
- addition of carbanions to tris(trimethylphosphine)-(cyclooctadiene)iridium(I) chloride: three different modes of reaction depending on the nature of the carbanion, **222**, 135
- Carbene complexes**
- silicon-containing carbene complexes 16. Intramolecular agostic stabilization versus nucleophilic addition to the 16-electron carbene complex (CO)<sub>4</sub>(L)W=C(NMe<sub>2</sub>)SiPh<sub>2</sub>Me, **220**, 73
- dirhodium(II) tetrakis[*N,N*-dimethyl-2-pyrrolidone-5(*S*)-carboxamide]. Structural effects on enantioselection in metal carbene transformations, **220**, 193
- stereoselective aldol addition reactions of Fischer carbene complexes via electronic tuning of the metal center for enolate reactivity, **220**, 215
- carbonyl carbene coupling on cationic η<sup>2</sup>-carbene complexes of molybdenum and tungsten, **222**, 77
- steric and electronic tuning of Michael addition reactions induced by pentacarbonyl metal fragments, **222**, 207
- synthesis of α-hydroxy acid derivatives by the photochemical reaction of *O*-silylchromium(0) carbene complexes and nucleophiles, **222**, 261
- manganese and tungsten carbene complex mediated intramolecular Diels–Alder reactions, **222**, 291
- a one-pot synthesis of cyclopropanes from aromatic aldehydes and styrene via iron carbene complexes, **223**, 159
- Carbene coupling**
- carbonyl carbene coupling on cationic η<sup>2</sup>-carbene complexes of molybdenum and tungsten, **222**, 77
- Carbido atoms**
- the formation of carbides in hexaruthenium carbonyl clusters, **227**, 197
- Carbohydrate complexes**
- coordination study of D-xylo 5-hexulosonic acid with borate, molybdate and tungstate in aqueous solution by nuclear magnetic resonance spectroscopy, **221**, 69
- Carbohydrate derivative complexes**
- equilibrium and spectroscopic studies of diethyltin(IV) complexes formed with carbohydrate derivatives of thiazolidine-4-carboxylic acid, **218**, 65
- Carbonato complexes**
- the steric effect on the ring opening process of the decarboxylation of *cis*-carbonato-bis(diamine)cobalt(III) ions, **221**, 9
- Carbon–carbon bond cleavage**
- oxidation of *cis*- and *trans*-pinane with RuO<sub>4</sub> generated in situ, **222**, 115
- Carbon–carbon bond formation**
- electrophilic addition to styrylsilanes: sequential carbon–carbon bond forming reactions, **220**, 145
- Carbon dioxide fixation**
- carbon dioxide fixation by Cp<sub>2</sub>(η<sup>3</sup>-allyl)Ti complexes generated from various dienes, **222**, 145
- Carbon–hydrogen insertion**

- dirhodium(II) tetrakis[*N,N*-dimethyl-2-pyrrolidone-5(*S*)-carboxamide]. Structural effects on enantioselection in metal carbene transformations, **220**, 193
- Carbon monoxide substitution  
ligand substitution at 19-electron organometallic centers. Electrocatalytic CO substitution reactions of (methylcyclopentadienyl)Mn(CO)<sub>2</sub>NO<sup>+</sup> and (indenyl)Mn(CO)<sub>2</sub>NO, **226**, 53
- Carbonylation  
cobalt chloride and phase-transfer catalyzed carbonylation of benzyl chlorides, **216**, 97  
labeling studies of some carbonylation reactions of styrene with cationic palladium complexes, **220**, 273
- Carbonyl clusters  
the formation of carbides in hexaruthenium carbonyl clusters, **227**, 197
- Carbonyl complexes  
electrochemical investigation of the electron-poor/precise ( $n=5/6$ ) complexes [Fe<sub>2</sub>(CO)<sub>*n*</sub>(μ-PR<sub>2</sub>)(μ-PR'<sub>2</sub>)] ( $n=5$ , R=R'=Bu<sup>1</sup>;  $n=6$ , R=R'=Ph; R=Bu<sup>1</sup>, R'=Ph; R=Bu<sup>1</sup>, R'=Cy). EPR study of the radical anion [Fe<sub>2</sub>(μ-PBu<sup>1</sup>)<sub>2</sub>(CO)<sub>5</sub>], **217**, 29  
molybdenum carbonyl derivatives containing eight- and nine-membered diphosphine chelating rings: the synthesis of [Mo(CO)<sub>4</sub>(dppp)<sub>2</sub>] (dppp = Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>3</sub>PPh<sub>2</sub>) and [Mo(CO)<sub>4</sub>(dpph)] (dpph = Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>6</sub>PPh<sub>2</sub>) and the crystal structure of the latter, **218**, 9  
a mechanistic study of the homogeneous catalytic hydroformylation of formaldehyde: synthesis and characterization of model intermediates, **218**, 89  
preparation and crystal structure of *anti*-(η<sup>3</sup>:η<sup>5</sup>-azulene)-pentakis(trimethylphosphine)dicobalt. Reactions with carbon monoxide and solution dynamics, **218**, 143  
coordination chemistry of perhalogenated cyclopentadienes and alkynes XIV. Synthesis of dinuclear ring-bridged cyanotrenes with -SiMe<sub>2</sub>- or -TiCp<sub>2</sub>- bridges. Structures of [(OC)<sub>3</sub>Mn(C<sub>5</sub>Br<sub>4</sub>-)]<sub>2</sub>SiMe<sub>2</sub> and [(OC)<sub>3</sub>Mn(C<sub>5</sub>Cl<sub>4</sub>-)]<sub>2</sub>TiCp<sub>2</sub>, **218**, 173  
organometallic chelating ligands. Synthesis of a new anionic tripod like ligand with an N,N',O-donor set. X-ray crystal structure of [(η<sup>5</sup>-C<sub>6</sub>Me<sub>6</sub>)Ru{μ-P(O)(OMe)<sub>2</sub>(μ-Pz)<sub>2</sub>Re(CO)<sub>3</sub>}] (Me = methyl, Pz = pyrazolate), **219**, 67  
transition metal complexes with sulfur ligands Part CII.  
Changes in configuration and reactivity caused by alkyl substituents R in [Fe(L)('N<sub>R</sub>S<sub>4</sub>') complexes (L = CO, NO<sup>+</sup>, NO, PMe<sub>3</sub>, N<sub>2</sub>H<sub>4</sub>; R = CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub><sup>-</sup>), **219**, 75  
synthesis and X-ray structure of [acenaphthene-chromium(0)-dicarbonyl-tri(2,3,5-trimethylphenyl)phosphite], **219**, 209  
a study of the reactivity of (methyl 2-acetamidoacrylate)-tricarbonyliron(0) leading to a novel synthesis of β,β,β-trialkyl α-amino acids, **220**, 137  
reactivity of allene at phosphine-bridged di-iron centres: X-ray crystal structures of [Fe<sub>2</sub>(CO)<sub>5</sub>{μ-σ,η<sup>3</sup>-C(O)C(CH<sub>2</sub>)<sub>2</sub>}(μ-dppm)] and [Fe<sub>2</sub>(CO)<sub>4</sub>{μ-η<sup>3</sup>,η<sup>3</sup>-(CH<sub>2</sub>)<sub>2</sub>C<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>}(μ-dppm)]·Et<sub>2</sub>O, **220**, 201  
stereoselective aldol addition reactions of Fischer carbene complexes via electronic tuning of the metal center for enolate reactivity, **220**, 215  
a study of the reactivity and structure of cyclic α,β-unsaturated Fischer-type carbene complexes, **220**, 233  
preparation of η<sup>1</sup>-1,3-dienyl, η<sup>1</sup>-1,2-dienyl and η<sup>3</sup>-cyclobutene complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305  
investigation of the dithiocarbamate-induced coupling of allylidene and carbonyl ligands on a tungsten center, **220**, 335  
N<sup>1</sup>,N<sup>7</sup>-expeditious dialkylation of cyclen (1,4,7,10-tetraazacyclododecane). An astonishing reactivity of cyclen tricarbonyl molybdenum and chromium complexes, **220**, 347  
an investigation of electron transfer properties of (Cp)Mn(CO)<sub>3</sub> and substituted Cp derivatives, **221**, 85  
exchange of azine ligands in *fac*-Re(CO)<sub>3</sub>(4-cyanopyridine)<sub>2</sub>Cl and *fac*-Re(CO)<sub>3</sub>(quinoline)<sub>2</sub>Cl, **221**, 161  
synthesis and structure of Cr(CO)<sub>3</sub> complexes of biphenyl compounds with axial chirality, **222**, 63  
conversion of cationic phosphite complexes into phosphonate complexes with selective deamination on the coordinating phosphorus. Reaction of [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Fe{P(NC<sub>4</sub>H<sub>8</sub>)<sub>*n*</sub>-(OMe)<sub>3-*n*</sub>}]<sup>+</sup> (L = CO, PMe<sub>3</sub>) with KOH to afford (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Fe{P(O)(NC<sub>4</sub>H<sub>8</sub>)<sub>*n*-1</sub>(OMe)<sub>3-*n*</sub>}], **222**, 123  
catalytic hydrosilylation of carbon monoxide with cobalt carbonyls, **222**, 131  
selective transformations of alkynols catalyzed by ruthenium complexes, **222**, 155  
steric and electronic tuning of Michael addition reactions induced by pentacarbonyl metal fragments, **222**, 207  
synthesis, structure, dynamics and first atropisomer-selective cleavage of the chromium tricarbonyl complex of a lactone-bridged biaryl, **222**, 255  
arene migration in ruthenium clusters: a kinetic study of the isomerisation of Ru<sub>5</sub>C(CO)<sub>12</sub>(μ<sub>3</sub>-η<sup>2</sup>:η<sup>2</sup>:η<sup>2</sup>-C<sub>6</sub>H<sub>6</sub>) to Ru<sub>5</sub>C(CO)<sub>12</sub>(η<sup>6</sup>-C<sub>6</sub>H<sub>6</sub>), **222**, 299  
transition metal complexes with sulfur ligands CIV. Protonation and alkylation of thiolate donors of [Fe(CO)<sub>2</sub>('S<sub>*i*</sub>')] and [Fe(CO)('S<sub>*i*</sub>')]: syntheses and properties of Fe(II) complexes with open-chain thioether ligands ('S<sub>4</sub><sup>2-</sup> = 1,2-bis(2-mercaptophenylthio)ethane(2-); 'S<sub>5</sub><sup>2-</sup> = 2,2'-bis(2-mercaptophenylthio)diethylsulfide(2-)), **224**, 45  
reactions of HCCO<sub>3</sub>(CO)<sub>9</sub> with silanes; synthesis and electrochemistry of X[SiMe<sub>2</sub>CCO<sub>3</sub>(CO)<sub>9</sub>]<sub>2</sub> (X = O, 1,4-C<sub>6</sub>H<sub>4</sub>), **224**, 73  
di-manganese carbonyl compounds containing bis(dicyclohexylphosphino)methane (dcpm) ligation, **224**, 105  
crystal and molecular structures of [{*exo*- and *endo*-(RO)<sub>2</sub>P(O)-η<sup>5</sup>-C<sub>6</sub>H<sub>6</sub>}Mn(CO)<sub>3</sub>], **224**, 147  
transition-metal silyl complexes 51. Preparation of the mixed silyl/stannyl complexes (π-C<sub>5</sub>H<sub>5</sub>)Fe(CO)(SnPh<sub>3</sub>)(SiR<sub>3</sub>)H, **224**, 177  
cage escape yields for photoinduced bimolecular electron transfer reactions of Re(I) complexes, **225**, 41  
the redox behaviour of the cluster anion [Fe<sub>4</sub>N(CO)<sub>12</sub>]<sup>-</sup>. Electron transfer chain catalytic substitution reactions. Crystal structure of [Fe<sub>4</sub>N(CO)<sub>11</sub>PPh<sub>3</sub>], **226**, 1  
ligand substitution at 19-electron organometallic centers. Electrocatalytic CO substitution reactions of (methylcyclopentadienyl)Mn(CO)<sub>2</sub>NO<sup>+</sup> and (indenyl)Mn(CO)<sub>2</sub>NO, **226**, 53  
photoinduced intramolecular electron transfer in bimetallic donor-acceptor assemblies, **226**, 79  
synthesis and structural analysis of *trans*-Cp(CO)<sub>2</sub>(PPh<sub>3</sub>)WHgCl, **227**, 145  
the equilibrium between acetyl- and methylcobalt tetracarbonyl, **227**, 211  
substitution kinetics of some tetracarbonyl(η<sup>2</sup>-alkene)osmium complexes, **227**, 215  
thermal decomposition of cobalt carbonyl complexes in viscous media, **227**, 233  
kinetics and mechanism of the methanolysis of acetylcobalt tetracarbonyl, **227**, 297
- Carboxamido complexes  
the reduction of pentaammine(phenanthroline-2-carboxamido-*N*)cobalt(III) by titanium(III) and vanadium(II), **225**, 245
- Carboxylate complexes  
preparation and structural characterization of Rh<sub>2</sub>(O<sub>2</sub>CCPh<sub>3</sub>)<sub>4</sub>(EtOH)<sub>2</sub>,



- $\text{Ru}_2(\text{O}_2\text{CCPh}_3)_4(\text{H}_2\text{O})(\text{EtOH}) \cdot 2\text{EtOH}$  and  $\text{Mo}_2(\text{O}_2\text{CCPh}_3)_4 \cdot 3\text{CH}_2\text{Cl}_2$ , **215**, 9
- preparation, structure and catecholase-mimetic activity of two mononuclear ferrocenecarboxylate copper(II) complexes, **215**, 131
- pyrazine-mediated antiferromagnetic intermolecular exchange in mixed-valent diruthenium tetracarboxylates, **215**, 203
- synthesis and characterization of symmetric and unsymmetric oxo-bridged trinuclear chromium benzoate complexes: crystal and molecular structure of  $[\text{Cr}_3\text{O}(\text{O}_2\text{CPh})_6(\text{py})_3]\text{ClO}_4$ , **217**, 171
- preparation, crystal structures and magnetic properties of a *syn-anti* carboxylate-bridged dinuclear copper(II) complex and its precursor, a hydrogen-bonded polynuclear copper(II) complex, **218**, 109
- synthesis and X-ray analysis of the dirhodium(II,II) carboxylate complex  $\text{Rh}_2(\text{O}_2\text{CCH}_3)_4(\text{formH})_2$  (formH = *N,N'*-di-*p*-tolylformamide), **223**, 145
- synthesis and properties of bis(amine)carboxyhydroboron(1+) cations, **223**, 155
- Carboxylato complexes
- crystal structure, magnetic and spectroscopic properties of the bis(dimethyl sulfoxide) adduct of tetra- $\mu$ -formato-dicopper(II), a new tetracarboxylato-bridged copper(II) dimer, **216**, 185
- selective transformations of alkynols catalyzed by ruthenium complexes, **222**, 155
- thermodynamics of complexation of the uranyl(VI) ion with benzoate. The crystal structure of sodium tris(benzoato) dioxouranate dihydrate,  $\text{Na}[(\text{PhCO}_2)_3\text{UO}_2] \cdot 2\text{H}_2\text{O}$ , **223**, 151
- synthesis of some dinuclear niobium chloro alkoxy carboxylates  $[\text{Nb}_2\text{Cl}_4(\text{OEt})_4(\text{O}_2\text{CR})_2]$ , and the X-ray crystal and molecular structure of the compound with  $\text{R} = \text{Ph}$ , **227**, 99
- Carboxylic acid
- selective transformations of alkynols catalyzed by ruthenium complexes, **222**, 155
- Carboxylic acid complexes
- photochemical and spectroscopic studies of complexes of iron(III) with citric acid and other carboxylic acids, **226**, 117
- Catalysis
- cobalt chloride and phase-transfer catalyzed carbonylation of benzyl chlorides, **216**, 97
- arylation of alkenes by the palladium(II) acetate–dialkyl sulfide system, **216**, 253
- a mechanistic study of the homogeneous catalytic hydroformylation of formaldehyde: synthesis and characterization of model intermediates, **218**, 89
- regio- and stereoselective dimerization of 1-alkynes catalyzed by an Os(II) complex, **220**, 5
- ruthenium complex catalyzed polymerization of OH or COOH group containing alkynes to give functionalized poly(acetylene)s, **220**, 35
- rhodium-catalyzed synthesis of 2(5*H*)-furanones from terminal alkynes and non-substituted alkynes under water-gas shift reaction conditions, **220**, 45
- asymmetric catalysis 87\*. Enantioselective allylation of 1,5-dimethylbarbituric acid with Pd catalysts and new optically active PN ligands, **220**, 63
- the importance of phosphine-to-rhodium ratios in enantioselective hydroborations, **220**, 93
- ultrasound-promoted palladium-catalyzed carbonyl allylation by allylic alcohols with tin(II) chloride in non-polar solvents, **220**, 155
- optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220**, 283
- synthesis and structural characterization of a new series of ruthenium(II) complexes containing the short bite  $\text{Ph}_2\text{PPy}$  ligand. Cooperative effect between the anionic rhodium and cationic ruthenium species in the catalytic hydroformylation of styrene by  $[\text{Ru}(\text{Ph}_2\text{PPy})_3\text{Cl}][\text{Rh}(\text{CO})_2\text{Cl}_2]$ , **221**, 109
- strain, silyl and steric effects on the regioselectivity of palladium(0) catalyzed allyl esters reduction as alternative to the Wittig reaction, **222**, 37
- a comparison of catalytic activity for imine hydrogenation using Ru ditertiary phosphine complexes, including chiral systems, **222**, 85
- uranium compounds as catalysts for aldolisation reactions, **222**, 91
- catalytic hydrosilylation of carbon monoxide with cobalt carbonyls, **222**, 131
- selective transformations of alkynols catalyzed by ruthenium complexes, **222**, 155
- intermediates in the palladium-catalysed reaction of 1,3-dienes Part 8. The reaction of palladium–butadiene complexes with ethyl methylacetoacetate, **222**, 179
- hydration of conjugated dienes to produce ketones catalyzed by ruthenium complexes, **222**, 225
- alkyne hydration using Pt(II) catalysts, **222**, 317
- an europium(III) complex as an efficient catalyst for the Michael reaction of  $\alpha,\beta$ -enones with ketene silyl acetals: LIS-NMR analysis for the transition state through complexation between europium(III) catalyst and enones, **222**, 377
- palladium complex catalysed synthesis of sulfinic acids, sulfinic acid esters, sulfonic acids and *S*-alkyl alkanethiosulfonates, **222**, 381
- a one-pot synthesis of cyclopropanes from aromatic aldehydes and styrene via iron carbene complexes, **223**, 159
- dynamical investigations of the catalytic mechanisms of water oxidation by the  $[(\text{bpy})_2\text{Ru}(\text{OH}_2)]_2\text{O}^{4+}$  ion, **226**, 179
- Catalytic activity
- the role of  $\text{H}_2\text{Ru}(\text{CO})_2(\text{PBu}_3)_2$  in the activation of alkynes and alkenes, **227**, 247
- Catecholate complexes
- synthesis and characterization of oxorhenium(V)–catecholate complexes. Crystal and molecular structures of  $(\text{CH}_3)_4\text{N}[\text{ReO}(\text{O}_2\text{C}_6\text{H}_4)_2]$  and  $(\text{CH}_3)_4\text{N}[\text{ReO}(\text{PPh}_3)(\text{O}_2\text{C}_6\text{H}_4)_2] \cdot \text{CH}_3\text{OH}$ , **218**, 157
- Catechol complexes
- valence tautomerism for catechol/semiquinone complexes of the *trans*- $\text{M}(\text{Bupy})_2(3,6\text{-DBQ})_2$  ( $\text{M} = \text{Mn}, \text{Fe}, \text{Co}$ ) series, **226**, 91
- Cationic palladium complexes
- labeling studies of some carbonylation reactions of styrene with cationic palladium complexes, **220**, 273
- CD spectroscopy
- oxidative quenching of luminescence from copper metallothionein, **226**, 275
- Ceria compounds
- electron transfer reactions in non-stoichiometric ceria and urania, **225**, 305
- Cerium complexes
- synthesis and molecular structure of two crystalline forms of hydrated cerium(III) picrate, **221**, 169
- Chalcogenide complexes
- evidence of the  $\text{W}(\text{O})(\text{S}_2)_2$  core as an intermediate in the acidification of  $\text{WS}_4^{2-}$ . Structural characterization of  $\text{W}(\text{O})(\text{S}_2)_2(\text{bpy})$  and  $\text{W}(\text{O})(\text{S}_2)_2(\text{phen})$ , **216**, 201
- extrusion of molecular clusters from solid-state materials: synthesis by application of  $\gamma$ -irradiation. Molecular and crystal structure of  $(\text{H}_6\text{O}_4)(\text{Et}_4\text{N})[\text{Mo}_3\text{S}_7\text{Br}_6]$ , **223**, 5
- Chalcogeno complexes

- the anomalous EPR spectra of the iron(III) dichalcogenocarbamates, **217**, 101
- Charge transfer  
charge and electron transfer from metal-to-carbon bonds of main group organometallics  $MR_n$  ( $M = Al, Ga, Zn$ ) to aromatic N-heterocycles: colored precursor compounds and radical complex formation, **225**, 23
- Charge transfer excitation  
photoredox behavior of a trinuclear Ru(III),Ru(II),Co(III) complex induced by metal-to-metal charge transfer excitations, **225**, 327
- Charge transfer excited state  
synthesis, electrochemistry and photoexcited-state properties of dinuclear ruthenium complexes bridged by 2,6'-bis(2-pyridyl)-2,2':6,2''-thiazolo[4,5-*d*]-benzothiazole, **226**, 17
- Chelate complexes  
neutral transition metal complexes with the tridentate ligand 4-(5-methyl-4-imidazolyl)-3-thiabutanoic acid. A series of isostructural octahedral compounds, **215**, 151  
mixed-ligand complexes of rhenium II. Synthesis and characterization of  $[ReN(X)(Me_2PhP)_2(R_2tcb)]$  complexes ( $X = I, N_3, SCN, CN$ ;  $R_2tcb^- = N-(N,N\text{-dialkylthiocarbamoylbenzamidinate})$ ). X-ray crystal structure of  $[ReN(N_3)(Me_2PhP)_2(Et_2tcb)]$ , **215**, 159  
solvent effects on acetylacetonato iron complexes, **215**, 185  
principal component analysis of the structures of the 1,2-diaminobenzene derivatives chelated to metal ions: correlation between structural parameters and redox behavior, **215**, 219  
transition metal compounds of the tridentate pyrazole substituted amine ligand bis(2-(3,5-dimethyl-1-pyrazolyl)-ethyl)ethylamine (ddae). X-ray structures of  $[Co(ddae)(NO_3)_2]$ ,  $[Cu(ddae)(NO_3)(H_2O)](NO_3)$  and  $[Cu(ddae)(Cl)_2] \cdot C_2H_5OH$ , **216**, 43  
unexpected five-coordination in di(1,3-diaminopropane)-3,5-dinitrobenzoatocopper(II) 3,5-dinitrobenzoate. Comparison between the coordinated and non-coordinated anions, **216**, 57  
spectroscopic and structural studies on volatile gallium  $\beta$ -diketonates as potential precursors for MOCVD, **217**, 71  
unusually stable peroxocopper complexes. Stoichiometry, products and kinetics of oxidation of the dimeric copper(I) complex  $[LCuBr]_2$  ( $L = N,N'$ -diethylethylenediamine) by dioxygen in methylene chloride from  $-51$  to  $30^\circ C$ , **217**, 109  
nucleophilic addition of water to coordinated di-(2-pyridyl)ketone (DPK) in rhenium(V) complexes. Synthesis and crystal structure of the adduct  $(DPK \cdot H)^+ \{ [ReOCl_2(DPK \cdot OH)]_2 Cl \}$ , **217**, 141  
complexes of mercuric halides with the tripod ligand  $np_3$  ( $= N(CH_2CH_2PPh_2)_3$ ). Molecular structure of  $\{ [np_3]HgI \} \cdot 0.5C_2H_5OH$ , **217**, 155  
the synthesis and crystal structure of  $[Cr(acacen)py_2][ZnCl_3py]$ , **217**, 195  
molybdenum carbonyl derivatives containing eight- and nine-membered diphosphine chelating rings: the synthesis of  $[Mo(CO)_4(dppp)_e]$  ( $dppp_e = Ph_2P(CH_2)_3PPh_2$ ) and  $[Mo(CO)_4(dpph)]$  ( $dpph = Ph_2P(CH_2)_6PPh_2$ ) and the crystal structure of the latter, **218**, 9  
new heptacoordinated tin complexes of 2,6-diacetylpyridine-bis(thiosemicarbazone),  $H_2daptsc$ , and of 2,6-diacetylpyridine-bis(semicarbazone),  $H_2dapsc$ . Crystal and molecular structures of  $[MeSnCl(Hdaptsc)]Cl \cdot MeOH$  and  $[MeSnCl(H_2dapsc)]Cl_2 \cdot 2H_2O$ , **218**, 139  
conformational flexibility of 1,3-diaminopropane in chelate rings. Redetermination of the structure of *trans*-dinitratobis(1,3-diaminopropane)copper(II), **218**, 151  
synthesis, crystal structure and spectroscopic properties of bis(diphenyldithiophosphinato)cadmium(II), **219**, 115  
synthesis and characterization of hexadentate cobalt(III) complexes with new edta-type ligands Part 4. Circular dichroism of the *trans*-( $O_5O_6$ ) isomer of a cobalt(III) complex of 1,3-propanediamine- $N,N,N'$ -triacetic- $N'$ -3-propionic acid, **219**, 147  
three-coordinate Pt(0)  $\eta^2$ -complexes: electrophilic hydrogen attack through oxidative-addition of protic acids, **219**, 169  
synthesis and X-ray crystal structure of *cis*-1,4-diaminocyclohexanetetrachloroplatinum(IV): a new antitumor agent, **219**, 193  
spectroscopic investigation of mixed-ligand nickel(II) chelates containing  $\beta$ -ketoenols and 1,2-diamines, **221**, 17  
an ethyl-cobalt complex with a buckled equatorial ligand, **221**, 35  
synthesis, spectral and redox properties of a new series of aqua complexes of ruthenium(II), **221**, 125  
metal complexes of 5-acetyl-2,2'-pyridylthiophene aroylhydrazones. Crystal structure of a cobalt(II) derivative, **223**, 97  
synthesis and characterization of metal complexes with  $N,N''$ -bis(2-hydroxybenzyl)-diethylenetriamine- $N,N',N''$ -triacetic acid and the stabilities of its complexes with divalent and trivalent metal ions, **223**, 109  
cage escape yields for photoinduced bimolecular electron transfer reactions of Re(I) complexes, **225**, 41  
electrochemical, photophysical and photochemical properties of a hexanuclear ruthenium(II) bis(2,2'-bipyridyl) complex containing a tris(pyrazyltriazole)benzene ligand, **225**, 67  
electron transfer 122. Reductions of an  $N_4$ -coordinated complex of silver(III), **225**, 83  
electron transfer properties of ReO(ECD), **225**, 323  
synthesis and characterization of a bidentate chelating diisonitrile ligand with two thioetheral sulfurs in a bridging group and its copper(I), rhodium(I) and palladium(II) complexes, **227**, 113  
preparation, structural and physical studies of mercury(II) halide complexes with the  $N,N'$ -chelating biheteroaromatic ligand 2-(2'-pyridyl)quinoxaline, **227**, 129
- Chelate ligand complexes  
structure and spectroscopic properties of  $[di(3\text{-aminopropyl})amine-N,N',N''] [iminodiacetato-N,O,O']chromium(III) perchlorate$ , **216**, 155  
the oxidation of  $[Co(edta)]^{2-}$  by  $[Co(phen)_3]$ , **225**, 75
- Chelate phosphine ligand complexes  
the crystal structure of a chelate silver(I) complex:  $\{ Ag[Ph_2P(CH_2)_2SCH_2CH_3]_2 ClO_4 \}$ , **215**, 199
- Chelating ligand complexes  
crystal and molecular structure of the solvate of bis(2,6-bis(1'-methylbenzimidazol-2'-yl)pyridine)cobalt(II) dihexafluorophosphate with 4-methyl-1,3-dioxolan-2-on, **219**, 11  
the first isolated antineoplastic Ru(IV) complex: synthesis and structure of  $[Cl_2(1,2\text{-cyclohexanediaminotetraacetate})Ru] \cdot 2H_2O$ , **224**, 15
- Chemistry  
chemical and electrochemical approaches to the investigation of redox reactions of simple electron transfer metalloproteins, **226**, 293
- Chemoselective reaction  
chemoselective functionalization of zirconacyclopentenes, **220**, 319
- Chemoselectivity  
chele- and anticheleselectivity IV. Reaction-accelerating neighboring-group effects in the cross-coupling of vinylbromides with alkyl-transition metal reagents, **220**, 85
- Chiral complexes

- EXAFS studies of a chiral rhenium lactone complex and its precursors including multiple scattering calculations, **222**, 27
- Chiral diamines  
 stereochemical investigation of cationic olefin complexes of platinum(II) containing chiral diamines: a sequence of four asymmetric centres (N, C, C and N) in the chelate ring, **227**, 261
- Chiral ligand complexes  
 chiral metal complexes 40. The sterically mediated hydrolysis of a coordinated imine, **221**, 191  
 synthesis and structure of new chiral ferrocenylphosphines for asymmetric catalysis, **222**, 213
- Chiral reagents  
 the preparation of chiral titanium reagents: a comparison of new and known ligands, **222**, 95
- Chloride complexes  
 Raman and infrared spectroscopic studies of tris triphenyl phosphine chloride complexes of transition metals, **216**, 191  
 spectroscopic studies of the *trans*-[VOCl<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>] chromophore in its hydrogen bonding adduct with 18-crown-6, **227**, 91
- Chloroaluminate ionic ligands  
 dependence of electron transfer kinetics of the ferrocene/ferrocenium couple on the viscosity in ambient temperature chloroaluminate ionic liquids, **225**, 9
- Chloro complexes  
 tTbPcCrCl<sub>2</sub>H a sparingly soluble tetra-*t*-butyl-phthalocyaninato-chloro-chromium complex, **224**, 185
- Chlorophyll complexes  
 ESR studies on distortion in the macrocyclic ring of copper(II) chlorophylls, **224**, 181
- Chromium complexes  
 kinetics and mechanism of the phosphate-catalyzed heterolysis of the metal-carbon  $\sigma$  bond of  $\alpha$ -hydroxy-alkyl-chromium(III) complexes. Effects of the non-participating ligands, temperature and pressure, **215**, 173  
 a polymeric two-dimensional mixed-metal network. Crystal structure and magnetic properties of {[P(Ph)<sub>4</sub>]-[MnCr(ox)<sub>3</sub>]}<sub>n</sub>, **216**, 65  
 interaction of cyanopentaamminechromium(III) with mercuric ion. Electrochemical, spectral and preparative study of adduct formation and linkage isomerism, **216**, 113  
 structure and spectroscopic properties of [di(3-aminopropyl)-amine-*N,N',N''*][iminodiacetato-*N,O,O'*]chromium(III) perchlorate, **216**, 155  
 synthesis and characterization of symmetric and unsymmetric oxo-bridged trinuclear chromium benzoate complexes: crystal and molecular structure of [Cr<sub>3</sub>O(O<sub>2</sub>CPh)<sub>6</sub>(py)<sub>3</sub>]ClO<sub>4</sub>, **217**, 171  
 the synthesis and crystal structure of [Cr(acacen)py<sub>2</sub>][ZnCl<sub>3</sub>py], **217**, 195  
 structural characterization of octahedral metal complexes of [MX<sub>3</sub>(py)<sub>3</sub>] type with three pyridines as revealed by <sup>2</sup>H NMR spectra, **218**, 179  
 synthesis and X-ray structure of [acenaphthene-chromium(0)-dicarbonyl-tri(2,3,5-trimethylphenyl)phosphite], **219**, 209  
 reactions of benzocyclobutene chromium complexes with carbon, nitrogen and oxygen nucleophiles: nucleophilic addition and unexpected proximal ring opening reactions, **220**, 175  
 stereoselective aldol addition reactions of Fischer carbene complexes via electronic tuning of the metal center for enolate reactivity, **220**, 215  
 a study of the reactivity and structure of cyclic  $\alpha,\beta$ -unsaturated Fischer-type carbene complexes, **220**, 233  
 N<sup>1</sup>,N<sup>7</sup>-expeditious dialkylation of cyclen (1,4,7,10-tetraazacyclododecane). An astonishing reactivity of cyclen tricarbonyl molybdenum and chromium complexes, **220**, 347  
 crystal structure and ligand field states of [Cr(NH<sub>3</sub>)<sub>5</sub>NCO](NO<sub>3</sub>)<sub>2</sub>, **221**, 93  
 synthesis and structure of Cr(CO)<sub>3</sub> complexes of biphenyl compounds with axial chirality, **222**, 63  
 synthesis, structure, dynamics and first atropisomer-selective cleavage of the chromium tricarbonyl complex of a lactone-bridged biaryl, **222**, 255  
 synthesis of  $\alpha$ -hydroxy acid derivatives by the photochemical reaction of *O*-silylchromium(0) carbene complexes and nucleophiles, **222**, 261  
 tTbPcCrCl<sub>2</sub>H a sparingly soluble tetra-*t*-butyl-phthalocyaninato-chloro-chromium complex, **224**, 185  
 the synthesis, X-ray structure and hydrolysis kinetics of chloro(2-aminomethylpyridine)(triamine)chromium(III) salts, **227**, 11  
 volumes of activation for  $\beta$ -elimination and related reactions of chromium(III)-alkyl complexes in aqueous solutions, **227**, 57  
 the preparation, properties and crystal structure of a salt containing the hexabromochromate(III) anion, **227**, 167
- Chromophore complexes  
 ruthenium chromophores containing terpyridine and a series of polyazine bridging ligands, **221**, 55
- Cinoxacin complexes  
 synthesis and characterization of Cd(II) complexes of cinoxacin. Crystal structure of a dicadmium complex containing two monoatomic-bridging carboxylate oxygen atoms, **217**, 149
- Cluster complexes  
 two silicon-bridged cyclopentadienyl Fe<sub>2</sub>S<sub>4</sub> cluster units connected through an Fe(III)S<sub>6</sub> unit. Crystal and molecular structure of [((CH<sub>3</sub>)<sub>2</sub>SiCp<sub>2</sub>)<sub>2</sub>Fe<sub>3</sub>( $\mu_3$ -S<sub>2</sub>)<sub>2</sub>( $\mu_4$ -S<sub>2</sub>)( $\mu_4$ -S<sub>4</sub>)]FeCl<sub>4</sub>, **216**, 1  
 oxo- and sulfido-bridged sandwich cubane-type molybdenum-indium mixed-metal cluster. Synthesis and X-ray structure of [(H<sub>2</sub>O)<sub>9</sub>Mo<sub>3</sub>S<sub>3</sub>OInOs<sub>3</sub>Mo<sub>3</sub>(H<sub>2</sub>O)<sub>9</sub>]- (CH<sub>3</sub>·C<sub>6</sub>H<sub>4</sub>·SO<sub>3</sub>)<sub>8</sub>·30H<sub>2</sub>O, **216**, 13  
 a novel tetranuclear iodoargentate(I) ion, [Ag<sub>4</sub>I<sub>8</sub>]<sup>4-</sup>, with a cubane-type core, **217**, 15  
 systematics of some properties of hexamolybdotellurate(VI) salt hydrates with di- and trivalent metals, M<sub>4</sub>[TeMo<sub>6</sub>O<sub>24</sub>]·nH<sub>2</sub>O, **217**, 93  
 mercury and platinum mixed cluster having a prismatic structure, [HgPt<sub>6</sub>(2,6-Me<sub>2</sub>C<sub>6</sub>H<sub>3</sub>NC)<sub>12</sub>], **217**, 121  
 synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129  
 ten-vertex polyhedral mixed diheteromonometallaborane chemistry. Directed synthesis of *arachno*-9,6,8-platinathiacarbadecaboranes, **218**, 1  
 the study of isolobal displacement reactions. Synthesis and structures of the heterobimetallic  $\mu$ -alkyne cluster complexes  $\eta^5$ -RC<sub>5</sub>H<sub>4</sub>(CO)<sub>2</sub>M( $\mu$ -R<sup>1</sup>C<sub>2</sub>R<sup>2</sup>)Co(CO)<sub>3</sub> (M = Mo, W; R = CH<sub>3</sub>C(O), CH<sub>3</sub>O<sub>2</sub>C), **219**, 93  
 transition metal complexes with sulfur ligands CIII. [( $\mu$ -S<sub>2</sub>)-{Fe( $\mu$ -S<sub>4</sub>)<sub>2</sub>}]<sub>2</sub>, a novel [2Fe-2S] cluster with octahedral low-spin [Fe<sup>II</sup>S<sub>6</sub>] centers ('S<sub>4</sub>'<sup>2-</sup> = 1,2-bis(2-mercaptophenylthio)ethane(2-)), **224**, 35  
 electron transfer reactivity of cuboidal heterometallic Mo<sub>3</sub>MS<sub>4</sub> clusters in aqueous acidic solutions, **225**, 157  
 syntheses and structures of mixed-ligand tetrairon-sulfur clusters, [Cp'<sub>2</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)<sub>2</sub>Fe<sub>4</sub>S<sub>4</sub>](PF<sub>6</sub>) and [Cp'<sub>3</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)Fe<sub>4</sub>S<sub>5</sub>](PF<sub>6</sub>)<sub>n</sub> (n = 1 and 2) (Cp' =  $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>). Structural change of the clusters accompanying electron transfer reactions, **225**, 229

- electron transfer 120. Reductions of the cuboidal mixed-valence molybdenum/sulfur cluster,  $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]$ , **225**, 297
- the redox behaviour of the cluster anion  $[\text{Fe}_4\text{N}(\text{CO})_{12}]^-$ . Electron transfer chain catalytic substitution reactions. Crystal structure of  $[\text{Fe}_4\text{N}(\text{CO})_{11}(\text{PPh}_3)]$ , **226**, 1
- redox behavior of the iron–sulfur cluster  $[\text{Fe}_4\text{Cp}_4\text{S}_5][\text{PF}_6]_2$  in protic organic solvents and aqueous micellar solutions, **226**, 341
- heteronuclear metal cluster complexes with a new coordination mode of the  $\text{Fe}_2\text{S}_2(\text{CO})_6$  unit. Synthesis, structure and possible reaction pathway of the first Fe–Cu–S cluster compound,  $[\text{Fe}_6\text{Cu}_5\text{S}_6(\text{CO})_{18}(\text{PPh}_3)_2]$ , **227**, 119
- a new type of face-sharing bioctahedral triply bonded dinitium cluster  $(\text{PyH})[\text{Nb}_2(\mu\text{-Cl})_2(\mu\text{-THT})\text{Cl}_3(\text{Py})_3](\text{THF})(\text{C}_6\text{H}_6)_{0.5}$ , made by a designed synthesis, **227**, 191
- metal cluster topology 16. Macropolygons and macropolyhedra in coinage metal iron carbonyl cluster anions and related compounds, **227**, 207
- Clusters**
- extrusion of molecular clusters from solid-state materials: synthesis by application of  $\gamma$ -irradiation. Molecular and crystal structure of  $(\text{H}_9\text{O}_4)(\text{Et}_4\text{N})[\text{Mo}_3\text{S}_7\text{Br}_6]$ , **223**, 5
- the vibrations of bridging carbonyl groups as more sensitive indicators of molecular geometry than terminal carbonyl groups in transition metal carbonyl cluster compounds, **227**, 241
- Cobaloxime complexes**
- evidence of the interaction between steric and electronic influence in rhodoximes and cobaloximes. Synthesis of  $\text{pyRh}(\text{DH})_2\text{I}$  and X-ray structure of  $\text{pyRh}(\text{DH})_2\text{Cl}$ ,  $\text{pyCo}(\text{DH})_2\text{Cl}$  and  $\text{pyRh}(\text{DH})_2\text{I}$ , **216**, 125
- Cobalt**
- solution structures of cobalt–rhodium clusters: the identification of a new hexanuclear cobalt–rhodium species, **227**, 187
- Cobalt complexes**
- neutral transition metal complexes with the tridentate ligand 4-(5-methyl-4-imidazolyl)-3-thiabutanoic acid. A series of isostructural octahedral compounds, **215**, 151
- cobalt chloride and phase-transfer catalyzed carbonylation of benzyl chlorides, **216**, 97
- evidence of the interaction between steric and electronic influence in rhodoximes and cobaloximes. Synthesis of  $\text{pyRh}(\text{DH})_2\text{I}$  and X-ray structure of  $\text{pyRh}(\text{DH})_2\text{Cl}$ ,  $\text{pyCo}(\text{DH})_2\text{Cl}$  and  $\text{pyRh}(\text{DH})_2\text{I}$ , **216**, 125
- cobalt(III) complexes with 2-acetyl- and 2-formylpyridine  $^4N$ -methylthiosemicarbazones, **216**, 229
- a novel amino acid-binding receptor based on weak non-covalent ligand–ligand interactions, **216**, 249
- preparation and dioxygen binding property of a new zeolite encapsulated cobalt tetramethylporphyrin complex, **217**, 135
- nitrogen-15 NMR coordination shifts of  $\beta$ -alanine and glycine in cobalt(III) complexes, **217**, 213
- stereoselectivity in reactions of metal complexes Part XVIII. Kinetics and stereoselectivity in electron-transfer reactions between  $[\text{Co}(\text{L})\text{H}_2\text{O}]^+$  ( $\text{L} = N, N'$ -[(pyridine-2,6-diyl)bis(methylene)]bis[proline] (promp) and  $N, N'$ -[(4-methoxy-pyridine-2,6-diyl)bis(methylene)]bis[proline] (MeO-promp) and optically active iron(II) complexes, **218**, 73
- preparation and crystal structure of *anti*-( $\eta^3$ : $\eta^5$ -azulene)-pentakis(trimethylphosphine)dicobalt. Reactions with carbon monoxide and solution dynamics, **218**, 143
- crystal and molecular structure of the solvate of bis(2,6-bis(1'-methylbenzimidazol-2'-yl)pyridine)cobalt(II) dihexafluorophosphate with 4-methyl-1,3-dioxolan-2-on, **219**, 11
- the study of isolobal displacement reactions. Synthesis and structures of the heterobimetallic  $\mu$ -alkyne cluster complexes  $\eta^5\text{-RC}_5\text{H}_4(\text{CO})_2\text{M}(\mu\text{-R}^1\text{C}_2\text{R}^2)\text{Co}(\text{CO})_3$  ( $\text{M} = \text{Mo}, \text{W}$ ;  $\text{R} = \text{CH}_3\text{C}(\text{O}), \text{CH}_3\text{O}_2\text{C}$ ), **219**, 93
- synthesis and characterization of hexadentate cobalt(III) complexes with new edta-type ligands Part 4. Circular dichroism of the *trans*-( $\text{O}_5\text{O}_6$ ) isomer of a cobalt(III) complex of 1,3-propanediamine-*N, N, N'*-triacetic-*N'*-3-propionic acid, **219**, 147
- the steric effect on the ring opening process of the decarboxylation of *cis*-carbonato-bis(diamine)cobalt(III) ions, **221**, 9
- an ethyl–cobalt complex with a buckled equatorial ligand, **221**, 35
- chiral metal complexes 40. The sterically mediated hydrolysis of a coordinated imine, **221**, 191
- cyclodimerisation and addition reactions of  $\text{Bu}^i\text{C}\equiv\text{P}$  at a cobalt(I) centre, **222**, 13
- catalytic hydrosilylation of carbon monoxide with cobalt carbonyls, **222**, 131
- the development of a quinone  $\alpha$ -anion synthon. Utilization of a maleoylcobalt complex as a quinone surrogate and a dominant role for ligand effects, **222**, 235
- metal complexes of 5-acetyl-2,2'-pyridylthiophene arylhydrazones. Crystal structure of a cobalt(II) derivative, **223**, 97
- synthesis, properties and molecular structure of copper(II) and cobalt(II) 1,2-bis(pyrazol-1-ylmethyl)benzene complexes, **223**, 139
- reactions of  $\text{HCCO}_3(\text{CO})_9$  with silanes; synthesis and electrochemistry of  $\text{X}[\text{SiMe}_2\text{CCO}_3(\text{CO})_9]_2$  ( $\text{X} = \text{O}, 1,4\text{-C}_6\text{H}_4$ ), **224**, 73
- the oxidation of  $[\text{Co}(\text{edta})]^{2-}$  by  $[\text{Co}(\text{phen})_3]$ , **225**, 75
- volume profile analysis for intermolecular electron transfer between cytochrome *c* and  $\text{Co}(\text{terpy})_2$ , **225**, 95
- intramolecular electron transfer from  $\text{Ru}^{\text{II}}(\text{EDTA})^{2-}$  to  $\text{Co}^{\text{III}}(\text{NH}_3)_5^{3+}$  via nitrogen heterocycle bridging ligands, **225**, 129
- electrocatalysts for the four-electron reduction of dioxygen based on adsorbed cobalt tetrapyrrolylporphyrin molecules linked by aquaammine complexes of ruthenium(II), **225**, 215
- the reduction of pentaammine(phenanthroline-2-carboxamido-*N*)cobalt(III) by titanium(III) and vanadium(II), **225**, 245
- photoredox behavior of a trinuclear  $\text{Ru}(\text{III}), \text{Ru}(\text{II}), \text{Co}(\text{III})$  complex induced by metal-to-metal charge transfer excitations, **225**, 327
- valence tautomerism for catechol/semiquinone complexes of the *trans*- $\text{M}(\text{Bupy})_2(3,6\text{-DBQ})_2$  ( $\text{M} = \text{Mn}, \text{Fe}, \text{Co}$ ) series, **226**, 91
- forbidden  $\pi$ - $\sigma$  donor–acceptor electronic coupling in linked transition metal complexes: contrasting behavior of ruthenium(II) donors with ruthenium(III) and cobalt(III) acceptors, **226**, 109
- electron transfer reduction of cobalt tetraphenylporphyrin by hydroquinone dianions and alkylation with alkyl halides, **226**, 145
- reduction of hexaazacyclophane complexes of  $\text{Co}(\text{II})$  and  $\text{Cu}(\text{II})$  by  $^-\text{CH}_2\text{OH}$  and  $e_{\text{sol}}^-$ ; a pulse radiolysis study, **226**, 151
- electron transfer quenching of the excited MLCT state of  $\text{Ru}(\text{II})$  in the double complex salts  $[\text{Ru}(\text{bpy})_3]_2[\text{M}(\text{CN})_6]\text{Cl}\cdot 8\text{H}_2\text{O}$  ( $\text{M} = \text{Co}$  and  $\text{Fe}$ ), **226**, 203
- spectroscopy and photoredox chemistry of ion-paired cobalt(III) am(m)ine complexes, **226**, 285
- equilibrium studies of 1,2-diaminobenzene and 3,4-diaminobenzoate anion with dinuclear cobalt(II) 1,4,7,13,16,19-hexaaza-10,22-dioxacyclotetrasolane, **227**, 49
- early/late bimetallic complexes:  $\text{MCO}_2$  ( $\text{M} = \text{Ti}, \text{Zr}$ ) derivatives, **227**, 201
- the equilibrium between acetyl- and methylcobalt tetracarbonyl, **227**, 211

- thermal decomposition of cobalt carbonyl complexes in viscous media, **227**, 233
- kinetics and mechanism of the methanolysis of acetylcobalt tetracarbonyl, **227**, 297
- Compartmental ligand complexes  
preparation, properties and coordination behaviour of planar or tridimensional compartmental Schiff bases, **219**, 135
- Condensation reactions  
a novel reaction of the cyano ligand opening an organometallic route to 4-amino imidazoles: four component condensation (4CC) with isocyanide, aldehyde and amine, **222**, 275
- Conjugated dienes  
synthesis of  $\eta^3$ -allyl-Pd complexes and subsequent formation of N-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- Contrast agent complexes  
stability constants of the lanthanide(III)-1,4,7,10-tetraazacyclododecane-*N,N',N'',N'''*-tetraacetate complexes, **221**, 165
- Cooperativity  
a spectroelectrochemical method for evaluating factors which regulate the redox potential of hemoglobins, **226**, 187
- Coordination compounds  
a unified view of inorganic stereochemistry and stereochemical changes, **219**, 187
- Coordination polymer  
spontaneous formation of a 1D-coordination polymer with helix structure starting from (*S*)-pyroglutamic acid and copper(II) acetate, **222**, 5
- Coordination polymer complexes  
predicting the effect of oxidative doping on the conductivities of metal-tetrakis(dimethylphosphino)benzene coordination polymers from the electrochemical properties of their related bimetallic complexes, **225**, 15
- Cope rearrangement  
bicyclic carbo- and heterocycles from an allylidene complex and cyclic 1,3-dienes by tandem cyclopropanation/Cope rearrangement, **220**, 327
- Copper  
EXAFS investigation on nickel substituted Cu,Zn bovine superoxide dismutase, **216**, 121  
stereoselective electron transfer between Cu(II) superoxide dismutase and  $\Lambda$ - and  $\Delta$ -[Fe(pdta)], **225**, 151
- Copper complexes  
towards a model for galactose oxidase: the crystal structure of {2-[bis(2-pyridylethyl)aminomethyl]phenolato} copper(II) perchlorate, **215**, 1  
synthesis and hydrolysis of an ester of the 'one ring open' hydrolysis product of the anticancer drug 1,2-bis(3,5-dioxopiperazin-1-yl)propane, 'Razoxane', **215**, 55  
mixed-ligand copper(II) complexes with iminodiacetate and 2- or 4-methylimidazole: molecular and crystal structure of the 'remote' isomer iminodiacetato(5-methylimidazole)copper(II) monohydrate, [Cu(IDA)(5MeImH)]·H<sub>2</sub>O, **215**, 73  
spectroscopic studies on copper(II) complexes of chiral cyclens: [CuN<sub>4</sub>Cl] chromophores varying from square pyramidal to trigonal bipyramidal stereochemistry, **215**, 109  
preparation, structure and catecholase-mimetic activity of two mononuclear ferrocenecarboxylate copper(II) complexes, **215**, 131  
transition metal compounds of the tridentate pyrazole substituted amine ligand bis(2-(3,5-dimethyl-1-pyrazolyl)ethyl)ethylamine (ddae). X-ray structures of [Co(ddae)(NO<sub>3</sub>)<sub>2</sub>], [Cu(ddae)(NO<sub>3</sub>)(H<sub>2</sub>O)](NO<sub>3</sub>) and [Cu(ddae)(Cl)<sub>2</sub>]·C<sub>2</sub>H<sub>5</sub>OH, **216**, 43  
unexpected five-coordination in di(1,3-diaminopropane)-3,5-dinitrobenzoatocopper(II) 3,5-dinitrobenzoate. Comparison between the coordinated and non-coordinated anions, **216**, 57
- crystal structure, magnetic and spectroscopic properties of the bis(dimethyl sulfoxide) adduct of tetra- $\mu$ -formato-dicopper(II), a new tetracarboxylato-bridged copper(II) dimer, **216**, 185  
synthesis and complexation behaviour of a double-armed crown thioether ligand, **217**, 1  
a seven-spin cluster formed by an alkyl nitronyl nitroxide biradical and copper(II): crystal structure and magnetic properties, **217**, 7  
interaction of the lysine side chain amino group with Cu<sup>II</sup> in (glycyl-L-lysine<sup>2-</sup>)Cu, **217**, 33  
magnetic properties and crystal structure of { $\mu$ -pyrazine}-bis[(aqua)(2,6-xylylamino)diacetato]copper(II)}. A pyrazine-bridged dinuclear copper(II) complex, **217**, 39  
transmetalation of targets ( $\mu$ -Y)N<sub>4</sub>Cu<sub>2</sub><sup>I</sup>Cu<sub>2</sub><sup>II</sup>Cl<sub>4</sub> and ( $\mu$ -Y, $\mu$ -O)N<sub>4</sub>Cu<sub>4</sub>Cl<sub>4</sub> (Y = 3,4,5,6-tetrachlorocatechol) with M(NS)<sub>2</sub> reagents, **217**, 85  
unusually stable peroxocopper complexes. Stoichiometry, products and kinetics of oxidation of the dimeric copper(I) complex [LCuBr]<sub>2</sub> (L = *N,N'*-diethylethylenediamine) by dioxygen in methylene chloride from -51 to 30 °C, **217**, 109  
functionalised phosphines: structure of bis(2-aminophenyl)di-phenylphosphine- $\kappa^2$ *N,P*copper(I) perchlorate ethanol solvate, **217**, 191  
substituent effect on the coordination ability of the amide group of N-protected amino acids, **218**, 53  
synthesis, structural characterization and dioxygen reactivity of imidazole-ligated Cu(I) complexes, **218**, 103  
preparation, crystal structures and magnetic properties of a *syn-anti* carboxylate-bridged dinuclear copper(II) complex and its precursor, a hydrogen-bonded polynuclear copper(II) complex, **218**, 109  
conformational flexibility of 1,3-diaminopropane in chelate rings. Redetermination of the structure of *trans*-dinitratobis(1,3-diaminopropane)copper(II), **218**, 151  
antiferromagnetism tuned by the topology of the intervening bridging ligand. Crystal structure and magnetic properties of the ( $\mu$ -phthalato){[bis(piperidine)]Cu(II)} chain complex, **218**, 167  
reactions of the polynuclear chain complex [Cu(L)(H<sub>2</sub>O)]<sub>n</sub> (H<sub>2</sub>L is 5-[(2'-carboxyphenyl)azo]-1,3-dimethylbarbituric acid) with imidazole. Crystal structures of the dinuclear and mononuclear derivatives and EPR study of the dimer-monomer equilibrium in solution, **219**, 127  
variation of the exchange interaction in oximate-bridged Cu<sup>II</sup>M<sup>II</sup> dimers (M = Cu, Ni, Mn). Crystal structure of [Cu(pdmg)Cu(bipy)(H<sub>2</sub>O)<sub>2</sub>](ClO<sub>4</sub>)<sub>2</sub>·H<sub>2</sub>O, **219**, 179  
synthesis of symmetric and unsymmetric 1,4-bis(*p*-R-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289  
formation of an anchored copper(II) complex on a thiourea-functionalized silica gel by *in situ* modification of the tethered ligating function, **221**, 183  
spontaneous formation of a 1D-coordination polymer with helix structure starting from (*S*)-pyroglutamic acid and copper(II) acetate, **222**, 5  
synthesis, spectroscopic and structural characterization of methyl pyruvate- and pyridoxal-hydrazinopyruvoylthiosemicarbazones, **223**, 77  
synthesis, properties and molecular structure of copper(II) and cobalt(II) 1,2-bis(pyrazol-1-ylmethyl)benzene complexes, **223**, 139

- synthesis, spectra and electrochemical behaviour of biomimetic copper(II) complexes with  $\text{CuN}_5$  and  $\text{CuN}_6$  chromophores, **224**, 19
- crystal structure and magnetic property of a 2D infinite layered copper(II) complex with bridging *trans*-oxamidate and pyrimidine ligands, **224**, 171
- ESR studies on distortion in the macrocyclic ring of copper(II) chlorophylls, **224**, 181
- photophysical properties of mono-, di- and tetranuclear copper(I)–polypyridine complexes, **225**, 251
- photoinduced electron transfer reaction between  $[\text{Cu}(\text{dmp})\text{P}_2]^{+\text{or}-}$  ( $\text{P} = \text{PPh}_3$  or  $\text{PPh}_2(m\text{-C}_6\text{H}_4\text{SO}_3^-)$ ) and viologen derivatives, **225**, 261
- reduction of hexaazacyclophane complexes of Co(II) and Cu(II) by  $^-\text{CH}_2\text{OH}$  and  $e_{\text{sol}}^-$ ; a pulse radiolysis study, **226**, 151
- electron transfer in surfactant films on electrodes; copper phthalocyaninetetrasulfonate–didodecyldimethylammonium bromide, **226**, 159
- synthesis and crystal structure of the  $\mu_2$ -{tris[bis-(diphenylphosphine)methane]}- $\mu_3$ -3-methyl-8-ethyl-xanthine-*N,O,O*}- $\mu_3$ -chloro-triangle-tricopper(I)-water complex exhibiting a novel oxopurine coordination mode, **227**, 43
- structural, spectroscopic and potentiometric investigation on the coordination properties of 4- $\text{NO}_2$ -hippuric acid towards Cu(II) in binary and ternary systems, **227**, 105
- synthesis and characterization of a bidentate chelating diisonitrile ligand with two thioetheral sulfurs in a bridging group and its copper(I), rhodium(I) and palladium(II) complexes, **227**, 113
- heteronuclear metal cluster complexes with a new coordination mode of the  $\text{Fe}_2\text{S}_2(\text{CO})_6$  unit. Synthesis, structure and possible reaction pathway of the first Fe–Cu–S cluster compound,  $[\text{Fe}_6\text{Cu}_5\text{S}_6(\text{CO})_{18}(\text{PPh}_3)_2]$ , **227**, 119
- NQR studies of polyhalocuprate(I) anions, **227**, 153
- synthesis and X-ray structure of the dinuclear  $\mu$ -5,5'-diamino-3,3'-bis-1,2,4-triazolato(1-)-bis(diethylenetriamine)copper(II) triperchlorate,  $[\{\text{Cu}(\text{dien})\}_2\text{Hdabt}](\text{ClO}_4)_3$ , **227**, 181
- structural characterization of bidentate carboxylate derivatives of copper(I) bistrisphenylphosphine, **227**, 223
- Copper metallothionein  
oxidative quenching of luminescence from copper metallothionein, **226**, 275
- Coupling reaction  
gold in organic synthesis Part 2. Preparation of benzyl-alkyl and -aryketones via C–C coupling, **220**, 1  
investigation of the dithiocarbamate-induced coupling of allylidene and carbonyl ligands on a tungsten center, **220**, 335
- Cross-coupling  
chele- and anticheleselectivity IV. Reaction-accelerating neighboring-group effects in the cross-coupling of vinylbromides with alkyl-transition metal reagents, **220**, 85
- Cross-coupling reaction  
synthesis of symmetric and unsymmetric 1,4-bis(*p*-R-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289
- Crown ether complexes  
synthesis and crystal structure of  $[\text{GdCl}_2(\text{dibenzo-18-crown-6})(\text{MeCN})][\text{SbCl}_6] \cdot 2\text{MeCN}$ , dibenzo-18-crown-6 = 2,3,11,12-dibenzo-1,4,7,10,13,16-hexaoxacyclooctadeca-2,11-diene, **215**, 209  
influence of pendant cyanoalkyl side-arms on the stabilities of metal ion complexes of monoaza- and diaza-18-crown-6 ethers in methanol, **219**, 31  
structures of isomeric substituted 14-crown-4 ethers and their LiSCN complexes, **223**, 125  
electrochemical recognition of Group 1 and 2 metal cations by redox-active ionophores, **225**, 137
- Crown thioether complexes  
synthesis and complexation behaviour of a double-armed crown thioether ligand, **217**, 1
- Cryptand complexes  
the complexation of divalent metal ions by the cryptands 4,7,13,16-tetraoxa-1,10-diazabicyclo[8.8.2]eicosane, -[8.8.5]tricosane and -[8.8.8]hexacosane in aqueous solution, **215**, 79
- Cryptate complexes  
syntheses, spectroscopy and crystal structure of several binuclear silver(I) cryptates, **223**, 181
- Crystal structures  
towards a model for galactose oxidase: the crystal structure of {2-[bis(2-pyridylethyl)aminomethyl]phenolato} copper(II) perchlorate, **215**, 1  
preparation and structural characterization of  $\text{Rh}_2(\text{O}_2\text{CCPh}_3)_4(\text{EtOH})_2$ ,  $\text{Ru}_2(\text{O}_2\text{CCPh}_3)_4(\text{H}_2\text{O})(\text{EtOH}) \cdot 2\text{EtOH}$  and  $\text{Mo}_2(\text{O}_2\text{CCPh}_3)_4 \cdot 3\text{CH}_2\text{Cl}_2$ , **215**, 9  
cyclometallated derivatives of platinum(II) derived from 1,4-benzodiazepin-2-ones. Crystal and molecular structure of  $\text{Pt}(\text{L})(\text{HL})\text{Cl}$  ( $\text{HL} = 7\text{-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one}$ , DIAZEPAM), a molecule containing a neutral and a deprotonated 1,4-benzodiazepin-2-one, **215**, 17  
the structure of  $(\text{CH}_3)_3\text{Si}_3$ . Comparison between the structure in the solid and liquid state, **215**, 33  
phosphine and phosphine oxide adducts of vanadium(IV) chloride: synthesis and structural studies, **215**, 47  
mixed-ligand copper(II) complexes with iminodiacetato and 2- or 4-methylimidazole: molecular and crystal structure of the 'remote' isomer iminodiacetato(5-methylimidazole)-copper(II) monohydrate,  $[\text{Cu}(\text{IDA})(5\text{MeImH})] \cdot \text{H}_2\text{O}$ , **215**, 73  
synthesis and characterization of lead(II)–semicarbazone complexes. Crystal structures of dinitrato di-2-pyridylketone semicarbazone lead(II) and dichloro cyclohexane-1,2-dione bis(semicarbazone) dimethylformamide lead(II), **215**, 85  
cyclometallation of *N,N*-dimethylbenzo[*b*]furan-2-carbothio (and seleno) amides with palladium(II), ruthenium(II) and rhodium(III), **215**, 91  
the unusual crystallographic disorder of  $\text{M}_2$  units in  $\text{M}_2\text{Cl}_4(\text{PEt}_3)_4$  ( $\text{M} = \text{Re}, \text{Mo}$ ), **215**, 103  
synthesis, crystal and molecular structures, UV–Vis spectroscopy and electrochemical properties of two iron(III) phenolate complexes, **215**, 123  
preparation, structure and catecholase-mimetic activity of two mononuclear ferrocenecarboxylate copper(II) complexes, **215**, 131  
neutral transition metal complexes with the tridentate ligand 4-(5-methyl-4-imidazolyl)-3-thiabutanonic acid. A series of isostructural octahedral compounds, **215**, 151  
mixed-ligand complexes of rhenium II. Synthesis and characterization of  $[\text{ReN}(\text{X})(\text{Me}_2\text{PhP})_2(\text{R}_2\text{tcb})]$  complexes ( $\text{X} = \text{I}, \text{N}_3, \text{SCN}, \text{CN}$ ;  $\text{R}_2\text{tcb}^- = N\text{-(}N,N\text{-dialkyl-thiocarbamoylbenzamidinate)}$ ). X-ray crystal structure of  $[\text{ReN}(\text{N}_3)(\text{Me}_2\text{PhP})_2(\text{Et}_2\text{tcb})]$ , **215**, 159  
preparation and characterization of  $[(\text{LL})\text{Cl}_2\text{Mo}]_2(\mu\text{-SPr})_2$  compounds with  $\text{Pr} = n\text{-propyl}$  and  $\text{LL} = \text{PrSCH}_2\text{CH}_2\text{SPr}$  and  $\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2$ . Further examples of C–S bond cleavage and oxidative addition across an Mo–Mo quadruple bond, **215**, 191

- the crystal structure of a chelate silver(I) complex:  $\{\text{Ag}[\text{Ph}_2\text{P}(\text{CH}_2)_2\text{SCH}_2\text{CH}_3]_2\text{ClO}_4\}$ , **215**, 199
- synthesis and crystal structure of  $[\text{GdCl}_2(\text{dibenzo-18-crown-6})\text{-}(\text{MeCN})][\text{SbCl}_6] \cdot 2\text{MeCN}$ , dibenzo-18-crown-6 = 2,3,11,12-dibenzo-1,4,7,10,13,16-hexaoxacyclooctadeca-2,11-diene, **215**, 209
- synthesis of bis- or tris(pentafluorophenyl) thallium(III) complexes with bidentate ligands. X-ray crystal structure of  $[\text{Tl}(\text{C}_6\text{F}_5)_2(\text{acac})\{(\text{O})\text{Ph}_2\text{PCH}_2\text{PPh}_2(\text{O})\}]$ , **215**, 229
- two silicon-bridged cyclopentadienyl  $\text{Fe}_2\text{S}_4$  cluster units connected through an  $\text{Fe}(\text{III})\text{S}_6$  unit. Crystal and molecular structure of  $[\{(\text{CH}_3)_2\text{SiCp}_2\}_2\text{Fe}_2(\mu_3\text{-S}_2)(\mu_4\text{-S}_2)(\mu_4\text{-S}_4)]\text{FeCl}_4$ , **216**, 1
- $[\{\text{Ni}(3,2,3\text{-tet})(\mu\text{-NCO})\}_n](\text{ClO}_4)_n$ : structural characterization of the first monodimensional nickel(II)–cyanato system with a description of its ferromagnetic coupling, **216**, 5
- a tris-( $\mu_2$ -I) face-sharing dizirconium(III) complex,  $[\text{I}_4\text{-Zr}_2\text{I}_7(\text{PEt}_3)_2]^-$ ; the first such dizirconium species, **216**, 9
- oxo- and sulfido-bridged sandwich cubane-type molybdenum–indium mixed-metal cluster. Synthesis and X-ray structure of  $[(\text{H}_2\text{O})_9\text{Mo}_3\text{S}_3\text{OInOs}_3\text{Mo}_3\text{-}(\text{H}_2\text{O})_9\text{-}(\text{CH}_3\text{-C}_6\text{H}_4\text{-SO}_3)_8 \cdot 30\text{H}_2\text{O}]$ , **216**, 13
- dithiocarbamates of dimethylthallium(III). Crystal structure of dimethyl(di-*n*-propyl)dithiocarbamate)thallium(III), **216**, 15
- binuclear oxo-bridged iron(III) complexes of *N,N'*-bis(2-pyridylmethyl)ethane-1,2-diamine (bispicen) and related ligands, **216**, 21
- mixed-ligand complexes of rhenium III. Synthesis, characterisation and structures of  $[\text{ReN}(\text{Me}_2\text{PhP})_2(\text{mnt})]$  and  $[\text{ReN}(\text{Me}_2\text{PhP})(2,6\text{-DMTP})(\text{morphtc})]$  ( $\text{Me}_2\text{PhP}$  = dimethylphenylphosphine; 2,6-DMTP<sup>−</sup> = 2,6-dimethylthiophenolate; morphtc<sup>−</sup> = *N*-(morpholinylthiocarbonyl)benzamidinate), **216**, 31
- transition metal compounds of the tridentate pyrazole substituted amine ligand bis(2-(3,5-dimethyl-1-pyrazolyl)ethyl)ethylamine (ddae). X-ray structures of  $[\text{Co}(\text{ddae})(\text{NO}_3)_2]$ ,  $[\text{Cu}(\text{ddae})(\text{NO}_3)(\text{H}_2\text{O})](\text{NO}_3)$  and  $[\text{Cu}(\text{ddae})(\text{Cl})_2] \cdot \text{C}_2\text{H}_5\text{OH}$ , **216**, 43
- unexpected five-coordination in di(1,3-diaminopropane)-3,5-dinitrobenzoatecopper(II) 3,5-dinitrobenzoate. Comparison between the coordinated and non-coordinated anions, **216**, 57
- a polymeric two-dimensional mixed-metal network. Crystal structure and magnetic properties of  $\{[\text{P}(\text{Ph})_4]\text{-}[\text{MnCr}(\text{ox})_3]\}_n$ , **216**, 65
- synthesis and characterization of platinum(II)–thiolate complexes. Crystal and molecular structures of *cis*- $\text{Pt}(\text{Ph}_3\text{P})_2(\text{SC}_6\text{H}_2\text{-2,4,6-iPr}_3)\text{Cl}$  and *cis*- $\text{Pt}(\text{Ph}_3\text{P})_2(\text{SC}_6\text{H}_2\text{-2,4,6-iPr}_3)_2$ , **216**, 83
- bulky substituent effects on the iron(III) complexation of 1,4,7-triazacyclononane, **216**, 89
- evidence of the interaction between steric and electronic influence in rhodoximes and cobaloximes. Synthesis of  $\text{pyRh}(\text{DH})_2\text{I}$  and X-ray structure of  $\text{pyRh}(\text{DH})_2\text{Cl}$ ,  $\text{pyCo}(\text{DH})_2\text{Cl}$  and  $\text{pyRh}(\text{DH})_2\text{I}$ , **216**, 125
- octahedral  $\mu$ -oxalato–nickel(II) dinuclear complexes with water and tridentate amines as blocking ligands: magnetostructural correlations, **216**, 139
- structure and spectroscopic properties of  $[\text{di}(3\text{-aminopropyl)amine-}N,N',N''][\text{iminodiacetato-}N,O,O']\text{chromium(III) perchlorate}$ , **216**, 155
- molecular and crystal structures of *cis*- $\text{Pt}(\text{DMSO})(\text{C}_2\text{H}_5\text{CN})\text{Cl}_2$  and  $\text{Cl}(\text{DMSO})\text{Pt}(\mu\text{-N}(\text{CH}_3)\text{CHO})_2\text{Pt}(\text{DMSO})\text{Cl}$ , **216**, 163
- diorganotin(IV) derivatives of salicylaldehydethiosemicarbazone. The crystal structure of dimethyl- and diphenyl(salicylaldehydethiosemicarbazonato)tin(IV), **216**, 169
- ionic cluster dithiophosphinates  $[\text{Mo}_3\text{S}_4(\text{R}_2\text{PS}_2)_3\text{py}_3]^+[\text{Ml}_3]^-$  ( $\text{M} = \text{Zn} \cdot \text{py}$ ,  $\text{Cd} \cdot \text{py}$ ,  $\text{Hg}$ ), **216**, 177
- crystal structure, magnetic and spectroscopic properties of the bis(dimethyl sulfoxide) adduct of tetra- $\mu$ -formato-dicopper(II), a new tetracarboxylato-bridged copper(II) dimer, **216**, 185
- evidence of the  $\text{W}(\text{O})(\text{S}_2)_2$  core as an intermediate in the acidification of  $\text{WS}_2^{2-}$ . Structural characterization of  $\text{W}(\text{O})(\text{S}_2)_2(\text{bpy})$  and  $\text{W}(\text{O})(\text{S}_2)_2(\text{phen})$ , **216**, 201
- polyfunctional phosphine ligands II. Iridium(I) complexes of the 7-diphenylphosphino-2,4-dimethyl-1,8-naphthyridine (dpnapy) ligand. X-ray crystal structure of  $[\{\text{Ir}(\text{cod})\text{Cl}\}_2(\mu\text{-dpnapy})]$ , **216**, 209
- isothiocyanato complexes of rhenium II. Synthesis, characterization and structures of  $\text{ReN}(\text{NCS})_2(\text{Me}_2\text{PhP})_3$  and  $\text{Re}(\text{NS})(\text{NCS})_2(\text{Me}_2\text{PhP})_3$ , **216**, 223
- cobalt(III) complexes with 2-acetyl- and 2-formylpyridine <sup>4</sup>*N*-methylthiosemicarbazones, **216**, 229
- a novel amino acid-binding receptor based on weak non-covalent ligand–ligand interactions, **216**, 249
- synthesis, characterization and crystal structure of  $(\text{C}_3\text{H}_5\text{N}_2)_2[\text{Sn}_2(\text{NO}_3)_4(\mu\text{-OH})_2\text{Me}_4]$ , a di- $\mu$ -hydroxo-bridged compound with heptacoordinated tin(IV), **216**, 257
- synthesis, crystal structure and thermal stability of the  $\text{MgCl}_2$  formamide adduct, **216**, 261
- trimethylgallium based phosphinogallanes. Synthesis and molecular structure of  $[\text{Me}_2\text{Ga-P}(\text{SiMe}_3)_2]_2$ , **216**, 267
- a seven-spin cluster formed by an alkyl nitronyl nitroxide biradical and copper(II): crystal structure and magnetic properties, **217**, 7
- a novel tetranuclear iodoargentate(I) ion,  $[\text{Ag}_4\text{I}_8]^{4-}$ , with a cubane-type core, **217**, 15
- interaction of the lysine side chain amino group with  $\text{Cu}^{\text{II}}$  in (glycyl-*L*-lysine<sup>2-</sup>)<sub>2</sub>Cu, **217**, 33
- magnetic properties and crystal structure of  $\{\mu\text{-pyrazine}\}\text{-bis}[(\text{aqua})(2,6\text{-xylylamino}(\text{diacetato})\text{copper(II)})]$ . A pyrazine-bridged dinuclear copper(II) complex, **217**, 39
- syntheses of dinuclear gold(I) ring complexes containing 1,1-dicyanoethene-2,2-dithiolate-*S,S* and bis(diphosphines) as bridging ligands. X-ray crystal structure of  $[\text{Au}_2\mu\text{-}(\text{i-MNT})\mu\text{-}(\text{dppee})]$ , **217**, 45
- bis(2-diphenylphosphinoethyl)phenylphosphineruthenium(II) complexes of amino acids and dipeptides, **217**, 51
- spectroscopic investigation of dichloro and hydrazido(2−) derivatives of bis(dithiocarbamate)dioxomolybdenum(VI) compounds. Crystal and molecular structure of bis(diisobutyl-dithiocarbamate)(*N,N*-dimethylhydrazido(2−)-*N'*)-oxomolybdenum(VI), **217**, 61
- spectroscopic and structural studies on volatile gallium  $\beta$ -diketonates as potential precursors for MOCVD, **217**, 71
- a mixed-metal salt comprised of metal–metal bonded dinuclear ions: structure and properties of  $[\text{Rh}_2(\text{OAc})_2(\text{MeCN})_6][\text{Re}_2\text{Cl}_8]$ , **217**, 79
- mercury and platinum mixed cluster having a prismatic structure,  $[\text{HgPt}_6(2,6\text{-Me}_2\text{C}_6\text{H}_3\text{NC})_{12}]$ , **217**, 121
- synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129
- nucleophilic addition of water to coordinated di-(2-pyridyl)-ketone (DPK) in rhenium(V) complexes. Synthesis and crystal structure of the adduct  $(\text{DPK} \cdot \text{H})^+[\{\text{ReOCl}_2\text{-}(\text{DPK} \cdot \text{OH})\}_2\text{Cl}]$ , **217**, 141
- synthesis and characterization of Cd(II) complexes of cinoxacin. Crystal structure of a dicadmium complex containing two monoatomic-bridging carboxylate oxygen atoms, **217**, 149

- complexes of mercuric halides with the tripod ligand  $np_3$  ( $=N(CH_2CH_2PPh_2)_3$ ). Molecular structure of  $[(np_3)HgI] \cdot 0.5C_2H_5OH$ , **217**, 155
- synthesis and characterization of symmetric and unsymmetric oxo-bridged trinuclear chromium benzoate complexes: crystal and molecular structure of  $[Cr_3O(O_2CPh)_6(py)_3]ClO_4$ , **217**, 171
- synthesis and structure of  $Zr_2Cl_4(\eta^5-C_6H_5PMe_2)_2(PMe_2Ph)_2$ , an arene complex of zirconium(II), **217**, 187
- functionalised phosphines: structure of bis(2-aminophenyl)diphenylphosphine- $\kappa^2N,P$ copper(I) perchlorate ethanol solvate, **217**, 191
- the synthesis and crystal structure of  $[Cr(acacen)py_2][ZnCl_3py]$ , **217**, 195
- synthesis and characterization of phosphane-substituted hydroxyruthenocenes: crystal structure of  $[Ru(\eta^5-C_5H_4PPh_3)(\eta^5-C_5H_4OH)]PF_6$ , **217**, 203
- azacryptand host for lead: NMR and structural studies, **218**, 5
- molybdenum carbonyl derivatives containing eight- and nine-membered diphosphine chelating rings: the synthesis of  $[Mo(CO)_4(dpppe)]$  ( $dpppe = Ph_2P(CH_2)_3PPh_2$ ) and  $[Mo(CO)_4(dpph)]$  ( $dpph = Ph_2P(CH_2)_6PPh_2$ ) and the crystal structure of the latter, **218**, 9
- alkyl for hydride exchange between alane-trimethylamine and Group IVB metal alkyls, **218**, 47
- substituent effect on the coordination ability of the amide group of N-protected amino acids, **218**, 53
- hydrothermal synthesis and crystal and molecular structure of a three-dimensional vanado-arsenate  $(NH_4)_3[(VO)(V_2O_3)(AsO_4)_2(HAsO_4)]$ , exhibiting both  $\{V-O-V\}_\infty$  chains and isolated  $\{V_2O_3\}$  units, **218**, 59
- organometallic complexes of ruthenium(IV) with bidentate oxygen and nitrogen donor ligands, **218**, 81
- a mechanistic study of the homogeneous catalytic hydroformylation of formaldehyde: synthesis and characterization of model intermediates, **218**, 89
- synthesis, structural characterization and dioxygen reactivity of imidazole-ligated Cu(I) complexes, **218**, 103
- preparation, crystal structures and magnetic properties of a *syn-anti* carboxylate-bridged dinuclear copper(II) complex and its precursor, a hydrogen-bonded polynuclear copper(II) complex, **218**, 109
- simultaneous binding of soft and hard metals to a pyrimidine nucleobase: *trans*- $K_2[Pt_2(1-MeU)_2] \cdot 6H_2O$ , an anionic Pt-1-methyluracil (1-MeU) complex arranged in circles, **218**, 117
- lithium coordination to amino acids and peptides. Synthesis, spectroscopic characterization and structure determination of lithium complexes of neutral and anionic glycine and diglycine, **218**, 121
- main group metal halide complexes with sterically hindered thioureas XIV. A reinvestigation of coordination sensitive modes in the solid state vibrational spectrum of 1,3-dimethyl-2(3*H*)-imidazolethione and related compounds, **218**, 133
- new heptacoordinated tin complexes of 2,6-diacetylpyridine-bis(thiosemicarbazone),  $H_2daptsc$ , and of 2,6-diacetylpyridine-bis(semicarbazone),  $H_2dapsc$ . Crystal and molecular structures of  $[MeSnCl(Hdaptsc)Cl] \cdot MeOH$  and  $[MeSnCl(H_2dapsc)Cl]_2 \cdot 2H_2O$ , **218**, 139
- preparation and crystal structure of *anti*-( $\eta^3$ : $\eta^5$ -azulene)-pentakis(trimethylphosphine)dibalt. Reactions with carbon monoxide and solution dynamics, **218**, 143
- conformational flexibility of 1,3-diaminopropane in chelate rings. Redetermination of the structure of *trans*-dinitratobis(1,3-diaminopropane)copper(II), **218**, 151
- synthesis and characterization of oxorhenium(V)-catecholate complexes. Crystal and molecular structures of  $(CH_3)_4N[ReO(O_2C_6H_4)_2]$  and  $(CH_3)_4N[ReO(PPh_3)(O_2C_6H_4)_2] \cdot CH_3OH$ , **218**, 157
- antiferromagnetism tuned by the topology of the intervening bridging ligand. Crystal structure and magnetic properties of the  $(\mu$ -phthalato){bis(piperidine)}Cu(II) chain complex, **218**, 167
- coordination chemistry of perhalogenated cyclopentadienes and alkynes XIV. Synthesis of dinuclear ring-bridged cyanotrenes with  $-SiMe_2-$  or  $-TiCp_2-$  bridges. Structures of  $[(OC)_3Mn(C_5Br_4^-)]_2SiMe_2$  and  $[(OC)_3Mn(C_5Cl_4^-)]_2TiCp_2$ , **218**, 173
- structural characterization of octahedral metal complexes of  $[MX_3(py)_3]$  type with three pyridines as revealed by  $^2H$  NMR spectra, **218**, 179
- crystal structure and magnetic properties of the first 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position:  $\{catena-(\mu_{N,S}-NCS)[Ni(en)_2]\}(PF_6)$ , **218**, 185
- orthometalation reactions in trifluoroacetate dirhodium(II) compounds. Molecular structure of  $Rh_2(O_2CCF_3)_2 \cdot [(C_6H_4)PPh_2]_2 \cdot (PPh_3)_2 \cdot 2(C_7H_8)$ , **218**, 189
- preparation and structure of  $[PhPt(OCO_2H)(PEt_3)_2]$  and  $[PhPt(OH_2)(PEt_3)_2]BF_4$ , **218**, 199
- cyclopalladated *p*-azoxyanisole complexes. Synthesis and structure of  $\{[bis(2\text{-phenylpyridine})][4'-(\text{methoxyphenyl-}NNO\text{-azoxy-}N^2)\text{-4-methoxyphenyl-2-ato}]palladium(II)\}$  (tetrafluoroborate), **218**, 211
- novel metal thiolates:  $[Mn_2(SC_6H_3[SiMe_3]_2)_4(C_4H_8O)]$ , a binuclear manganese thiolate complex with tetrahedral and trigonal-planar metal coordination, **219**, 1
- a new class of dinuclear compounds: the synthesis and X-ray structural characterization of tris( $\mu$ -diphenylformamidinato)diiron, **219**, 7
- crystal and molecular structure of the solvate of bis(2,6-bis(1'-methylbenzimidazol-2'-yl)pyridine)cobalt(II) dihexafluorophosphate with 4-methyl-1,3-dioxolan-2-on, **219**, 11
- dimolybdenum(II) and diruthenium(II) complexes with 2-substituted 7-methyl-1,8-naphthyridines as bridging ligands, **219**, 23
- electrochemical and structural studies of nickel(II) complexes with  $N_2O_2$  Schiff base ligands 2. Crystal and molecular structure of *N,N'*-1,2-ethane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II), *N,N'*-1,2-*cis*-cyclohexane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II) and *N,N'*-1,2-benzene-1,2-diyl-bis(3,5-dichlorosalicylideneimine)nickel(II), **219**, 43
- zinc(II) and cadmium(II) complexes of the potentially tridentate ligands 2-[[2-pyridinylmethyl]amino]methyl]phenol (HSALAMP) and 2-[[[2-(2-pyridinyl)ethyl]amino]methyl]phenol (HSALAEP). X-ray crystal structures of the  $[Zn(SALAMP)(NO_3)]_2$  and the  $[Cd(SALAEP)(NO_3)]_2$  dimers, **219**, 55
- organometallic chelating ligands. Synthesis of a new anionic tripod like ligand with an *N,N',O*-donor set. X-ray crystal structure of  $[(\eta^6-C_6Me_6)Ru(\mu-P(O)(OMe)_2)(\mu-Pz)_2Re(CO)_3]$  (*Me* = methyl, *Pz* = pyrazolate), **219**, 67
- transition metal complexes with sulfur ligands Part CII. Changes in configuration and reactivity caused by alkyl substituents *R* in  $[Fe(L)(N_R S_4)]$  complexes (*L* = CO,  $NO^+$ , NO,  $PMe_3$ ,  $N_2H_4$ ; *R* =  $CH_3$ ,  $CH_2CH_2CO_2CH_3$ ,  $CH_2CH_2CO_2^-$ ), **219**, 75
- the study of isolobal displacement reactions. Synthesis and structures of the heterobimetallic  $\mu$ -alkyne cluster complexes  $\eta^5-R_3C_5H_4(CO)_2M(\mu-R^1C_2R^2)Co(CO)_3$  (*M* = Mo, W; *R* =  $CH_3C(O)$ ,  $CH_3O_2C$ ), **219**, 93
- synthesis, crystal structure and spectroscopic properties of bis(diphenyldithiophosphinato)cadmium(II), **219**, 115



- reactions of the polynuclear chain complex  $[\text{Cu}(\text{L})(\text{H}_2\text{O})]_n$  ( $\text{H}_2\text{L}$  is 5-[(2'-carboxyphenyl)azo]-1,3-dimethylbarbituric acid) with imidazole. Crystal structures of the dinuclear and mononuclear derivatives and EPR study of the dimer–monomer equilibrium in solution, **219**, 127
- preparation, properties and coordination behaviour of planar or tridimensional compartmental Schiff bases, **219**, 135
- new methods for synthesis of platinum(II) dimethyl sulfide complexes, **219**, 155
- coordination polymers of Mn(II) with the ligand pyrazine-2,3,5,6-tetracarboxylic acid, **219**, 161
- variation of the exchange interaction in oximate-bridged  $\text{Cu}^{\text{II}}\text{M}^{\text{II}}$  dimers ( $\text{M}=\text{Cu}, \text{Ni}, \text{Mn}$ ). Crystal structure of  $[\text{Cu}(\text{pdmg})\text{Cu}(\text{bipy})(\text{H}_2\text{O})_2](\text{ClO}_4)_2 \cdot \text{H}_2\text{O}$ , **219**, 179
- synthesis and X-ray crystal structure of *cis*-1,4-diaminocyclohexanetetrachloroplatinum(IV): a new antitumor agent, **219**, 193
- the synthesis, structure and hydrolytic activity of a functional pentacoordinate zinc complex, **219**, 205
- synthesis and X-ray structure of [acenaphthene–chromium(0)-dicarbonyl-tri(2,3,5-trimethylphenyl)phosphite], **219**, 209
- bis(2-mercaptopyridine-*N*-oxide)oxovanadium(IV): synthesis, structure and its conversion to the dichloro complex, **219**, 213
- synthesis of 5,6-dihydro-4*H*-1,3-oxazines from neutral and cationic platinum(II) nitrile complexes. X-ray structure of *trans*- $[\text{Pt}(\text{CF}_3)(\text{N}=\text{C}(\text{Ph})\text{OCH}_2\text{CH}_2\text{CH}_2)(\text{PPh}_3)_2]\text{BF}_4$ , **220**, 21
- photochemical reactions of transition metal organyl complexes with olefins Part 10. Light-induced formal [5 + 2] cycloadditions at manganese, **220**, 77
- preparation, properties and reactions of metal-containing heterocycles Part LXXXIX. Metalla[*n*]- and metalla[*m,n*]-orthocyclophanes as mediators for the synthesis of cyclic ketones, **220**, 107
- synthesis of  $\eta^3$ -allyl–Pd complexes and subsequent formation of N-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- reactions of benzocyclobutene chromium complexes with carbon, nitrogen and oxygen nucleophiles: nucleophilic addition and unexpected proximal ring opening reactions, **220**, 175
- dirhodium(II) tetrakis[*N,N*-dimethyl-2-pyrrolidone-5(*S*)-carboxamide]. Structural effects on enantioselection in metal carbene transformations, **220**, 193
- reactivity of allene at phosphine-bridged di-iron centres: X-ray crystal structures of  $[\text{Fe}_2(\text{CO})_5\{\mu\text{-}\sigma, \eta^3\text{-C}(\text{O})\text{C}(\text{CH}_2)_2\}(\mu\text{-dppm})]$  and  $[\text{Fe}_2(\text{CO})_4\{\mu\text{-}\eta^3, \eta^3\text{-}(\text{CH}_2)_2\text{C}_2(\text{CH}_2)_2\}(\mu\text{-dppm})] \cdot \text{Et}_2\text{O}$ , **220**, 201
- a study of the reactivity and structure of cyclic  $\alpha, \beta$ -unsaturated Fischer-type carbene complexes, **220**, 233
- direct nitration of aryl groups  $\sigma$ -bound to ruthenium(II), **220**, 261
- the synthesis and reactions of A-ring aromatic steroid complexes of manganese tricarbonyl, **220**, 297
- $[\text{Rh}_2(\text{O}_2\text{CCH}_3)_4\text{TCNE}]_n$  polymers: isomers with either 1,1- or *trans*-1,2-TCNE bridges, **221**, 1
- ten-vertex polyhedral dicarborane chemistry: new positional isomers of cluster constituents in the ten-vertex *arachno* dicarborane family: the parent [*arachno*-5,10- $\text{C}_2\text{B}_8\text{H}_{13}$ ]<sup>−</sup> anion and the isomeric ligand derivatives *exo*-9-*L*-*arachno*-5,6- $\text{C}_2\text{B}_8\text{H}_{12}$  and *exo*-6-*L*-*arachno*-5,10- $\text{C}_2\text{B}_8\text{H}_{12}$ , **221**, 5
- an ethyl–cobalt complex with a buckled equatorial ligand, **221**, 35
- $\pi$ -facial selectivities in the coordination of an optically pure fused cyclopentadienyl ligand to arene ruthenium moieties. NMR spectral and structural characterization of a single diastereomer of  $[(\text{PCp})\text{Ru}(\text{Me}_6\text{C}_6)]\text{PF}_6$ , **221**, 41
- synthesis, characterization and X-ray crystal structure of the benzenedithiolatohafnium(IV) porphyrin complex, **221**, 51
- synthesis and spectroscopic properties of diorganotin(IV) derivatives of 2,6-diacetylpyridine bis(thiosemicarbazone). Crystal structure of diphenyl{2,6-diacetylpyridine bis(thiosemicarbazonato)}tin(IV) bis(dimethylformamide) solvate, **221**, 61
- crystal structure and ligand field states of  $[\text{Cr}(\text{NH}_3)_5\text{NCO}(\text{NO}_3)_2]$ , **221**, 93
- nickel(II) and manganese(II) complexes of substituted phenanthroline ligands, **221**, 99
- synthesis and structural characterization of a new series of ruthenium(II) complexes containing the short bite  $\text{Ph}_2\text{PPy}$  ligand. Cooperative effect between the anionic rhodium and cationic ruthenium species in the catalytic hydroformylation of styrene by  $[\text{Ru}(\text{Ph}_2\text{PPy})\text{Cl}][\text{Rh}(\text{CO})_2\text{Cl}_2]$ , **221**, 109
- synthesis and X-ray structural determination of pentamethylcyclopentadienyl dichloro oxo molybdenum, **221**, 147
- binuclear complexes from 1,8-substituted anthracene derivatives. The crystal structure of  $\{[\mu\text{-}[1,8\text{-anthracenedicarboxaldehydebis(dimethylhydrazone)]]\text{tetrachlorobis}(\eta^2\text{-ethylene})\text{di-platinum}]\}$ , **221**, 151
- the reaction of  $\text{CpRu}(\text{PPh}_3)_2^+$  with trimethylenesulfide. Synthesis and molecular and crystal structure of  $[\text{CpRu}(\text{PPh}_3)_2(\text{SC}_3\text{H}_6)]\text{CF}_3\text{SO}_3$ , **221**, 157
- synthesis and molecular structure of two crystalline forms of hydrated cerium(III) picrate, **221**, 169
- synthesis, characterization and X-ray crystal structure of ethylenediamine(oroato)palladium(II), **221**, 173
- zn(II) complexes with thiopyrimidine derivatives: solution study, synthesis and crystal structure of a zig-zag chain zinc(II) complex with the ligand 4,6-dimethyl-2-thiopyrimidine, **221**, 177
- chiral metal complexes 40. The sterically mediated hydrolysis of a coordinated imine, **221**, 191
- palladium-assisted formation of carbon–carbon bonds Part 2. Stoichiometric synthesis of spirocyclic compounds. X-ray structure of a  $\pi$ -allylic palladium intermediate, **222**, 1
- spontaneous formation of a 1D-coordination polymer with helix structure starting from (*S*)-pyroglutamic acid and copper(II) acetate, **222**, 5
- synthesis and structure of  $\text{Cr}(\text{CO})_3$  complexes of biphenyl compounds with axial chirality, **222**, 63
- carbonyl carbene coupling on cationic  $\eta^2$ -carbene complexes of molybdenum and tungsten, **222**, 77
- addition of carbanions to tris(trimethylphosphine)(cyclooctadiene)-iridium(I) chloride: three different modes of reaction depending on the nature of the carbanion, **222**, 135
- intermediates in the palladium-catalysed reaction of 1,3-dienes Part 8. The reaction of palladium–butadiene complexes with ethyl methylacetoacetate, **222**, 179
- insertion of NH into the rhenium–carbon bonds of a Fischer-type rhenacyclobutadiene. Preparation and characterization of azarhenacyclic complexes, **222**, 199
- synthesis and structure of new chiral ferrocenylphosphines for asymmetric catalysis, **222**, 213
- synthesis, structure, dynamics and first atropisomer-selective cleavage of the chromium tricarbonyl complex of a lactone-bridged biaryl, **222**, 255
- reactions of  $[\text{Au}(\text{C}_6\text{F}_5)(\text{SC}_4\text{H}_8)]$  with diazoalkanes. Synthesis and molecular structures of  $[\text{Au}(\text{C}_6\text{F}_5)(\text{Ph}_2\text{C}=\text{N}=\text{N}=\text{CPh}_2)]$ , **222**, 267
- a novel reaction of the cyano ligand opening an organometallic route to 4-amino imidazoles: four component condensation (4CC) with isocyanide, aldehyde and amine, **222**, 275

- nitrosoarene and nitrosoalkane insertion reactions of titanacyclobutenes, **222**, 305
- reactions of aziridine with platinum(II) nitriles. Formation of (aziridino)amidines and 2-imidazolines and X-ray structure of *trans*-[PtCl<sub>2</sub>{N(H)=C(Ph)NCH<sub>2</sub>CH<sub>2</sub>}]<sub>2</sub>, **222**, 327
- synthesis, structure and solid state polymerization of diaquabis(propynoato)magnesium, **222**, 339
- reaction of primary silanes with dinuclear rhodium hydride complexes: silane coupling reactions, **222**, 345
- extrusion of molecular clusters from solid-state materials: synthesis by application of  $\gamma$ -irradiation. Molecular and crystal structure of (H<sub>3</sub>O<sub>4</sub>)(Et<sub>4</sub>N)[Mo<sub>3</sub>S<sub>7</sub>Br<sub>6</sub>], **223**, 5
- binuclear platinum(II) complexes of 1,1,4,7,10,10-hexaphenyl-1,4,7,10-tetraphosphadecane (P4) containing various bridging ligands as spacers between the P4 coordination units, **223**, 13
- stabilities of 1,2-diethyl-3-hydroxy-4-pyridinone chelates of divalent and trivalent metal ions, **223**, 21
- synthesis, structure and stability of complexes of lanthanide nitrates with [(MeO)<sub>2</sub>P(O)]<sub>2</sub>C(OH)Ph and the isomeric (MeO)<sub>2</sub>P(O)OCHPhP(O)(OMe)<sub>2</sub>, **223**, 43
- di- and tetranuclear gold(II) complexes with dithiocarbamate and related ligands. X-ray structure of [Au<sub>2</sub>( $\mu$ -CH<sub>2</sub>PPh<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>(S<sub>2</sub>CNMe<sub>2</sub>)<sub>2</sub>], **223**, 55
- the reactions of thiolate ligands with rhenium hydrides Part 1. The syntheses and structures of [ReO(SC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>)<sub>3</sub>(PPh<sub>3</sub>)] and [Re(SC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>)<sub>3</sub>(PPh<sub>2</sub>H)(PPh<sub>3</sub>)], **223**, 63
- synthesis, spectroscopic and structural characterization of methyl pyruvate- and pyridoxal-hydrazinopyruvylthiosemicarbazones, **223**, 77
- direct electrochemical synthesis of tin(IV) complexes of 2-(2-mercaptoethyl)iminomethyl-phenols; the molecular structure of bis[2-(2-mercaptoethyl)iminomethyl-5-bromo-phenoxy]-tin(IV), **223**, 87
- the Zr<sub>6</sub>Cl<sub>12</sub>(PMe<sub>2</sub>Ph)<sub>6</sub> molecule revisited: a directed synthesis, **223**, 93
- metal complexes of 5-acetyl-2,2'-pyridylthiophene aroylhydrazones. Crystal structure of a cobalt(II) derivative, **223**, 97
- structures of isomeric substituted 14-crown-4 ethers and their LiSCN complexes, **223**, 125
- synthesis, properties and molecular structure of copper(II) and cobalt(II) 1,2-bis(pyrazol-1-ylmethyl)benzene complexes, **223**, 139
- synthesis and X-ray analysis of the dirhodium(II,II) carboxylate complex Rh<sub>2</sub>(O<sub>2</sub>CCH<sub>3</sub>)<sub>4</sub>(formH)<sub>2</sub> (formH = N<sub>2</sub>N'-di-*p*-tolylformamide), **223**, 145
- thermodynamics of complexation of the uranyl(VI) ion with benzoate. The crystal structure of sodium tris(benzoato)dioxouranate dihydrate, Na[(PhCO<sub>2</sub>)<sub>3</sub>UO<sub>2</sub>·2H<sub>2</sub>O], **223**, 151
- synthesis and characterization of the new peroxo complex aquaoxidiperoxoimidazole molybdenum(VI), **223**, 173
- syntheses, spectroscopy and crystal structure of several binuclear silver(I) cryptates, **223**, 181
- synthesis, characterization and crystal structures of two *cis*-dioxo vanadium(V) complexes of monoanionic tridentate Schiff base ligands, **223**, 193
- divalent iron formamidinato complexes: a highly distorted dinuclear compound, **224**, 5
- a new functionalised oligopyridine ligand containing ferrocene as a ball-bearing spacer for metallosupramolecular chemistry, **224**, 11
- transition metal complexes with sulfur ligands CIV. Protonation and alkylation of thiolate donors of [Fe(CO)<sub>2</sub>(S<sub>4</sub>')] and [Fe(CO)(S<sub>5</sub>')] complexes: syntheses and properties of Fe(II) complexes with open-chain thioether ligands (S<sub>4</sub>'<sup>2-</sup> = 1,2-bis(2-mercaptophenylthio)ethane(2-); S<sub>5</sub>'<sup>2-</sup> = 2,2'-bis(2-mercaptophenylthio)diethylsulfide(2-)), **224**, 45
- transition metal complexes with sulfur ligands CVI. [( $\mu$ -S<sub>2</sub>){Fe( $\mu$ -S<sub>4</sub>')}]<sub>2</sub>, a novel [2Fe-2S] cluster with octahedral low-spin [Fe<sup>II</sup>S<sub>4</sub>] centers (S<sub>4</sub>'<sup>2-</sup> = 1,2-bis(2-mercaptophenylthio)ethane(2-)), **224**, 35
- transition metal complexes with sulfur ligands CVI. Phosphine and azido [Fe(L)(N<sub>H</sub>S<sub>4</sub>')] complexes; structural characterization of the key complex [Fe(N<sub>H</sub>S<sub>4</sub>')·THF (L = PMe<sub>3</sub>, PBu<sub>3</sub>, PMe<sub>2</sub>Ph, PMePh<sub>2</sub>, N<sub>3</sub><sup>-</sup>; N<sub>H</sub>S<sub>4</sub>'<sup>2-</sup> = dianion of 2,2'-bis(2-mercaptophenylthio)diethylamine), **224**, 61
- phenylimido derivatives of [Mo<sub>6</sub>O<sub>19</sub>]<sup>2-</sup>: syntheses, X-ray structures, vibrational, electrochemical, <sup>95</sup>Mo and <sup>14</sup>N NMR studies, **224**, 81
- di-manganese carbonyl compounds containing bis(dicyclohexylphosphino)methane (dcpm) ligation, **224**, 105
- electrochemical behavior and X-ray crystal structure of the unusual dimeric ruthenium(cyclooctadiene) complex, (tri- $\mu$ -chloro)(chloro)(acetonitrile)bis( $\eta^4$ -cyclooctadiene)diruthenium(II), **224**, 125
- crystal and molecular structures of [*exo*- and *endo*-(RO)<sub>2</sub>P(O)- $\eta^5$ -C<sub>6</sub>H<sub>6</sub>]Mn(CO)<sub>3</sub>, **224**, 147
- synthesis, structure and properties of the polychalcogenides [M<sub>4</sub>Te<sub>12</sub>]<sup>4-</sup> (M = Cd, Hg), **224**, 163
- crystal structure and magnetic property of a 2D infinite layered copper(II) complex with bridging *trans*-oxamidate and pyrimidine ligands, **224**, 171
- isothiocyanato complexes of rhenium III. Synthesis and characterization of [ReN(NCS)<sub>2</sub>(Ph<sub>3</sub>P)<sub>2</sub>] and [ReN(NCS)<sub>2</sub>(CH<sub>3</sub>CN)(Ph<sub>3</sub>P)<sub>2</sub>]. X-ray crystal structure of [ReN(NCS)<sub>2</sub>(CH<sub>3</sub>CN)(Ph<sub>3</sub>P)<sub>2</sub>]·CH<sub>3</sub>CN, **224**, 193
- crystal structure of a binuclear vanadium(III) complex with a new tripodal ligand, [V<sub>2</sub>Cl<sub>4</sub>(tped)(EtOH)<sub>2</sub>]·2EtOH, **224**, 199
- the oxidation of [Co(edta)]<sup>2-</sup> by [Co(phen)<sub>3</sub>], **225**, 75
- mixed valence state associated with halogen migration in diruthenocylmethane and its related compounds, **225**, 103
- electrochemical recognition of Group 1 and 2 metal cations by redox-active ionophores, **225**, 137
- transition metal complexes with sulfur ligands Part CVII. Complexes of the [Mo(NO)(S<sub>4</sub>')] fragment with 16, 17 and 18 valence electrons (S<sub>4</sub>'<sup>2-</sup> = 1,2-bis(2-mercaptophenylthio)ethane(2-)), **225**, 163
- indium(III) halide-3,5-di-*tert*-butyl-*o*-benzosemiquinone systems, **225**, 207
- syntheses and structures of mixed-ligand tetrairon-sulfur clusters, [Cp'<sub>2</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)<sub>2</sub>Fe<sub>4</sub>S<sub>4</sub>](PF<sub>6</sub>) and [Cp'<sub>3</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)-Fe<sub>4</sub>S<sub>4</sub>](PF<sub>6</sub>)<sub>*n*</sub> (*n* = 1 and 2) (Cp' =  $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>). Structural change of the clusters accompanying electron transfer reactions, **225**, 229
- the redox behaviour of the cluster anion [Fe<sub>4</sub>N(CO)<sub>12</sub>]<sup>-</sup>. Electron transfer chain catalytic substitution reactions. Crystal structure of [Fe<sub>4</sub>N(CO)<sub>11</sub>PPh<sub>3</sub>], **226**, 1
- crystal structure and solid-state fluorescence of an oxomolybdenum(V)-benzylviologen electron donor-acceptor salt, **226**, 25
- valence tautomerism for catechol/semiquinone complexes of the *trans*-M(Bupy)<sub>2</sub>(3,6-DBQ)<sub>2</sub> (M = Mn, Fe, Co) series, **226**, 91
- simultaneous binding of soft and hard metals to a pyrimidine nucleobase: preparation and X-ray structure of [(en)Pd(1-MeT)<sub>2</sub>Na<sub>2</sub>](NO<sub>3</sub>)<sub>2</sub>·H<sub>2</sub>O (1-MeT = 1-methylthymine anion), **227**, 5
- the synthesis, X-ray structure and hydrolysis kinetics of chloro(2-aminomethylpyridine)(triamine)chromium(III) salts, **227**, 11
- synthesis and characterization of neutral oxorhenium(V) and nitridotchnetium(V) complexes with a tetradentate N<sub>2</sub>S<sub>2</sub>

- unsaturated ligand derived from dithiocarboxylic acid, **227**, 25
- one-equivalent oxidative charge storage on an  $[\text{Mn}_2\text{O}_2]^{3+}$  core: synthesis, crystal structure and resonance Raman spectra of  $[\text{Mn}_2\text{O}_2(\text{pbz})_4]^{n+}$  ( $n=3, 4$ ), **227**, 33
- synthesis and crystal structure of the  $\mu_2$ -{tris[bis-(diphenylphosphine)methane]}- $\mu_3$ -3-methyl-8-ethyl-xanthine-*N,O,O*]- $\mu_3$ -chloro-triangle-tricopper(I)-water complex exhibiting a novel oxopurine coordination mode, **227**, 43
- solid state and solution studies of a T-shaped gold(I) complex:  $\text{Au}[(\text{C}_6\text{H}_5)_2\text{PCH}_2\text{N}(\text{C}_6\text{H}_5)_2]_2\text{Cl}$ , **227**, 85
- synthesis of some dinuclear niobium chloro alkoxy carboxylates  $[\text{Nb}_2\text{Cl}_4(\text{OEt})_4(\text{O}_2\text{CR})_2]$ , and the X-ray crystal and molecular structure of the compound with  $\text{R}=\text{Ph}$ , **227**, 99
- structural, spectroscopic and potentiometric investigation on the coordination properties of 4- $\text{NO}_2$ -hippuric acid towards Cu(II) in binary and ternary systems, **227**, 105
- heteronuclear metal cluster complexes with a new coordination mode of the  $\text{Fe}_2\text{S}_2(\text{CO})_6$  unit. Synthesis, structure and possible reaction pathway of the first Fe–Cu–S cluster compound,  $[\text{Fe}_6\text{Cu}_3\text{S}_6(\text{CO})_{18}(\text{PPh}_3)_2]$ , **227**, 119
- preparation, structural and physical studies of mercury(II) halide complexes with the  $\text{N,N}'$ -chelating biheteroaromatic ligand 2-(2'-pyridyl)quinoxaline, **227**, 129
- synthesis, crystal structure and characterization of nickel(0) acrylonitrile complexes with a bidentate N-donor ligand, **227**, 137
- synthesis and structural analysis of *trans*- $\text{Cp}(\text{CO})_2$ - $(\text{PPh}_3)\text{WHgCl}$ , **227**, 145
- alkylation of 2-vinylpyridine and crystal structure of bis{(acetylacetonate)(1-pyridyl)butyl}nickel(II) complex, **227**, 149
- NQR studies of polyhalocuprate(I) anions, **227**, 153
- structures of a monomeric and a dimeric Ni(II) complex containing 2-acetylpyrrole, **227**, 159
- ten-vertex polyhedral azametallaborane chemistry. An *arachno*-6,9-azaplatinadecaborane and a *nido*-6,8-azaruthenadecaborane, **227**, 163
- the preparation, properties and crystal structure of a salt containing the hexabromochromate(III) anion, **227**, 167
- the synthesis and crystal structure of ethylenedinitrilotetra-acetamidoxime nickel(II) sulfate trihydrate, **227**, 171
- synthesis and X-ray structure of the dinuclear  $\mu$ -5,5'-diamino-3,3'-bis-1,2,4-triazolato(1-)-bis(diethylene-triamine)copper(II) triperchlorate,  $[\{\text{Cu}(\text{dien})_2\text{Hdabt}\}(\text{ClO}_4)_3]$ , **227**, 181
- a new type of face-sharing biocahedral triply bonded diniobium cluster  $(\text{PyH})[\text{Nb}_2(\mu\text{-Cl})_2(\mu\text{-THT})\text{Cl}_3(\text{Py})_3](\text{THF})\text{-}(\text{C}_6\text{H}_6)_{0.5}$ , made by a designed synthesis, **227**, 191
- structural characterization of bidentate carboxylate derivatives of copper(I) bistrisphenylphosphine, **227**, 223
- synthesis and experimental/theoretical investigation of the high-nuclearity cubic  $T_d$   $[\text{Au}_6\text{Ni}_{12}(\text{CO})_{24}]^{2-}$  cluster, an initial example of a discrete gold–nickel bimetallic-bonded species: comparative analysis of the results of electron-counting methods and the Fenske–Hall MO model in rationalizing the bonding interactions of its  $\text{Au}_6\text{Ni}_{12}$  core consisting of five face-fused metal octahedra, **227**, 269
- Cuboidal complexes
- electron transfer reactivity of cuboidal heterometallic  $\text{Mo}_3\text{MS}_4$  clusters in aqueous acidic solutions, **225**, 157
- Cyanate complexes
- $\{[\text{Ni}(3,2,3\text{-tet})(\mu\text{-NCO})_n](\text{ClO}_4)_n\}$ : structural characterization of the first monodimensional nickel(II)–cyanato system with a description of its ferromagnetic coupling, **216**, 5
- Cyanato complexes
- crystal structure and ligand field states of  $[\text{Cr}(\text{NH}_3)_5\text{NCO}](\text{NO}_3)_2$ , **221**, 93
- Cyanide complexes
- cobalt chloride and phase-transfer catalyzed carbonylation of benzyl chlorides, **216**, 97
- Cyanoalkyl complexes
- influence of pendant cyanoalkyl side-arms on the stabilities of metal ion complexes of monoaza- and diaza-18-crown-6 ethers in methanol, **219**, 31
- Cyano complexes
- interaction of cyanopentaamminechromium(III) with mercuric ion. Electrochemical, spectral and preparative study of adduct formation and linkage isomerism, **216**, 113
- synthesis and characterisation of cyanide bridged heterobinuclear complexes based on Group VI octacyanometalate(IV) and pentaammineosmium(III) ions: a comparison with Group VIII hexacyano analogues, **219**, 121
- a novel reaction of the cyano ligand opening an organometallic route to 4-amino imidazoles: four component condensation (4CC) with isocyanide, aldehyde and amine, **222**, 275
- properties of the mixed-valence binuclear complex ion,  $[(\text{NH}_3)_5\text{Ru}^{\text{III}}\text{-NC-Os}^{\text{II}}(\text{CN})_5]$ , **223**, 37
- the use of pressure-tuning spectroscopy to distinguish between one- and two-electron transfer processes, **225**, 89
- the construction of molecular systems capable of photoinduced multielectron charge transfer: a photochemical analysis of the reactivity of  $[(\text{CN})_5\text{Fe}^{\text{II}}\text{-(CN)-Pt}^{\text{IV}}(\text{L})_4\text{-}(\text{NC)-Fe}^{\text{II}}(\text{CN})_5]^{4-}$  ( $\text{L}=\text{NH}_3$ , ethylenediamine), **226**, 251
- Cyclic voltammetry
- electrocatalytic reduction of  $\text{CO}_2$  and  $\text{O}_2$  with electropolymerized films of vinyl-terpyridine complexes of Fe, Ni and Co, **225**, 285
- Cyclization
- electrophilic addition to styrylsilanes: sequential carbon–carbon bond forming reactions, **220**, 145
- Cycloaddition reaction
- preparation of  $\eta^1$ -1,3-dienyl,  $\eta^1$ -1,2-dienyl and  $\eta^3$ -cyclobutenone complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305
- Cycloaddition reactions
- a study of the reactivity and structure of cyclic  $\alpha,\beta$ -unsaturated Fischer-type carbene complexes, **220**, 233
- Cyclodextrin complexes
- the kinetics of electron-transfer reactions of the  $[\text{FeCp}(\text{CpCH}_2\text{N}(\text{CH}_3)_3)]^{+/2+}$  couple in the presence of cyclodextrins in aqueous media, **225**, 51
- Cyclohexadienyl complexes
- crystal and molecular structures of  $\{exo\text{- and }endo\text{-}(\text{RO})_2\text{P}(\text{O})\text{-}\eta^5\text{-C}_6\text{H}_6\}\text{Mn}(\text{CO})_3]$ , **224**, 147
- Cyclometallated complexes
- cyclometallation of *N,N*-dimethylbenzo[*b*]furan-2-carbothio (and seleno) amides with palladium(II), ruthenium(II) and rhodium(III), **215**, 91
- Cyclometallated ligand complexes
- cyclometallated derivatives of platinum(II) derived from 1,4-benzodiazepin-2-ones. Crystal and molecular structure of  $\text{Pt}(\text{L})(\text{HL})\text{Cl}$  ( $\text{HL}=\text{7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one}$ , DIAZEPAM), a molecule containing a neutral and a deprotonated 1,4-benzodiazepin-2-one, **215**, 17
- Cyclometallation
- synthesis of  $\eta^3$ -allyl–Pd complexes and subsequent formation of N-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- Cyclopalladate complexes
- cyclopalladated *p*-azoxyanisole complexes. Synthesis and structure of  $\{[\text{bis}(2\text{-phenylpyridine})][4'\text{-}(\text{methoxyphenyl})\text{-NNO}$

- azoxy-*N*<sup>2</sup>-4-methoxyphenyl-2-ato]palladium(II)}-(tetrafluoroborate), **218**, 211
- Cyclopentadienyl complexes
- NMR parameters as steric probes for arylcyclopentadienyl iron complexes, **215**, 139
- synthesis and fluxional behaviour of (cyclopentadienyl) and (pentamethylcyclopentadienyl)ruthenium(II) complexes with hemilabile ether-phosphines. <sup>31</sup>P DNMR spectroscopic studies and line shape analysis, **215**, 165
- two silicon-bridged cyclopentadienyl Fe<sub>2</sub>S<sub>4</sub> cluster units connected through an Fe(III)S<sub>6</sub> unit. Crystal and molecular structure of [(CH<sub>3</sub>)<sub>2</sub>SiCp<sub>2</sub>]<sub>2</sub>Fe<sub>5</sub>(μ<sub>3</sub>-S<sub>2</sub>)<sub>2</sub>(μ<sub>4</sub>-S<sub>2</sub>)(μ<sub>4</sub>-S<sub>4</sub>)FeCl<sub>4</sub>, **216**, 1
- <sup>1</sup>H NMR analysis of some η<sup>6</sup>-(substituted aryl phenylsulfonyl-acetonitrile)-η<sup>5</sup>-cyclopentadienylium cations, **219**, 221
- ruthenium complex catalyzed polymerization of OH or COOH group containing alkynes to give functionalized poly(acetylene)s, **220**, 35
- π-facial selectivities in the coordination of an optically pure fused cyclopentadienyl ligand to arene ruthenium moieties. NMR spectral and structural characterization of a single diastereomer of [(PCp)Ru(Me<sub>6</sub>C<sub>6</sub>)]PF<sub>6</sub>, **221**, 41
- an investigation of electron transfer properties of (Cp)Mn(CO)<sub>3</sub> and substituted Cp derivatives, **221**, 85
- synthesis and X-ray structural determination of pentamethylcyclopentadienyl dichloro oxo molybdenum, **221**, 147
- the reaction of CpRu(PPh<sub>3</sub>)<sub>2</sub><sup>+</sup> with trimethylenesulfide. Synthesis and molecular and crystal structure of {CpRu-(PPh<sub>3</sub>)<sub>2</sub>(SC<sub>3</sub>H<sub>6</sub>)]CF<sub>3</sub>SO<sub>3</sub>, **221**, 157
- carbon dioxide fixation by Cp<sub>2</sub>(η<sup>3</sup>-allyl)Ti complexes generated from various dienes, **222**, 145
- transition-metal silyl complexes 51. Preparation of the mixed silyl/stannyl complexes (π-C<sub>5</sub>H<sub>5</sub>)Fe(CO)(SnPh<sub>3</sub>)(SiR<sub>3</sub>)H, **224**, 177
- syntheses and structures of mixed-ligand tetrairon-sulfur clusters, [Cp'<sub>2</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)<sub>2</sub>Fe<sub>4</sub>S<sub>4</sub>](PF<sub>6</sub>) and [Cp'<sub>3</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)Fe<sub>4</sub>S<sub>5</sub>](PF<sub>6</sub>)<sub>n</sub> (n = 1 and 2) (Cp' = η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>). Structural change of the clusters accompanying electron transfer reactions, **225**, 229
- ligand substitution at 19-electron organometallic centers. Electrocatalytic CO substitution reactions of (methylcyclopentadienyl)Mn(CO)<sub>2</sub>NO<sup>+</sup> and (indenyl)Mn(CO)<sub>2</sub>NO, **226**, 53
- synthesis and structural analysis of *trans*-Cp(CO)<sub>2</sub>(PPh<sub>3</sub>)-WHgCl, **227**, 145
- Cyclopentadienylruthenium complexes
- synthesis, molecular structure and electrochemical investigation of the 1,2,5-thiadiazole-3,4-dithiolate-substituted complex Cp<sup>\*</sup>Ru(NO)(tdas). Evidence for nitrosyl bending during reduction, **226**, 267
- Cyclopropanation
- dirhodium(II) tetrakis[*N,N*-dimethyl-2-pyrrolidone-5(*S*)-carboxamide]. Structural effects on enantioselection in metal carbene transformations, **220**, 193
- bicyclic carbo- and heterocycles from an allylidene complex and cyclic 1,3-dienes by tandem cyclopropanation/Cope rearrangement, **220**, 327
- a one-pot synthesis of cyclopropanes from aromatic aldehydes and styrene via iron carbene complexes, **223**, 159
- Cytochrome *c*
- volume profile analysis for intermolecular electron transfer between cytochrome *c* and Co(terpy)<sub>2</sub>, **225**, 95
- intramolecular electron transfer between Ru(I) and Ru(III) and the heme iron of cytochrome *c* labeled with ruthenium(II) polypyridine complexes, **226**, 129
- Decarboxylation
- the steric effect on the ring opening process of the decarboxylation of *cis*-carbonato-bis(diamine)cobalt(III) ions, **221**, 9
- Depalladation
- synthesis of η<sup>3</sup>-allyl-Pd complexes and subsequent formation of N-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- Dextran complexes
- Mössbauer spectroscopy of iron dextran: comparison of solid and frozen solution, **223**, 9
- Dialkyl complexes
- organoplatinum(II) complexes containing disubstituted dithioamides: evidence for an S,S' Pt coordinated neutral dithioxamide acting as an anion binding agent, **227**, 63
- Diamine complexes
- the steric effect on the ring opening process of the decarboxylation of *cis*-carbonato-bis(diamine)cobalt(III) ions, **221**, 9
- a potentiometric study on the complexation of Ag(I) with various N-methyl and C-methyl substituted ethylenediamines in 1 M NO<sub>3</sub>, **224**, 97
- equilibrium studies of 1,2-diaminobenzene and 3,4-diaminobenzoate anion with dinuclear cobalt(II) 1,4,7,13,16,19-hexaaza-10,22-dioxacyclotetacosane, **227**, 49
- Diazoalkane complexes
- reactions of [Au(C<sub>6</sub>F<sub>5</sub>)(SC<sub>4</sub>H<sub>8</sub>)] with diazoalkanes. Synthesis and molecular structures of [Au(C<sub>6</sub>F<sub>5</sub>)(Ph<sub>2</sub>C=N=N=CPh<sub>2</sub>)], **222**, 267
- Dicarbaborane complexes
- ten-vertex polyhedral dicarbaborane chemistry: new positional isomers of cluster constituents in the ten-vertex *arachno* dicarbaborane family: the parent [*arachno*-5,10-C<sub>2</sub>B<sub>8</sub>H<sub>13</sub>]<sup>-</sup> anion and the isomeric ligand derivatives *exo*-9-*L*-*arachno*-5,6-C<sub>2</sub>B<sub>8</sub>H<sub>12</sub> and *exo*-6-*L*-*arachno*-5,10-C<sub>2</sub>B<sub>8</sub>H<sub>12</sub>, **221**, 5
- Dicyanamido complexes
- solvent trapped valency in the [bis(pentaammineruthenium)-(μ-1,4-dicyanamidobenzene)] trication, **225**, 123
- Diels-Alder reactions
- manganese and tungsten carbene complex mediated intramolecular Diels-Alder reactions, **222**, 291
- Diene
- preparation of η<sup>1</sup>-1,3-dienyl, η<sup>1</sup>-1,2-dienyl and η<sup>3</sup>-cyclobutene complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305
- Diene complexes
- addition of carbanions to tris(trimethylphosphine)-(cyclooctadiene)iridium(I) chloride: three different modes of reaction depending on the nature of the carbanion, **222**, 135
- intermediates in the palladium-catalysed reaction of 1,3-dienes Part 8. The reaction of palladium-butadiene complexes with ethyl methylacetoacetate, **222**, 179
- electrochemical behavior and X-ray crystal structure of the unusual dimeric ruthenium(cyclooctadiene) complex, (tri-μ-chloro)(chloro)(acetonitrile)bis(η<sup>4</sup>-cyclooctadiene)-diruthenium(II), **224**, 125
- Diimine complexes
- light induced exchange energy transfer reactions of micelle forming Ru(II) diimine complexes, **225**, 275
- photoinduced intramolecular electron transfer in bimetallic donor-acceptor assemblies, **226**, 79
- β-Diketonate complexes
- a novel route to the second-generation alkaline-earth metal precursors for metal-organic chemical vapour deposition: one-step synthesis of M(hfa)<sub>2</sub> tetraglyme (M = Ba, Sr, Ca and Hhfa = 1,1,1,5,5,5-hexafluoro-2,4-pentanedione), **224**, 203
- Dimeric complexes

- structures of a monomeric and a dimeric Ni(II) complex containing 2-acetylpyrrole, **227**, 159
- Dimerization**
- regio- and stereoselective dimerization of 1-alkynes catalyzed by an Os(II) complex, **220**, 5
- Dimethyl sulfoxide complexes**
- metal complexes of benzodiazepines. Kinetic properties of bromazepam toward square planar dimethyl sulfoxide complexes of platinum(II) and palladium(II), **215**, 41
- Dinuclear complexes**
- preparation and structural characterization of  $\text{Rh}_2(\text{O}_2\text{CCPh}_3)_4(\text{EtOH})_2$ ,  $\text{Ru}_2(\text{O}_2\text{CCPh}_3)_4(\text{H}_2\text{O})(\text{EtOH}) \cdot 2\text{EtOH}$  and  $\text{Mo}_2(\text{O}_2\text{CCPh}_3)_4 \cdot 3\text{CH}_2\text{Cl}_2$ , **215**, 9
- preparation and characterization of  $[(\text{LL})\text{Cl}_2\text{Mo}]_2(\mu\text{-SPr})_2$  compounds with Pr = n-propyl and LL = PrSCH<sub>2</sub>CH<sub>2</sub>SPr and Et<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PEt<sub>2</sub>. Further examples of C–S bond cleavage and oxidative addition across an Mo–Mo quadruple bond, **215**, 191
- pyrazine-mediated antiferromagnetic intermolecular exchange in mixed-valent diruthenium tetracarboxylates, **215**, 203
- a tris-(μ<sub>2</sub>-I) face-sharing dizirconium(III) complex,  $[\text{Zr}_2\text{I}_7(\text{PEt}_3)_2]^-$ ; the first such dizirconium species, **216**, 9
- a polymeric two-dimensional mixed-metal network. Crystal structure and magnetic properties of  $\{[\text{P}(\text{Ph})_4]_n[\text{MnCr}(\text{ox})_3]_n\}$ , **216**, 65
- dirhenium(III) complexes with *N,N*-dimethyladenine and 7-azaindole, **216**, 131
- octahedral μ-oxalato-nickel(II) dinuclear complexes with water and tridentate amines as blocking ligands: magnetostructural correlations, **216**, 139
- crystal structure, magnetic and spectroscopic properties of the bis(dimethyl sulfoxide) adduct of tetra-μ-formato-dicopper(II), a new tetracarboxylato-bridged copper(II) dimer, **216**, 185
- geometrical and optical isomers of the dinickel complexes of two chiral ligands of the *N,N'*-disubstituted oxamide type: 1D and 2D NMR study, **216**, 237
- synthesis, characterization and crystal structure of  $(\text{C}_3\text{H}_5\text{N}_2)_2[\text{Sn}_2(\text{NO}_3)_4(\mu\text{-OH})_2\text{Me}_4]$ , a di-μ-hydroxo-bridged compound with heptacoordinated tin(IV), **216**, 257
- electrochemical investigation of the electron-poor/precise (*n* = 5/6) complexes  $[\text{Fe}_2(\text{CO})_n(\mu\text{-PR}_2)(\mu\text{-PR}'_2)]$  (*n* = 5, R = R' = Bu<sup>t</sup>; *n* = 6, R = R' = Ph; R = Bu<sup>t</sup>, R' = Ph; R = Bu<sup>t</sup>, R' = Cy). EPR study of the radical anion  $[\text{Fe}_2(\mu\text{-PBu}^t)_2(\text{CO})_5]$ , **217**, 29
- magnetic properties and crystal structure of {μ-pyrazine}-bis[aqua](2,6-xylylamino)diacetato)copper(II)]. A pyrazine-bridged dinuclear copper(II) complex, **217**, 39
- syntheses of dinuclear gold(I) ring complexes containing 1,1-dicyanoethene-2,2-dithiolate-*S,S* and bis(diphosphines) as bridging ligands. X-ray crystal structure of  $[\text{Au}_2\mu\text{-}(i\text{-MNT})\mu\text{-}(dppee)]$ , **217**, 45
- a mixed-metal salt comprised of metal–metal bonded dinuclear ions: structure and properties of  $[\text{Rh}_2(\text{OAc})_2(\text{MeCN})_6][\text{Re}_2\text{Cl}_8]$ , **217**, 79
- synthesis and characterization of Cd(II) complexes of cinoxacin. Crystal structure of a dicadmium complex containing two monoatomic-bridging carboxylate oxygen atoms, **217**, 149
- insertion reaction of dimethylacetylenedicarboxylate with the head-to-tail isomer of  $\text{Pt}_2\text{I}_2(\mu\text{-Ppy}_3)_2$ ; Ppy<sub>3</sub> = tris(2-pyridyl)phosphine, **217**, 209
- preparation, crystal structures and magnetic properties of a *syn-anti* carboxylate-bridged dinuclear copper(II) complex and its precursor, a hydrogen-bonded polynuclear copper(II) complex, **218**, 109
- preparation and crystal structure of *anti*-(η<sup>3</sup>:η<sup>5</sup>-azulene)-pentakis(trimethylphosphine)dicobalt. Reactions with carbon monoxide and solution dynamics, **218**, 143
- coordination chemistry of perhalogenated cyclopentadienes and alkynes XIV. Synthesis of dinuclear ring-bridged cyanotrenes with –SiMe<sub>2</sub>– or –TiCp<sub>2</sub>– bridges. Structures of  $[(\text{OC})_3\text{Mn}(\text{C}_5\text{Br}_4)]_2\text{SiMe}_2$  and  $[(\text{OC})_3\text{Mn}(\text{C}_5\text{Cl}_4)]_2\text{TiCp}_2$ , **218**, 173
- orthometalation reactions in trifluoroacetate dirhodium(II) compounds. Molecular structure of  $\text{Rh}_2(\text{O}_2\text{CCF}_3)_2 \cdot [(\text{C}_6\text{H}_4)\text{PPh}_2]_2 \cdot (\text{PPh}_3)_2 \cdot 2(\text{C}_7\text{H}_8)$ , **218**, 189
- a new class of dinuclear compounds: the synthesis and X-ray structural characterization of tris(μ-diphenylformamido)diiron, **219**, 7
- dimolybdenum(II) and diruthenium(II) complexes with 2-substituted 7-methyl-1,8-naphthyridines as bridging ligands, **219**, 23
- variation of the exchange interaction in oximate-bridged Cu<sup>II</sup>M<sup>II</sup> dimers (M = Cu, Ni, Mn). Crystal structure of  $[\text{Cu}(\text{pdmg})\text{Cu}(\text{bipy})(\text{H}_2\text{O})_2](\text{ClO}_4)_2 \cdot \text{H}_2\text{O}$ , **219**, 179
- dirhodium(II) tetrakis[*N,N*-dimethyl-2-pyrrolidone-5(*S*)-carboxamide]. Structural effects on enantioselection in metal carbene transformations, **220**, 193
- reactivity of allene at phosphine-bridged di-iron centres: X-ray crystal structures of  $[\text{Fe}_2(\text{CO})_5\{\mu\text{-}\sigma\text{-C}(\text{O})\text{C}(\text{CH}_2)_2\}(\mu\text{-dppm})]$  and  $[\text{Fe}_2(\text{CO})_4\{\mu\text{-}\eta^3\text{-}\eta^3\text{-}(\text{CH}_2)_2\text{C}_2(\text{CH}_2)_2\}(\mu\text{-dppm})] \cdot \text{Et}_2\text{O}$ , **220**, 201
- synthesis and structural characterization of a new series of ruthenium(II) complexes containing the short bite Ph<sub>2</sub>PPy ligand. Cooperative effect between the anionic rhodium and cationic ruthenium species in the catalytic hydroformylation of styrene by  $[\text{Ru}(\text{Ph}_2\text{PPy})_3\text{Cl}][\text{Rh}(\text{CO})_2\text{Cl}_2]$ , **221**, 109
- reaction of primary silanes with dinuclear rhodium hydride complexes: silane coupling reactions, **222**, 345
- synthesis and reactions of the novel sulfonium-methylidene complex  $[\text{Fe}_2(\text{CO})_2(\text{cp})_2(\mu\text{-CO})\{\mu\text{-C}(\text{H})(\text{SMe}_2)\}]\text{SO}_3\text{CF}_3$ , **223**, 31
- synthesis and X-ray analysis of the dirhodium(II,II) carboxylate complex  $\text{Rh}_2(\text{O}_2\text{CCH}_3)_4(\text{formH})_2$  (formH = *N,N'*-di-*p*-tolylformamidine), **223**, 145
- divalent iron formamidinato complexes: a highly distorted dinuclear compound, **224**, 5
- di-manganese carbonyl compounds containing bis(dicyclohexylphosphino)methane (dcpm) ligation, **224**, 105
- electrochemical behavior and X-ray crystal structure of the unusual dimeric ruthenium(cyclooctadiene) complex, (tri-μ-chloro)(chloro)(acetonitrile)bis(η<sup>4</sup>-cyclooctadiene)diruthenium(II), **224**, 125
- crystal structure of a binuclear vanadium(III) complex with a new tripodal ligand,  $[\text{V}_2\text{Cl}_4(\text{tped})(\text{EtOH})_2] \cdot 2\text{EtOH}$ , **224**, 199
- synthesis and electrochemical properties of oxo-bridged manganese dimers incorporating alkali and alkaline earth cations, **225**, 191
- metal–metal interactions of aqueous lead(II) and tin(II) ions with the ground and excited states of tetrakis(μ-diphosphito)diplatin(II), **225**, 201
- synthesis, electrochemistry and photoexcited-state properties of dinuclear ruthenium complexes bridged by 2,6'-bis(2-pyridyl)-2,2':6,2''-thiazolo[4,5-*d*]-benzothiazole, **226**, 17
- solvent, electrolyte and solute shape effects on optical electron transfer in mixed-valence ruthenium ammine dimers, **226**, 137
- synthesis of some dinuclear niobium chloro alkoxy carboxylates  $[\text{Nb}_2\text{Cl}_4(\text{OEt})_4(\text{O}_2\text{CR})_2]$ , and the X-ray crystal and molecular structure of the compound with R = Ph, **227**, 99

- preparation, structural and physical studies of mercury(II) halide complexes with the  $N,N'$ -chelating biheteroaromatic ligand 2-(2'-pyridyl)quinoxaline, **227**, 129
- synthesis and X-ray structure of the dinuclear  $\mu$ -5,5'-diamino-3,3'-bis-1,2,4-triazolato(1-)-bis(diethylenetriamine-copper(II)) triperchlorate,  $[\{\text{Cu}(\text{dien})\}_2\text{Hdabt}](\text{ClO}_4)_3$ , **227**, 181
- a new type of face-sharing bioctahedral triply bonded dinobium cluster  $(\text{PyH})[\text{Nb}_2(\mu\text{-Cl})_2(\mu\text{-THT})\text{Cl}_3(\text{Py})_3](\text{THF})(\text{C}_6\text{H}_6)_{0.5}$ , made by a designed synthesis, **227**, 191
- Dinucleating ligand complexes
- geometrical and optical isomers of the dinickel complexes of two chiral ligands of the  $N,N'$ -disubstituted oxamide type: 1D and 2D NMR study, **216**, 237
- Dioxygen complexes
- preparation and dioxygen binding property of a new zeolite encapsulated cobalt tetramethylporphyrin complex, **217**, 135
- synthesis, structural characterization and dioxygen reactivity of imidazole-ligated Cu(I) complexes, **218**, 103
- Dioxygen reduction
- electrocatalysts for the four-electron reduction of dioxygen based on adsorbed cobalt tetrapyrrolylporphyrin molecules linked by aquaammine complexes of ruthenium(II), **225**, 215
- Diphosphine complexes
- syntheses of dinuclear gold(I) ring complexes containing 1,1-dicyanoethene-2,2-dithiolate- $S,S$  and bis(diphosphines) as bridging ligands. X-ray crystal structure of  $[\text{Au}_2\mu\text{-}(i\text{-MNT})\mu\text{-}(dppee)]$ , **217**, 45
- bite angle dependence of the rate of reductive elimination from diphosphine palladium complexes, **220**, 249
- palladium complex catalysed synthesis of sulfinic acids, sulfinic acid esters, sulfonic acids and  $S$ -alkyl alkanethiosulfonates, **222**, 381
- Diphosphito complexes
- metal–metal interactions of aqueous lead(II) and tin(II) ions with the ground and excited states of tetrakis( $\mu$ -diphosphito)diplatinate(II), **225**, 201
- Dithiocarbamate complexes
- methyl dithiocarbamate gold(I) and gold(III) complexes. Synthesis and reactivity with amines, **215**, 215
- dithiocarbamates of dimethylthallium(III). Crystal structure of dimethyl(di- $n$ -propyldithiocarbamato)thallium(III), **216**, 15
- spectroscopic investigation of dichloro and hydrazido(2-)-derivatives of bis(dithiocarbamato)dioxomolybdenum(VI) compounds. Crystal and molecular structure of bis(diisobutyl dithiocarbamato)( $N,N$ -dimethylhydrazido(2-)- $N'$ -oxomolybdenum(VI)), **217**, 61
- Dithiocarbamate ligand
- investigation of the dithiocarbamate-induced coupling of allylidene and carbonyl ligands on a tungsten center, **220**, 335
- Dithiocarbamato complexes
- di- and tetranuclear gold(II) complexes with dithiocarbamate and related ligands. X-ray structure of  $[\text{Au}_2(\mu\text{-CH}_2\text{PPh}_2\text{CH}_2)_2(\text{S}_2\text{CNMe}_2)_2]$ , **223**, 55
- Dithiocarboxylic acid derived ligand
- synthesis and characterization of neutral oxorhenium(V) and nitridotechnetium(V) complexes with a tetradentate  $\text{N}_2\text{S}_2$  unsaturated ligand derived from dithiocarboxylic acid, **227**, 25
- Dithiolate complexes
- syntheses of dinuclear gold(I) ring complexes containing 1,1-dicyanoethene-2,2-dithiolate- $S,S$  and bis(diphosphines) as bridging ligands. X-ray crystal structure of  $[\text{Au}_2\mu\text{-}(i\text{-MNT})\mu\text{-}(dppee)]$ , **217**, 45
- Dithiolato complexes
- synthesis, characterization and X-ray crystal structure of the benzenedithiolatohafnium(IV) porphyrin complex, **221**, 51
- Dithiolene complexes
- preparation and properties of a metal dithiolene conductor, partially oxidized salt of tetra( $n$ -butyl)ammonium bis(5,6-dithio-benzo[ $d$ ]-1,3-dithiole-2-thione) nickelate,  $(n\text{-Bu}_4\text{N})_{0.29}[\text{Ni}(\text{dmbit})_2]$ , **216**, 147
- Dithiophosphinate complexes
- ionic cluster dithiophosphinates  $[\text{Mo}_3\text{S}_4(\text{R}_2\text{PS}_2)_3\text{py}_3]^+[\text{Ml}_3]^-$  ( $\text{M} = \text{Zn}\cdot\text{py}$ ,  $\text{Cd}\cdot\text{py}$ ,  $\text{Hg}$ ), **216**, 177
- Dithioamide complexes
- organoplatinum(II) complexes containing disubstituted dithioamides: evidence for an  $S,S'$  Pt coordinated neutral dithioamide acting as an anion binding agent, **227**, 63
- Divalent ion complexes
- stabilities of 1,2-diethyl-3-hydroxy-4-pyridinone chelates of divalent and trivalent metal ions, **223**, 21
- Divinylcyclopropane complexes
- bicyclic carbo- and heterocycles from an allylidene complex and cyclic 1,3-dienes by tandem cyclopropanation/Cope rearrangement, **220**, 327
- $\pi$ - $\sigma$  Donor–acceptor electronic coupling
- forbidden  $\pi$ - $\sigma$  donor–acceptor electronic coupling in linked transition metal complexes: contrasting behavior of ruthenium(II) donors with ruthenium(III) and cobalt(III) acceptors, **226**, 109
- Double complex salt
- electron transfer quenching of the excited MLCT state of Ru(II) in the double complex salts  $[\text{Ru}(\text{bpy})_3]_2\text{-}[\text{M}(\text{CN})_6]\text{Cl}\cdot 8\text{H}_2\text{O}$  ( $\text{M} = \text{Co}$  and  $\text{Fe}$ ), **226**, 203
- $\beta$ -Effect
- electrophilic addition to styrylsilanes: sequential carbon–carbon bond forming reactions, **220**, 145
- Electric properties
- preparation and properties of a metal dithiolene conductor, partially oxidized salt of tetra( $n$ -butyl)ammonium bis(5,6-dithio-benzo[ $d$ ]-1,3-dithiole-2-thione) nickelate,  $(n\text{-Bu}_4\text{N})_{0.29}[\text{Ni}(\text{dmbit})_2]$ , **216**, 147
- Electrocatalysis
- electrocatalysts for the four-electron reduction of dioxygen based on adsorbed cobalt tetrapyrrolylporphyrin molecules linked by aquaammine complexes of ruthenium(II), **225**, 215
- electrocatalytic reduction of  $\text{CO}_2$  and  $\text{O}_2$  with electropolymerized films of vinyl-terpyridine complexes of Fe, Ni and Co, **225**, 285
- ligand substitution at 19-electron organometallic centers. Electrocatalytic CO substitution reactions of (methylcyclopentadienyl) $\text{Mn}(\text{CO})_2\text{NO}^+$  and (indenyl) $\text{Mn}(\text{CO})_2\text{NO}$ , **226**, 53
- Electrochemical Peltier heat
- thermometric determination of electrochemical Peltier heat (thermal effect associated with electron transfer) of some redox couples, **226**, 69
- Electrochemistry
- redox reactions of an Mo(V) tetrasulfophthalocyanine, **215**, 27
- synthesis, crystal and molecular structures, UV–Vis spectroscopy and electrochemical properties of two iron(III) phenolate complexes, **215**, 123
- interaction of cyanopentaamminechromium(III) with mercuric ion. Electrochemical, spectral and preparative study of adduct formation and linkage isomerism, **216**, 113
- electrochemical investigation of the electron-poor/precise ( $n = 5/6$ ) complexes  $[\text{Fe}_2(\text{CO})_n(\mu\text{-PR}_2)(\mu\text{-PR}'_2)]$  ( $n = 5$ ,  $\text{R} = \text{R}' = \text{Bu}^t$ ;  $n = 6$ ,  $\text{R} = \text{R}' = \text{Ph}$ ;  $\text{R} = \text{Bu}^t$ ,  $\text{R}' = \text{Ph}$ ;  $\text{R} = \text{Bu}^t$ ,  $\text{R}' = \text{Cy}$ ). EPR study of the radical anion  $[\text{Fe}_2(\mu\text{-P}^t\text{Bu}_2)_2(\text{CO})_5]$ , **217**, 29

- synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129
- heterodinuclear Ni(II)M(II) (M=Pb, Mn, Fe, Co, Ni, Cu, Zn) complexes of a phenol-based dinucleating macrocycle with dissimilar 4- and 5-coordination sites, **218**, 27
- electrochemistry of organometallic halide complexes III. Oxidation of  $[C_6Me_6RuCl_2]_2$ , **218**, 33
- estrogen derivatives of transition metal complexes for analytical detection enhancement. Part II, **218**, 207
- electrochemical and structural studies of nickel(II) complexes with  $N_2O_2$  Schiff base ligands 2. Crystal and molecular structure of *N,N'*-1,2-ethane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II), *N,N'*-1,2-*cis*-cyclohexane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II) and *N,N'*-1,2-benzene-1,2-diyl-bis(3,5-dichlorosalicylideneimine)nickel(II), **219**, 43
- ruthenium chromophores containing terpyridine and a series of polyazine bridging ligands, **221**, 55
- molybdenum complexes with sterically-hindered thio-carboxylate ligands. Electrochemical properties of the anionic complex bis(2,2-diphenyl-2-mercaptoethanoate)dioxomolybdate(VI) in protic solvents, **221**, 117
- direct electrochemical synthesis of tin(IV) complexes of 2-(2-mercaptoethyl)iminomethyl-phenols; the molecular structure of bis{2-(2-mercaptoethyl)iminomethyl-5-bromo-phenoxy}tin(IV), **223**, 87
- synthesis, spectra and electrochemical behaviour of biomimetic copper(II) complexes with  $CuN_5$  and  $CuN_6$  chromophores, **224**, 19
- the synthesis, and UV-Vis, electrochemical, photochemical and intervalence characterization of tetraammine and tris chelated ruthenium(II) complexes bound to the bridging ligand dipyrido(2,3-*a*;2',3'-*h*)phenazine (dpop), **224**, 27
- reactions of  $HCCO_3(CO)_9$  with silanes; synthesis and electrochemistry of  $X[SiMe_2CCO_3(CO)_9]_2$  (X=O, 1,4- $C_6H_4$ ), **224**, 73
- electrochemical behavior and X-ray crystal structure of the unusual dimeric ruthenium(cyclooctadiene) complex, (tri- $\mu$ -chloro)(chloro)(acetonitrile)bis( $\eta^4$ -cyclooctadiene)diruthenium(II), **224**, 125
- multinuclear ( $^{31}P$ ,  $^{77}Se$ ,  $^{113}Cd$ ,  $^{199}Hg$ ) magnetic resonance, electrospray mass spectrometric and electrochemical studies on the reactions of  $M(S_2P\{OEt\}_2)_2$  (M=Cd, Hg) with potentially bidentate mixed Group 15/Group 16 and Group 16 donor ligands, **224**, 137
- electron transfer in *trans*-[Pt(PPh<sub>3</sub>)<sub>2</sub>(-C≡C-Fc)<sub>2</sub>] and related compounds, **225**, 35
- electrochemistry of nitrite reductase model compounds 5. Electrochemistry and spectroelectrochemistry of iron porphyrone, porphindione and isobacteriochlorin complexes, **225**, 57
- electrochemical, photophysical and photochemical properties of a hexanuclear ruthenium(II) bis(2,2'-bipyridyl) complex containing a tris(pyrazyltriazole)benzene ligand, **225**, 67
- ligand nature of coordinated  $NO_2^-$  in the oxidation reaction of Ru(II) complexes,  $[Ru(NO_2)(OH_2)(py)_{4-2n}(bpy)_n]^+$  (n=0, 1, 2), and their related complexes, **225**, 111
- solvent trapped valency in the [bis(pentaammineruthenium)-( $\mu$ -1,4-dicyanamidobenzene)] trication, **225**, 123
- electrochemical recognition of Group 1 and 2 metal cations by redox-active ionophores, **225**, 137
- synthesis and electrochemical properties of oxo-bridged manganese dimers incorporating alkali and alkaline earth cations, **225**, 191
- syntheses and structures of mixed-ligand tetrairon-sulfur clusters,  $[Cp'_2(Ph_2C_2S_2)_2Fe_4S_4](PF_6)$  and  $[Cp'_3(Ph_2C_2S_2)Fe_4S_3](PF_6)_n$  (n=1 and 2) (Cp' =  $\eta^5$ - $C_5Me_5$ ). Structural change of the clusters accompanying electron transfer reactions, **225**, 229
- controlling the acidity of the carboxylic group by a ferrocene based redox switch, **225**, 239
- the redox behaviour of the cluster anion  $[Fe_4N(CO)_{12}]^-$ . Electron transfer chain catalytic substitution reactions. Crystal structure of  $[Fe_4N(CO)_{11}PPh_3]$ , **226**, 1
- electron-transfer activation of the aminocarbyne and the hydrogen isocyanide complexes *trans*-[ReCl(CNH<sub>n</sub>)-(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>][BF<sub>4</sub>]<sub>n-1</sub> (n=2 or 1). Interconversion of coordinated CNH<sub>2</sub> and CNH, **226**, 9
- electron transfer in the electrochemistry of metals, metal compounds and metal complexes, **226**, 99
- quenching of a polypyridyl-ruthenium(II) chromophore by covalently attached {ML(NO)Cl} fragments (M=Mo, W; L=tris(3,5-dimethylpyrazolyl)hydroborate), **226**, 171
- reaction of low-valent iron porphyrins with alkyl containing supporting electrolytes, **226**, 195
- chemical and electrochemical approaches to the investigation of redox reactions of simple electron transfer metalloproteins, **226**, 293
- redox behavior of the iron-sulfur cluster  $[Fe_4Cp_4S_5][PF_6]_2$  in protic organic solvents and aqueous micellar solutions, **226**, 341
- Electrode
- electron transfer in surfactant films on electrodes; copper phthalocyaninetetrasulfonate-didodecyltrimethylammonium bromide, **226**, 159
- Electronic delocalization
- the electronic structure of pentaammine(pyrazine) ruthenium(II) and (III): the metal-ligand  $\pi$  conjugation and its implications in electron delocalization, **226**, 43
- Electronic excitation spectroscopy
- spectroscopic investigation of mixed-ligand nickel(II) chelates containing  $\beta$ -ketoenols and 1,2-diamines, **221**, 17
- Electronic interactions
- inter-chromophore electronic interactions in ligand-bridged polynuclear complexes: a comparative study of various bridging ligands, **226**, 213
- Electronic structures
- crystal structure and ligand field states of  $[Cr(NH_3)_5NCO](NO_3)_2$ , **221**, 93
- the electronic structure of pentaammine(pyrazine) ruthenium(II) and (III): the metal-ligand  $\pi$  conjugation and its implications in electron delocalization, **226**, 43
- Electron-localized states
- the influence of spin-forbidden monomer excitations on spin-allowed electron transfer and electron-localized states of mixed-valence and single-valence dimeric systems, **226**, 33
- Electron transfer
- stereoselectivity in reactions of metal complexes Part XVIII. Kinetics and stereoselectivity in electron-transfer reactions between  $[Co(L)H_2O]^+$  (L=*N,N'*-[(pyridine-2,6-diyl)bis(methylene)]bis[proline] (promp) and *N,N'*-[(4-methoxy-pyridine-2,6-diyl)bis(methylene)]bis[proline] (MeO-promp) and optically active iron(II) complexes, **218**, 73
- an investigation of electron transfer properties of (Cp)Mn(CO)<sub>3</sub> and substituted Cp derivatives, **221**, 85
- effect of substituting sulfur for oxygen on the heterogeneous electron transfer kinetics of oxomolybdenum(V) hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes, **225**, 1
- dependence of electron transfer kinetics of the ferrocene/ferrocenium couple on the viscosity in ambient temperature chloroaluminate ionic liquids, **225**, 9

- charge and electron transfer from metal-to-carbon bonds of main group organometallics  $MR_n$  ( $M = Al, Ga, Zn$ ) to aromatic N-heterocycles: colored precursor compounds and radical complex formation, **225, 23**
- electron transfer in *trans*-[Pt(PPh<sub>3</sub>)<sub>2</sub>(-C≡C-Fc)<sub>2</sub>] and related compounds, **225, 35**
- the kinetics of electron-transfer reactions of the [FeCp(CpCH<sub>2</sub>N(CH<sub>3</sub>)<sub>3</sub>)<sup>+2+</sup>] couple in the presence of cyclodextrins in aqueous media, **225, 51**
- the oxidation of [Co(edta)]<sup>2-</sup> by [Co(phen)<sub>3</sub>], **225, 75**
- electron transfer 122. Reductions of an N<sub>4</sub>-coordinated complex of silver(III), **225, 83**
- the use of pressure-tuning spectroscopy to distinguish between one- and two-electron transfer processes, **225, 89**
- volume profile analysis for intermolecular electron transfer between cytochrome *c* and Co(terpy)<sub>2</sub>, **225, 95**
- stereoselective electron transfer between Cu(II) superoxide dismutase and  $\Lambda$ - and  $\Delta$ -[Fe(pdta)], **225, 151**
- electron transfer reactivity of cuboidal heterometallic Mo<sub>3</sub>MS<sub>4</sub> clusters in aqueous acidic solutions, **225, 157**
- comparison of the rates of substitution in [Ru(NH<sub>3</sub>)<sub>5</sub>H<sub>2</sub>O]<sup>3+</sup> and [Os(NH<sub>3</sub>)<sub>5</sub>H<sub>2</sub>O]<sup>3+</sup> by hexacyano complexes: substitution coupled to electron transfer, **225, 173**
- indium(III) halide-3,5-di-*tert*-butyl-*o*-benzosemiquinone systems, **225, 207**
- syntheses and structures of mixed-ligand tetrairon-sulfur clusters, [Cp'<sub>2</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)<sub>2</sub>Fe<sub>4</sub>S<sub>4</sub>](PF<sub>6</sub>) and [Cp'<sub>3</sub>(Ph<sub>2</sub>C<sub>2</sub>S<sub>2</sub>)Fe<sub>4</sub>S<sub>4</sub>](PF<sub>6</sub>)<sub>*n*</sub> ( $n = 1$  and  $2$ ) (Cp' =  $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>). Structural change of the clusters accompanying electron transfer reactions, **225, 229**
- electron transfer 120. Reductions of the cuboidal mixed-valence molybdenum/sulfur cluster, [Mo<sub>4</sub>S<sub>4</sub>(H<sub>2</sub>O)<sub>12</sub>], **225, 297**
- electron transfer properties of ReO(ECD), **225, 323**
- infrared spectroelectrochemistry of [Rh(TM4)<sub>4</sub>M(CO)<sub>5</sub>]<sub>2</sub><sup>2+</sup> (TM4 = 2,5-diiisocyno-2,5-dimethylhexane; M = Re, Mn): electron-transfer-promoted metal-metal bond cleavage and formation, **226, 61**
- electron transfer in the electrochemistry of metals, metal compounds and metal complexes, **226, 99**
- electron transfer reduction of cobalt tetraphenylporphyrin by hydroquinone dianions and alkylation with alkyl halides, **226, 145**
- electron transfer in surfactant films on electrodes; copper phthalocyaninetetrasulfonate-didodecyldimethylammonium bromide, **226, 159**
- chemical and electrochemical approaches to the investigation of redox reactions of simple electron transfer metalloproteins, **226, 293**
- Electron transfer activation**
- electron-transfer activation of the aminocarbyne and the hydrogen isocyanide complexes *trans*-[ReCl(CNH<sub>*n*</sub>)-(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>][BF<sub>4</sub>]<sub>*n-1*</sub> ( $n = 2$  or  $1$ ). Interconversion of coordinated CNH<sub>2</sub> and CNH, **226, 9**
- Electron transfer catalyst**
- use of aluminium as an electron pool. Reductive coupling of iodobenzenes promoted by electron transfer in an Ni/Pb/Al multi redox system, **222, 323**
- Electron transfer process**
- light-induced electron transfer in lead(II)gold(I) dicyanide, **226, 345**
- Electron transfer reactions**
- relationship between four-state delineation and two-state delineation in electron transfer reactions and excitation transfer, **225, 185**
- basis set effects on the direct calculation of intermolecular coupling elements for electron transfer reactions, **226, 237**
- direct solution calorimetric measurements of enthalpies of proton and electron transfer reactions for transition metal complexes. Thermochemical study of metal-hydride and metal-metal bond energies, **227, 285**
- Electrophilic addition**
- electrophilic addition to styrylsilanes: sequential carbon-carbon bond forming reactions, **220, 145**
- Electrospray mass spectrometry**
- $\pi$ -bonded alkene and arene complexes of silver(I): an electrospray mass spectrometric study, **220, 99**
- positive and negative ion electrospray mass spectrometric studies of some amino acids and glutathione, and their interactions with alkali metal ions and methylmercury(II), **223, 103**
- multinuclear (<sup>31</sup>P, <sup>77</sup>Se, <sup>113</sup>Cd, <sup>199</sup>Hg) magnetic resonance, electrospray mass spectrometric and electrochemical studies on the reactions of M(S<sub>2</sub>P{OEt}<sub>2</sub>)<sub>2</sub> (M = Cd, Hg) with potentially bidentate mixed Group 15/Group 16 and Group 16 donor ligands, **224, 137**
- $\beta$ -Elimination**
- volumes of activation for  $\beta$ -elimination and related reactions of chromium(III)-alkyl complexes in aqueous solutions, **227, 57**
- Enamines**
- synthesis of stable tail-to-tail-conjugated primary enamines containing heterene substituents, **222, 193**
- Enantioselective synthesis**
- enantioselective synthesis of ferrocene analogs of hexestrol and estradiol; recognition towards estradiol receptors, **222, 51**
- Enantioselectivity**
- the importance of phosphine-to-rhodium ratios in enantioselective hydroborations, **220, 93**
- dirhodium(II) tetrakis[*N,N*-dimethyl-2-pyrrolidone-5(*S*)-carboxamide]. Structural effects on enantioselection in metal carbene transformations, **220, 193**
- optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220, 283**
- Energy gap law**
- relationship between four-state delineation and two-state delineation in electron transfer reactions and excitation transfer, **225, 185**
- Energy transfer reactions**
- light induced exchange energy transfer reactions of micelle forming Ru(II) diimine complexes, **225, 275**
- Enolate complexes**
- the preparation of chiral titanium reagents: a comparison of new and known ligands, **222, 95**
- Enthalpy**
- direct solution calorimetric measurements of enthalpies of proton and electron transfer reactions for transition metal complexes. Thermochemical study of metal-hydride and metal-metal bond energies, **227, 285**
- Enzyme complexes**
- methane as the product of reaction of methyl-coenzyme-M with monovalent nickel complexes in aqueous solutions. A model for the *in vivo* activity of cofactor F<sub>430</sub>, **227, 1**
- Epoxidation**
- optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220, 283**
- EPR spectra**
- synthesis, spectra and electrochemical behaviour of biomimetic copper(II) complexes with CuN<sub>5</sub> and CuN<sub>6</sub> chromophores, **224, 19**
- EPR spectroscopy**



- the anomalous EPR spectra of the iron(III) dichalcogenocarbamates, **217**, 101
- Equilibrium studies
- equilibrium and spectroscopic studies of diethyltin(IV) complexes formed with carbohydrate derivatives of thiazolidine-4-carboxylic acid, **218**, 65
  - equilibrium studies of 1,2-diaminobenzene and 3,4-diaminobenzoate anion with dinuclear cobalt(II) 1,4,7,13,16,19-hexaaza-10,22-dioxacyclotetracosane, **227**, 49
  - the equilibrium between acetyl- and methylcobalt tetracarbonyl, **227**, 211
- ESR study
- ESR studies on distortion in the macrocyclic ring of copper(II) chlorophylls, **224**, 181
- Esterification
- selective transformations of alkynols catalyzed by ruthenium complexes, **222**, 155
- Estradiol complexes
- the synthesis and reactions of A-ring aromatic steroid complexes of manganese tricarbonyl, **220**, 297
- Estrogen complexes
- estrogen derivatives of transition metal complexes for analytical detection enhancement. Part II, **218**, 207
- Estrone complexes
- the synthesis and reactions of A-ring aromatic steroid complexes of manganese tricarbonyl, **220**, 297
- Ether-phosphine complexes
- synthesis and fluxional behaviour of (cyclopentadienyl) and (pentamethylcyclopentadienyl)ruthenium(II) complexes with hemilabile ether-phosphines. <sup>31</sup>P DNMR spectroscopic studies and line shape analysis, **215**, 165
- Europium complexes
- an europium(III) complex as an efficient catalyst for the Michael reaction of  $\alpha,\beta$ -enones with ketene silyl acetals: LIS-NMR analysis for the transition state through complexation between europium(III) catalyst and enones, **222**, 377
  - luminescent europium(III) and terbium(III) ions encapsulated in a 2-aminoethoxy or carbamoyloxy substituted calixarene host, **227**, 293
- EXAFS
- EXAFS investigation on nickel substituted Cu<sub>2</sub>Zn bovine superoxide dismutase, **216**, 121
  - EXAFS studies of a chiral rhenium lactone complex and its precursors including multiple scattering calculations, **222**, 27
- Excitation transfer
- relationship between four-state delineation and two-state delineation in electron transfer reactions and excitation transfer, **225**, 185
- Exo* and *endo* isomers
- crystal and molecular structures of [*exo*- and *endo*-(RO)<sub>2</sub>P(O)- $\eta^5$ -C<sub>6</sub>H<sub>6</sub>]Mn(CO)<sub>3</sub>, **224**, 147
- Fast atom bombardment mass spectrometry
- liquid matrix induced processes in fast atom bombardment mass spectrometry of polyazacrown transition metal complexes, **219**, 85
- Fenske–Hall MO calculation
- synthesis and experimental/theoretical investigation of the high-nuclearity cubic T<sub>d</sub> [Au<sub>6</sub>Ni<sub>12</sub>(CO)<sub>24</sub>]<sup>2-</sup> cluster, an initial example of a discrete gold–nickel bimetallic-bonded species: comparative analysis of the results of electron-counting methods and the Fenske–Hall MO model in rationalizing the bonding interactions of its Au<sub>6</sub>Ni<sub>12</sub> core consisting of five face-fused metal octahedra, **227**, 269
- Ferrate complexes
- kinetics and mechanism of the ferrate oxidation of hydrazine and monomethylhydrazine, **225**, 145
- Ferrocene
- enantioselective synthesis of ferrocene analogs of hexestrol and estradiol; recognition towards estradiol receptors, **222**, 51
- Ferrocene/ferrocenium couple
- dependence of electron transfer kinetics of the ferrocene/ferrocenium couple on the viscosity in ambient temperature chloroaluminate ionic liquids, **225**, 9
- Ferrocene complexes
- preparation, structure and catecholase-mimetic activity of two mononuclear ferrocenecarboxylate copper(II) complexes, **215**, 131
  - a new functionalised oligopyridine ligand containing ferrocene as a ball-bearing spacer for metallosupramolecular chemistry, **224**, 11
  - electrochemical recognition of Group 1 and 2 metal cations by redox-active ionophores, **225**, 137
  - controlling the acidity of the carboxylic group by a ferrocene based redox switch, **225**, 239
- Ferrocene substituted complexes
- the kinetics of electron-transfer reactions of the [FeCp(CpCH<sub>2</sub>N(CH<sub>3</sub>)<sub>3</sub>)<sup>+2+</sup> couple in the presence of cyclodextrins in aqueous media, **225**, 51
- Ferrocenylcarboxylic acid complexes
- controlling the acidity of the carboxylic group by a ferrocene based redox switch, **225**, 239
- Ferrocenyl complexes
- electron transfer in *trans*-[Pt(PPh<sub>3</sub>)<sub>2</sub>(-C≡C-Fc)<sub>2</sub>] and related compounds, **225**, 35
- Ferrocenyl phosphine complexes
- synthesis and structure of new chiral ferrocenylphosphines for asymmetric catalysis, **222**, 213
- Fischer carbene complexes
- stereoselective aldol addition reactions of Fischer carbene complexes via electronic tuning of the metal center for enolate reactivity, **220**, 215
  - a study of the reactivity and structure of cyclic  $\alpha,\beta$ -unsaturated Fischer-type carbene complexes, **220**, 233
- Fluoride complexes
- uranium hexafluoride: laboratory scale synthesis, coordination behaviour towards nitriles and Raman spectra, **215**, 5
- Formamide complexes
- synthesis, crystal structure and thermal stability of the MgCl<sub>2</sub> formamide adduct, **216**, 261
- Formamidinato complexes
- a new class of dinuclear compounds: the synthesis and X-ray structural characterization of tris( $\mu$ -diphenylformamidinato)diiron, **219**, 7
  - divalent iron formamidinato complexes: a highly distorted dinuclear compound, **224**, 5
- Formamido complexes
- molecular and crystal structures of *cis*-Pt(DMSO)(C<sub>2</sub>H<sub>5</sub>CN)Cl<sub>2</sub> and Cl(DMSO)Pt( $\mu$ -N(CH<sub>3</sub>)CHO)<sub>2</sub>Pt(DMSO)Cl, **216**, 163
- Formation constants
- equilibrium and spectroscopic studies of diethyltin(IV) complexes formed with carbohydrate derivatives of thiazolidine-4-carboxylic acid, **218**, 65
- Four-state delineation
- relationship between four-state delineation and two-state delineation in electron transfer reactions and excitation transfer, **225**, 185
- Free radical transfer
- free radical-mediated electron transfer in organometallic complexes: homolysis and alkyl group crossover reactions for alkyliron(II) porphyrins, **226**, 231

- Friedel–Crafts reaction  
electrophilic addition to styrylsilanes: sequential carbon–carbon bond forming reactions, **220**, 145
- Functionalized-phosphine complexes  
functionalised phosphines: structure of bis(2-aminophenyl)diphenylphosphine- $\kappa^2N,P$ copper(I) perchlorate ethanol solvate, **217**, 191
- Furanone  
rhodium-catalyzed synthesis of 2(5*H*)-furanones from terminal alkynes and non-substituted alkynes under water-gas shift reaction conditions, **220**, 45
- Gadolinium complexes  
synthesis and crystal structure of [GdCl<sub>2</sub>(dibenzo-18-crown-6)-(MeCN)][SbCl<sub>6</sub>] $\cdot$ 2MeCN, dibenzo-18-crown-6 = 2,3,11,12-dibenzo-1,4,7,10,13,16-hexaoxacyclooctadeca-2,11-diene, **215**, 209  
stability constants of the lanthanide(III)-1,4,7,10-tetraazacyclododecane-*N,N',N'',N'''*-tetraacetate complexes, **221**, 165
- Galactose oxidase complexes  
towards a model for galactose oxidase: the crystal structure of {2-[bis(2-pyridylethyl)aminomethyl]phenolato} copper(II) perchlorate, **215**, 1
- Gallium complexes  
trimethylgallium based phosphinogallanes. Synthesis and molecular structure of [Me<sub>2</sub>Ga–P(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, **216**, 267  
spectroscopic and structural studies on volatile gallium  $\beta$ -diketonates as potential precursors for MOCVD, **217**, 71
- Gamma-irradiation  
extrusion of molecular clusters from solid-state materials: synthesis by application of  $\gamma$ -irradiation. Molecular and crystal structure of (H<sub>2</sub>O<sub>4</sub>)(Et<sub>4</sub>N)[Mo<sub>3</sub>S<sub>7</sub>Br<sub>6</sub>], **223**, 5
- Gold complexes  
methyl dithiocarbamate gold(I) and gold(III) complexes. Synthesis and reactivity with amines, **215**, 215  
syntheses of dinuclear gold(I) ring complexes containing 1,1-dicyanoethene-2,2-dithiolate-*S,S* and bis(diphosphines) as bridging ligands. X-ray crystal structure of [Au<sub>2</sub> $\mu$ -(i-MNT) $\mu$ -(dppe)]<sub>2</sub>, **217**, 45  
synthesis of water-soluble (tri(hydroxymethyl)phosphine)gold(I) complexes containing a nucleoside ligand, **217**, 201  
gold in organic synthesis Part 2. Preparation of benzyl-alkyl and -arylketonates via C–C coupling, **220**, 1  
reactions of [Au(C<sub>6</sub>F<sub>5</sub>)(SC<sub>4</sub>H<sub>8</sub>)] with diazoalkanes. Synthesis and molecular structures of [Au(C<sub>6</sub>F<sub>5</sub>)(Ph<sub>2</sub>C=N–N=CPh<sub>2</sub>)]<sub>2</sub>, **222**, 267  
di- and tetranuclear gold(II) complexes with dithiocarbamate and related ligands. X-ray structure of [Au<sub>2</sub>( $\mu$ -CH<sub>2</sub>PPh<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>(S<sub>2</sub>CNMe<sub>2</sub>)<sub>2</sub>]<sub>2</sub>, **223**, 55  
solid state and solution studies of a T-shaped gold(I) complex: Au[(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>PCH<sub>2</sub>N(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>]<sub>2</sub>Cl, **227**, 85
- Gold derivatives  
bimolecular electron transfer at high driving forces. Photoreactions of gold(I) and iridium(I) complexes with pyridinium derivatives, **225**, 319
- Gold–nickel cluster  
synthesis and experimental/theoretical investigation of the high-nuclearity cubic T<sub>d</sub> [Au<sub>6</sub>Ni<sub>12</sub>(CO)<sub>24</sub>]<sup>2-</sup> cluster, an initial example of a discrete gold–nickel bimetallic-bonded species: comparative analysis of the results of electron-counting methods and the Fenske–Hall MO model in rationalizing the bonding interactions of its Au<sub>6</sub>Ni<sub>12</sub> core consisting of five face-fused metal octahedra, **227**, 269
- Grignard reagent  
magnesium bromide in Grignard reagent formation, **222**, 365
- Hafnium complexes  
alkyl for hydride exchange between alane-trimethylamine and Group IVB metal alkyls, **218**, 47  
synthesis, characterization and X-ray crystal structure of the benzenedithiolatohafnium(IV) porphyrin complex, **221**, 51
- Halide complexes  
phosphine and phosphine oxide adducts of vanadium(IV) chloride: synthesis and structural studies, **215**, 47  
the unusual crystallographic disorder of M<sub>2</sub> units in M<sub>2</sub>Cl<sub>4</sub>(PEt<sub>3</sub>)<sub>4</sub> (M = Re, Mo), **215**, 103  
cobalt chloride and phase-transfer catalyzed carbonylation of benzyl chlorides, **216**, 97  
molecular mechanical and quantum chemical study on the species involved in the hydrolysis of *cis*-diamminedichloroplatinum(II) and substituted bis(ethylenediamine)dichloroplatinum(II) complexes Part I. Reactants and products, **217**, 159  
electrochemistry of organometallic halide complexes III. Oxidation of [C<sub>6</sub>Me<sub>6</sub>RuCl<sub>2</sub>]<sub>2</sub>, **218**, 33  
ruthenium complex catalyzed polymerization of OH or COOH group containing alkynes to give functionalized poly(acetylene)s, **220**, 35  
uranium compounds as catalysts for aldolisation reactions, **222**, 91  
indium(III) halide-3,5-di-*tert*-butyl-*o*-benzosemiquinone systems, **225**, 207  
NQR studies of polyhalocuprate(I) anions, **227**, 153
- Helicates  
photophysical properties of mono-, di- and tetranuclear copper(I)–polypyridine complexes, **225**, 251
- Hemocyanin model compounds  
hemocyanin model compounds; a quantum mechanical calculation of geometry-dependent exchange interaction, **221**, 141
- Hemoglobins  
a spectroelectrochemical method for evaluating factors which regulate the redox potential of hemoglobins, **226**, 187
- Hemoprotein enzyme  
mesoferriheme regeneration from its hemoprotein enzyme compound I analog: absorption and kinetic properties of one- and two-electron oxidation products, **225**, 255
- Heteroatom  
synthesis, properties and reactions of sila-functional  $\alpha$ -heteroatom-substituted organosilicon compounds, **220**, 129
- Heterobimetallic complexes  
syntheses of ruthenium–iridium heterobimetallic complexes using tridentate phosphine ligands, **227**, 79
- Heterobinuclear complexes  
hydroformylation of alkenes and alkynes using a heterobinuclear Rh–W catalyst, **220**, 187
- Heterocyclic ligand complexes  
intramolecular electron transfer from Ru<sup>II</sup>(EDTA)<sup>2-</sup> to Co<sup>III</sup>(NH<sub>3</sub>)<sub>5</sub><sup>3+</sup> via nitrogen heterocycle bridging ligands, **225**, 129
- Hexaazacyclophane complexes  
reduction of hexaazacyclophane complexes of Co(II) and Cu(II) by  $\cdot$ CH<sub>2</sub>OH and e<sub>solv</sub><sup>-</sup>; a pulse radiolysis study, **226**, 151
- Hexadentate ligand complexes  
synthesis and characterization of hexadentate cobalt(III) complexes with new edta-type ligands Part 4. Circular dichroism of the *trans*-(O<sub>5</sub>O<sub>6</sub>) isomer of a cobalt(III) complex of 1,3-propanediamine-*N,N,N'*-triacetic-*N'*-3-propionic acid, **219**, 147
- Hexamethylbenzene complexes  
electrochemistry of organometallic halide complexes III. Oxidation of [C<sub>6</sub>Me<sub>6</sub>RuCl<sub>2</sub>]<sub>2</sub>, **218**, 33
- Hexaruthenium clusters

- the formation of carbides in hexaruthenium carbonyl clusters, **227**, 197
- Hippuric acid complexes  
structural, spectroscopic and potentiometric investigation on the coordination properties of 4-NO<sub>2</sub>-hippuric acid towards Cu(II) in binary and ternary systems, **227**, 105
- Host  
A 'calix[4]arene' porphyrin as a new host and an oxygen carrier model, **224**, 1
- Hückel molecular orbital  
synthesis, molecular structure and electrochemical investigation of the 1,2,5-thiadiazole-3,4-dithiolate-substituted complex Cp\*Ru(NO)(tdas). Evidence for nitrosyl bending during reduction, **226**, 267
- Hydration  
hydration of conjugated dienes to produce ketones catalyzed by ruthenium complexes, **222**, 225
- Hydrazine  
kinetics and mechanism of the ferrate oxidation of hydrazine and monomethylhydrazine, **225**, 145
- Hydrazone complexes  
metal complexes of 5-acetyl-2,2'-pyridylthiophene aroylhydrazones. Crystal structure of a cobalt(II) derivative, **223**, 97
- Hydride complexes  
alkyl for hydride exchange between alane-trimethylamine and Group IVB metal alkyls, **218**, 47  
hydrozirconation/transmetalation of acetylenic stannanes. New 1,1-dimetallio reagents, **220**, 41  
reaction of primary silanes with dinuclear rhodium hydride complexes: silane coupling reactions, **222**, 345
- Hydrido complexes  
cyclometallated derivatives of platinum(II) derived from 1,4-benzodiazepin-2-ones. Crystal and molecular structure of Pt(L)(HL)Cl (HL = 7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one, DIAZEPAM), a molecule containing a neutral and a deprotonated 1,4-benzodiazepin-2-one, **215**, 17
- Hydroboration  
the importance of phosphine-to-rhodium ratios in enantioselective hydroborations, **220**, 93
- Hydrodehalogenation  
synthesis of symmetric and unsymmetric 1,4-bis(*p*-R-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289
- Hydroformylation  
a mechanistic study of the homogeneous catalytic hydroformylation of formaldehyde: synthesis and characterization of model intermediates, **218**, 89  
hydroformylation of alkenes and alkynes using a heterobinuclear Rh–W catalyst, **220**, 187
- Hydrogenation  
the role of H<sub>2</sub>Ru(CO)<sub>2</sub>(PBu<sub>3</sub>)<sub>2</sub> in the activation of alkynes and alkenes, **227**, 247
- Hydrogen isocyanide complexes  
electron-transfer activation of the aminocarbene and the hydrogen isocyanide complexes *trans*-[ReCl(CNH<sub>n</sub>)<sub>n</sub>-(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>][BF<sub>4</sub>]<sub>n-1</sub> (*n* = 2 or 1). Interconversion of coordinated CNH<sub>2</sub> and CNH, **226**, 9
- Hydrolysis  
synthesis and hydrolysis of an ester of the 'one ring open' hydrolysis product of the anticancer drug 1,2-bis(3,5-dioxopiperazin-1-yl)propane, 'Razoxane', **215**, 55  
chiral metal complexes 40\*. The sterically mediated hydrolysis of a coordinated imine, **221**, 191  
the synthesis, X-ray structure and hydrolysis kinetics of chloro(2-aminomethylpyridine)(triamine)chromium(III) salts, **227**, 11
- Hydrolytic activity  
the synthesis, structure and hydrolytic activity of a functional pentacoordinate zinc complex, **219**, 205
- Hydrosilylation  
catalytic hydrosilylation of carbon monoxide with cobalt carbonyls, **222**, 131
- Hydrosulfination  
palladium complex catalysed synthesis of sulfinic acids, sulfinic acid esters, sulfonic acids and *S*-alkyl alkanethiosulfonates, **222**, 381
- Hydroxo complexes  
synthesis, characterization and crystal structure of (C<sub>3</sub>H<sub>3</sub>N<sub>2</sub>)<sub>2</sub>[Sn<sub>2</sub>(NO<sub>3</sub>)<sub>4</sub>(μ-OH)<sub>2</sub>Me<sub>4</sub>], a di-μ-hydroxo-bridged compound with heptacoordinated tin(IV), **216**, 257
- Hydroxylation  
oxidation of *cis*- and *trans*-pinane with RuO<sub>4</sub> generated in situ, **222**, 115
- Hydroxypyridinones  
stabilities of 1,2-diethyl-3-hydroxy-4-pyridinone chelates of divalent and trivalent metal ions, **223**, 21
- Hydrozirconation  
hydrozirconation/transmetalation of acetylenic stannanes. New 1,1-dimetallio reagents, **220**, 41
- Imidazole complexes  
mixed-ligand copper(II) complexes with iminodiacetato and 2- or 4-methylimidazole: molecular and crystal structure of the 'remote' isomer iminodiacetato(5-methylimidazole)-copper(II) monohydrate, [Cu(IDA)(5MeImH)]·H<sub>2</sub>O, **215**, 73  
synthesis, structural characterization and dioxygen reactivity of imidazole-ligated Cu(I) complexes, **218**, 103  
steric and electronic tuning of Michael addition reactions induced by pentacarbonyl metal fragments, **222**, 207
- Imidazoles  
a novel reaction of the cyano ligand opening an organometallic route to 4-amino imidazoles: four component condensation (4CC) with isocyanide, aldehyde and amine, **222**, 275
- 2-Imidazolines  
reactions of aziridine with platinum(II) nitriles. Formation of (aziridino)amidines and 2-imidazolines and X-ray structure of *trans*-[PtCl<sub>2</sub>{N(H)=C(Ph)NCH<sub>2</sub>CH<sub>2</sub>}]<sub>2</sub>, **222**, 327
- Imido complexes  
reactions of the technetium(V) phenylimido complex [TcX<sub>5</sub>(NPh)(PPh<sub>3</sub>)<sub>2</sub>] with aromatic thiolate ligands (where X = Cl, Br), **218**, 97  
phenylimido derivatives of [Mo<sub>6</sub>O<sub>19</sub>]<sup>2-</sup>: syntheses, X-ray structures, vibrational, electrochemical, <sup>95</sup>Mo and <sup>14</sup>N NMR studies, **224**, 81
- Imine complexes  
chiral metal complexes 40\*. The sterically mediated hydrolysis of a coordinated imine, **221**, 191  
a novel sodium template approach for preparing tetraimine macrocycles of 2,6-diformyl-4-methylphenol and diamino derivatives, **223**, 169
- Imine hydrogenation  
a comparison of catalytic activity for imine hydrogenation using Ru ditertiary phosphine complexes, including chiral systems, **222**, 85
- Inclusion complexes  
the kinetics of electron-transfer reactions of the [FeCp(CpCH<sub>2</sub>N(CH<sub>3</sub>)<sub>3</sub>)<sup>+2+</sup> couple in the presence of cyclodextrins in aqueous media, **225**, 51
- Indenyl complexes  
ligand substitution at 19-electron organometallic centers. Electrocatalytic CO substitution reactions of (methylcyclopenta-

- diényl)Mn(CO)<sub>2</sub>NO<sup>+</sup> and (indenyl)Mn(CO)<sub>2</sub>NO, **226**, 53
- Indium complexes**
- oxo- and sulfido-bridged sandwich cubane-type molybdenum–indium mixed-metal cluster. Synthesis and X-ray structure of [(H<sub>2</sub>O)<sub>9</sub>Mo<sub>3</sub>S<sub>3</sub>OInOS<sub>3</sub>Mo<sub>3</sub>(H<sub>2</sub>O)<sub>9</sub>](CH<sub>3</sub>·C<sub>6</sub>H<sub>4</sub>·SO<sub>3</sub>)<sub>8</sub>·30H<sub>2</sub>O, **216**, 13
- indium(III) halide–3,5-di-tert-butyl-*o*-benzosemiquinone systems, **225**, 207
- Infrared Raman spectroscopy**
- the vibrations of bridging carbonyl groups as more sensitive indicators of molecular geometry than terminal carbonyl groups in transition metal carbonyl cluster compounds, **227**, 241
- Infrared spectroscopy**
- Raman and infrared spectroscopic studies of tris triphenyl phosphine chloride complexes of transition metals, **216**, 191
- spectroscopic investigation of mixed-ligand nickel(II) chelates containing β-ketoenols and 1,2-diamines, **221**, 17
- Insertion**
- synthesis of η<sup>3</sup>-allyl–Pd complexes and subsequent formation of N-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- Insertion reactions**
- insertion reaction of dimethylacetylenedicarboxylate with the head-to-tail isomer of Pt<sub>2</sub>I<sub>2</sub>(μ-Ppy)<sub>2</sub>; Ppy<sub>3</sub> = tris(2-pyridyl)phosphine, **217**, 209
- nitrosoarene and nitrosoalkane insertion reactions of titanacyclobutenes, **222**, 305
- alkylation of 2-vinylpyridine and crystal structure of bis{(acetylacetonate)(1-pyridylbutyl)nickel(II)} complex, **227**, 149
- Intermolecular coupling elements**
- basis set effects on the direct calculation of intermolecular coupling elements for electron transfer reactions, **226**, 237
- Interstitial clusters**
- electron transfer reactions in non-stoichiometric ceria and urania, **225**, 305
- Intervalence spectra**
- electron transfer reactions in non-stoichiometric ceria and urania, **225**, 305
- Intervalence transfer**
- charge type effects on the temperature dependence of intervalence transfer in ferrocyanide substituted ruthenium(III) amine and aminopolycarboxylate compounds, **224**, 189
- Intramolecular electron transfer**
- intramolecular electron transfer from Ru<sup>II</sup>(EDTA)<sup>2-</sup> to Co<sup>III</sup>(NH<sub>3</sub>)<sub>5</sub><sup>3+</sup> via nitrogen heterocycle bridging ligands, **225**, 129
- specific ion-pairing effects in weakly exoergic intramolecular electron transfer, **225**, 269
- intramolecular electron transfer between Ru(I) and Ru(III) and the heme iron of cytochrome *c* labeled with ruthenium(II) polypyridine complexes, **226**, 129
- Iodide complexes**
- a novel tetranuclear iodoargentate(I) ion, [Ag<sub>4</sub>I<sub>8</sub>]<sup>4-</sup>, with a cubane-type core, **217**, 15
- synthesis and characterization of uranium triiodide tetrakis(acetonitrile), **218**, 203
- Iodo complexes**
- photoredox chemistry of iodoargentate(II) complexes in acetonitrile, **221**, 79
- Ion-pairing**
- outer-sphere coordination of polycyanometallate anions with polyammonium macrocycles: a spectrophotometric study, **227**, 71
- Ion-pairing effect**
- specific ion-pairing effects in weakly exoergic intramolecular electron transfer, **225**, 269
- Ion pairs**
- spectroscopy and photoredox chemistry of ion-paired cobalt(III) am(m)ine complexes, **226**, 285
- Iridium complexes**
- polyfunctional phosphine ligands II. Iridium(I) complexes of the 7-diphenylphosphino-2,4-dimethyl-1,8-naphthyridine (dpnpy) ligand. X-ray crystal structure of {[Ir(cod)Cl]<sub>2</sub>(μ-dpnpy)}, **216**, 209
- addition of carbanions to tris(trimethylphosphine)-(cyclooctadiene)iridium(I) chloride: three different modes of reaction depending on the nature of the carbanion, **222**, 135
- syntheses of ruthenium–iridium heterobimetallic complexes using tridentate phosphine ligands, **227**, 79
- Iridium derivatives**
- bimolecular electron transfer at high driving forces. Photoreactions of gold(I) and iridium(I) complexes with pyridinium derivatives, **225**, 319
- Iron complexes**
- synthesis, crystal and molecular structures, UV–Vis spectroscopy and electrochemical properties of two iron(III) phenolate complexes, **215**, 123
- NMR parameters as steric probes for arylcyclopentadienyl iron complexes, **215**, 139
- solvent effects on acetylacetonato iron complexes, **215**, 185
- two silicon-bridged cyclopentadienyl Fe<sub>2</sub>S<sub>4</sub> cluster units connected through an Fe(III)S<sub>6</sub> unit. Crystal and molecular structure of [(CH<sub>3</sub>)<sub>2</sub>SiCp<sub>2</sub>]<sub>2</sub>Fe<sub>2</sub>(μ<sub>3</sub>-S<sub>2</sub>)(μ<sub>4</sub>-S<sub>2</sub>)(μ<sub>4</sub>-S<sub>4</sub>)]FeCl<sub>4</sub>, **216**, 1
- binuclear oxo-bridged iron(III) complexes of *N,N'*-bis(2-pyridylmethyl)ethane-1,2-diamine (bispicen) and related ligands, **216**, 21
- bulky substituent effects on the iron(III) complexation of 1,4,7-triazaacyclononane, **216**, 89
- 'insulation' to lattice effects on quadrupole splitting in the Mössbauer spectra of salts of [Fe<sub>4</sub>S<sub>4</sub>(Cys)<sub>4</sub>]<sup>2-</sup>, a ferredoxin model, **217**, 3
- electrochemical investigation of the electron-poor/precise (*n* = 5/6) complexes [Fe<sub>2</sub>(CO)<sub>*n*</sub>(μ-PR<sub>2</sub>)(μ-PR'<sub>2</sub>)] (*n* = 5, R = R' = Bu<sup>1</sup>; *n* = 6, R = R' = Ph; R = Bu<sup>1</sup>, R' = Ph; R = Bu<sup>1</sup>, R' = Cy). EPR study of the radical anion [Fe<sub>2</sub>(μ-PBu<sup>1</sup>)<sub>2</sub>(CO)<sub>3</sub>], **217**, 29
- the anomalous EPR spectra of the iron(III) dichalcogenocarbamates, **217**, 101
- stereoselectivity in reactions of metal complexes Part XVIII. Kinetics and stereoselectivity in electron-transfer reactions between [Co(L)H<sub>2</sub>O]<sup>+</sup> (L = *N,N'*-[(pyridine-2,6-diyl)bis(methylene)]bis[proline] (promp) and *N,N'*-[(4-methoxy-pyridine-2,6-diyl)bis(methylene)]bis[proline] (MeO-promp) and optically active iron(II) complexes, **218**, 73
- a new class of dinuclear compounds: the synthesis and X-ray structural characterization of tris(μ-diphenylformamidinato)diiron, **219**, 7
- transition metal complexes with sulfur ligands Part CII. Changes in configuration and reactivity caused by alkyl substituents R in [Fe(L)(<sup>1</sup>N<sub>R</sub>S<sub>4</sub>')] complexes (L = CO, NO<sup>+</sup>, NO, PMe<sub>3</sub>, N<sub>2</sub>H<sub>4</sub>; R = CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub><sup>-</sup>), **219**, 75
- preparation, properties and coordination behaviour of planar or tridimensional compartmental Schiff bases, **219**, 135
- equilibrium and kinetic study of piperidine binding to phthalocyaninatoiron(II) in dimethyl sulfoxide, **219**, 199
- <sup>1</sup>H NMR analysis of some η<sup>6</sup>-(substituted aryl phenylsulfonyl-acetonitrile)-η<sup>5</sup>-cyclopentadienyliron cations, **219**, 221

- preparation, properties and reactions of metal-containing heterocycles Part LXXXIX. Metalla[n]- and metalla[m,n]-orthocyclophanes as mediators for the synthesis of cyclic ketones, **220**, 107
- a study of the reactivity of (methyl 2-acetamidoacrylate)-tricarbonyliron(0) leading to a novel synthesis of  $\beta,\beta,\beta$ -tri-alkyl  $\alpha$ -amino acids, **220**, 137
- reactivity of allene at phosphine-bridged di-iron centres: X-ray crystal structures of  $[\text{Fe}_2(\text{CO})_5\{\mu\text{-}\sigma,\eta^3\text{-C(O)C(CH}_2)_2\}(\mu\text{-dppm})]$  and  $[\text{Fe}_2(\text{CO})_4\{\mu\text{-}\eta^3,\eta^3\text{-(CH}_2)_2\text{C}_2(\text{CH}_2)_2\}(\mu\text{-dppm})]\cdot\text{Et}_2\text{O}$ , **220**, 201
- preparation of  $\eta^1$ -1,3-dienyl,  $\eta^1$ -1,2-dienyl and  $\eta^3$ -cyclobutene complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305
- activation of molecular oxygen by iron nitrosyls in the presence of the bidentate phosphines 1,2-bis(diphenylphosphino)ethane and ethene, **221**, 133
- conversion of cationic phosphite complexes into phosphonate complexes with selective deamination on the coordinating phosphorus. Reaction of  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{L}_2\text{Fe}\{\text{P}(\text{NC}_4\text{H}_8)_n\text{-}(\text{OMe})_{3-n}\}]^+$  (L = CO, PMe<sub>3</sub>) with KOH to afford  $(\eta^5\text{-C}_5\text{H}_5)_2\text{L}_2\text{Fe}\{\text{P}(\text{O})(\text{NC}_4\text{H}_8)_{n-1}(\text{OMe})_{3-n}\}$ , **222**, 123
- Mössbauer spectroscopy of iron dextran: comparison of solid and frozen solution, **223**, 9
- synthesis and reactions of the novel sulfonium-methylidene complex  $[\text{Fe}_2(\text{CO})_2(\text{cp})_2(\mu\text{-CO})\{\mu\text{-C(H)(SMe}_2)\}]\text{SO}_3\text{CF}_3$ , **223**, 31
- a one-pot synthesis of cyclopropanes from aromatic aldehydes and styrene via iron carbene complexes, **223**, 159
- divalent iron formamidinato complexes: a highly distorted dinuclear compound, **224**, 5
- transition metal complexes with sulfur ligands CIV. Protonation and alkylation of thiolate donors of  $[\text{Fe}(\text{CO})_2(\text{S}_4^{\prime})]$  and  $[\text{Fe}(\text{CO})(\text{S}_5^{\prime})]$ : syntheses and properties of Fe(II) complexes with open-chain thioether ligands ( $\text{S}_4^{\prime 2-} = 1,2$ -bis(2-mercaptophenylthio)ethane(2-);  $\text{S}_5^{\prime 2-} = 2,2'$ -bis(2-mercaptophenylthio)diethylsulfide(2-)), **224**, 45
- transition metal complexes with sulfur ligands CIII.  $[(\mu\text{-S}_2)\{\text{Fe}(\mu\text{-S}_4^{\prime})\}_2]$ , a novel  $[2\text{Fe}\text{-}2\text{S}]$  cluster with octahedral low-spin  $[\text{Fe}^{\text{II}}\text{S}_6]$  centers ( $\text{S}_4^{\prime 2-} = 1,2$ -bis(2-mercaptophenylthio)ethane(2-)), **224**, 35
- transition metal complexes with sulfur ligands CVI. Phosphine and azido  $[\text{Fe}(\text{L})(\text{N}_3\text{S}_4^{\prime})]$  complexes; structural characterization of the key complex  $[\text{Fe}(\text{N}_3\text{S}_4^{\prime})]\cdot\text{THF}$  (L = PMe<sub>3</sub>, PBu<sub>3</sub>, PMe<sub>2</sub>Ph, PMePh<sub>2</sub>, N<sub>3</sub><sup>-</sup>;  $\text{N}_3\text{S}_4^{\prime 2-} =$  dianion of 2,2'-bis(2-mercaptophenylthio)diethylamine), **224**, 61
- rotation and orientation of axially coordinated imidazoles in low-spin ferric porphyrin complexes, **224**, 113
- transition-metal silyl complexes 51. Preparation of the mixed silyl/stannyl complexes  $(\pi\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})(\text{SnPh}_3)(\text{SiR}_3)\text{H}$ , **224**, 177
- electrochemistry of nitrite reductase model compounds 5. Electrochemistry and spectroelectrochemistry of iron porphyrinone, porphindione and isobacteriochlorin complexes, **225**, 57
- the use of pressure-tuning spectroscopy to distinguish between one- and two-electron transfer processes, **225**, 89
- stereoselective electron transfer between Cu(II) superoxide dismutase and  $\Lambda$ - and  $\Delta$ - $[\text{Fe}(\text{pdta})]$ , **225**, 151
- syntheses and structures of mixed-ligand tetrairon-sulfur clusters,  $[\text{Cp}'_2(\text{Ph}_2\text{C}_2\text{S}_2)_2\text{Fe}_4\text{S}_4](\text{PF}_6)$  and  $[\text{Cp}'_3(\text{Ph}_2\text{C}_2\text{S}_2)\text{Fe}_4\text{S}_3](\text{PF}_6)_n$  ( $n = 1$  and  $2$ ) ( $\text{Cp}' = \eta^5\text{-C}_5\text{Me}_5$ ). Structural change of the clusters accompanying electron transfer reactions, **225**, 229
- electrocatalytic reduction of CO<sub>2</sub> and O<sub>2</sub> with electropolymerized films of vinyl-terpyridine complexes of Fe, Ni and Co, **225**, 285
- the redox behaviour of the cluster anion  $[\text{Fe}_4\text{N}(\text{CO})_{12}]^-$ . Electron transfer chain catalytic substitution reactions. Crystal structure of  $[\text{Fe}_4\text{N}(\text{CO})_{11}\text{PPh}_3]$ , **226**, 1
- valence tautomerism for catechol/semiquinone complexes of the *trans*-M(Bupy)<sub>2</sub>(3,6-DBQ)<sub>2</sub> (M = Mn, Fe, Co) series, **226**, 91
- photochemical and spectroscopic studies of complexes of iron(III) with citric acid and other carboxylic acids, **226**, 117
- reaction of low-valent iron porphyrins with alkyl containing supporting electrolytes, **226**, 195
- electron transfer quenching of the excited MLCT state of Ru(II) in the double complex salts  $[\text{Ru}(\text{bpy})_3]_2[\text{M}(\text{CN})_6]\text{Cl}\cdot 8\text{H}_2\text{O}$  (M = Co and Fe), **226**, 203
- free radical-mediated electron transfer in organometallic complexes: homolysis and alkyl group crossover reactions for alkyliron(II) porphyrins, **226**, 231
- the construction of molecular systems capable of photoinduced multielectron charge transfer: a photochemical analysis of the reactivity of  $[(\text{CN})_5\text{Fe}^{\text{II}}(\text{CN})\text{-Pt}^{\text{IV}}(\text{L})_4(\text{NC})\text{-Fe}^{\text{II}}(\text{CN})_5]^4-$  (L = NH<sub>3</sub>, ethylenediamine), **226**, 251
- heteronuclear metal cluster complexes with a new coordination mode of the Fe<sub>2</sub>S<sub>2</sub>(CO)<sub>6</sub> unit. Synthesis, structure and possible reaction pathway of the first Fe-Cu-S cluster compound,  $[\text{Fe}_6\text{Cu}_5\text{S}_6(\text{CO})_{18}(\text{PPh}_3)_2]$ , **227**, 119
- metal cluster topology 16. Macropolygons and macropolyhedra in coinage metal iron carbonyl cluster anions and related compounds, **227**, 207
- Iron-sulfur complexes
- redox behavior of the iron-sulfur cluster  $[\text{Fe}_4\text{Cp}_3\text{S}_5][\text{PF}_6]_2$  in protic organic solvents and aqueous micellar solutions, **226**, 341
- Irradiation
- synthesis, structure and solid state polymerization of diaqua-bis(propynoato)magnesium, **222**, 339
- Isobacteriochlorin complexes
- electrochemistry of nitrite reductase model compounds 5. Electrochemistry and spectroelectrochemistry of iron porphyrinone, porphindione and isobacteriochlorin complexes, **225**, 57
- Isoocyanide complexes
- pressure and concentration dependent formation of oligomers of tetrakis(*p*-methylphenylisocyanide)rhodium(I), **216**, 245
- mercury and platinum mixed cluster having a prismatic structure,  $[\text{HgPt}_6(2,6\text{-Me}_2\text{C}_6\text{H}_3\text{NC})_{12}]$ , **217**, 121
- solid-state isomerization of ruthenium(II) isocyanide complexes, **217**, 181
- infrared spectroelectrochemistry of  $[\text{Rh}(\text{TM}_4)_4\text{M}(\text{CO})_5]_2^{2+}$  (TM<sub>4</sub> = 2,5-diisocyno-2,5-dimethylhexane; M = Re, Mn): electron-transfer-promoted metal-metal bond cleavage and formation, **226**, 61
- Isomerization
- solid-state isomerization of ruthenium(II) isocyanide complexes, **217**, 181
- non-isothermal spectrophotometric kinetics applied to inorganic reactions, **221**, 31
- Isothiocyanato complexes
- isothiocyanato complexes of rhenium II. Synthesis, characterization and structures of  $\text{ReN}(\text{NCS})_2(\text{Me}_2\text{PhP})_3$  and  $\text{Re}(\text{NS})(\text{NCS})_2(\text{Me}_2\text{PhP})_3$ , **216**, 223
- isothiocyanato complexes of rhenium III. Synthesis and characterization of  $[\text{ReN}(\text{NCS})_2(\text{Ph}_3\text{P})_2]$  and  $[\text{ReN}(\text{NCS})_2(\text{CH}_3\text{CN})(\text{Ph}_3\text{P})_2]$ . X-ray crystal structure of  $[\text{ReN}(\text{NCS})_2(\text{CH}_3\text{CN})(\text{Ph}_3\text{P})_2]\cdot\text{CH}_3\text{CN}$ , **224**, 193
- Ketimido complexes

- synthesis of stable tail-to-tail-conjugated primary enamines containing hetero substituents, **222**, 193
- Ketone complexes**
- nucleophilic addition of water to coordinated di-(2-pyridyl)ketone (DPK) in rhenium(V) complexes. Synthesis and crystal structure of the adduct  $(\text{DPK} \cdot \text{H})^+ \cdot [\{\text{ReOCl}_2(\text{DPK} \cdot \text{OH})\}_2\text{Cl}]$ , **217**, 141
- Ketones**
- gold in organic synthesis Part 2. Preparation of benzyl-alkyl and -arylketones via C–C coupling, **220**, 1
- labeling studies of some carbonylation reactions of styrene with cationic palladium complexes, **220**, 273
- hydration of conjugated dienes to produce ketones catalyzed by ruthenium complexes, **222**, 225
- Kinetics**
- hydrolytic cleavage of peptides by palladium(II) complexes is enhanced as coordination of peptide nitrogen to palladium(II) is suppressed, **217**, 21
- stereoselectivity in reactions of metal complexes Part XVIII. Kinetics and stereoselectivity in electron-transfer reactions between  $[\text{Co}(\text{L})\text{H}_2\text{O}]^+$  ( $\text{L} = \text{N}, \text{N}'$ -[(pyridine-2,6-diyl)bis(methylene)]bis[proline] (promp) and  $\text{N}, \text{N}'$ -[(4-methoxy-pyridine-2,6-diyl)bis(methylene)]bis[proline] (MeO-promp) and optically active iron(II) complexes, **218**, 73
- effect of substituting sulfur for oxygen on the heterogeneous electron transfer kinetics of oxomolybdenum(V) hydrotris-(3,5-dimethyl-1-pyrazolyl)borate complexes, **225**, 1
- dependence of electron transfer kinetics of the ferrocene/ferrocenium couple on the viscosity in ambient temperature chloroaluminate ionic liquids, **225**, 9
- the kinetics of electron-transfer reactions of the  $[\text{FeCp}(\text{CpCH}_2\text{N}(\text{CH}_3)_3)^{+/2+}]$  couple in the presence of cyclodextrins in aqueous media, **225**, 51
- volume profile analysis for intermolecular electron transfer between cytochrome *c* and  $\text{Co}(\text{terpy})_2$ , **225**, 95
- redox reactions between oxomolybdenum(IV) and tetratolylporphinatoiron(III) complexes, **226**, 259
- Kinetics and mechanism**
- metal complexes of benzodiazepines. Kinetic properties of bromazepam toward square planar dimethyl sulfoxide complexes of platinum(II) and palladium(II), **215**, 41
- synthesis and hydrolysis of an ester of the 'one ring open' hydrolysis product of the anticancer drug 1,2-bis(3,5-dioxopiperazin-1-yl)propane, 'Razoxane', **215**, 55
- the complexation of divalent metal ions by the cryptands 4,7,13,16-tetraoxa-1,10-diazabicyclo[8.8.2]eicosane, -[8.8.5]tricosane and -[8.8.8]hexacosane in aqueous solution, **215**, 79
- kinetics and mechanism of the phosphate-catalyzed heterolysis of the metal-carbon  $\sigma$  bond of  $\alpha$ -hydroxyalkyl-chromium(III) complexes. Effects of the non-participating ligands, temperature and pressure, **215**, 173
- unusually stable peroxocopper complexes. Stoichiometry, products and kinetics of oxidation of the dimeric copper(I) complex  $[\text{LCuBr}]_2$  ( $\text{L} = \text{N}, \text{N}'$ -diethylethylenediamine) by dioxygen in methylene chloride from  $-51$  to  $30$  °C, **217**, 109
- ternary Pt(II)-amino acid-nucleotide complexes: kinetics of formation, **218**, 11
- novel metal thiolates:  $[\text{Mn}_2(\text{SC}_6\text{H}_3\{\text{SiMe}_3\}_2)_4(\text{C}_4\text{H}_8\text{O})]$ , a binuclear manganese thiolate complex with tetrahedral and trigonal-planar metal coordination, **219**, 1
- equilibrium and kinetic study of piperidine binding to phthalocyaninatoiron(II) in dimethyl sulfoxide, **219**, 199
- the steric effect on the ring opening process of the decarboxylation of *cis*-carbonato-bis(diamine)cobalt(III) ions, **221**, 9
- the decomposition of peroxomonosulfate induced by manganese(II) in acid solution, **221**, 25
- non-isothermal spectrophotometric kinetics applied to inorganic reactions, **221**, 31
- steric and electronic tuning of Michael addition reactions induced by pentacarbonyl metal fragments, **222**, 207
- arene migration in ruthenium clusters: a kinetic study of the isomerisation of  $\text{Ru}_5\text{C}(\text{CO})_{12}(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_6)$  to  $\text{Ru}_5\text{C}(\text{CO})_{12}(\eta^6\text{-C}_6\text{H}_6)$ , **222**, 299
- electron transfer 122. Reductions of an  $\text{N}_4$ -coordinated complex of silver(III), **225**, 83
- kinetics and mechanism of the ferrate oxidation of hydrazine and monomethylhydrazine, **225**, 145
- electron transfer reactivity of cuboidal heterometallic  $\text{Mo}_3\text{MS}_4$  clusters in aqueous acidic solutions, **225**, 157
- comparison of the rates of substitution in  $[\text{Ru}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  and  $[\text{Os}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  by hexacyano complexes: substitution coupled to electron transfer, **225**, 173
- the reduction of pentaammine(phenanthroline-2-carboxamido-*N*)cobalt(III) by titanium(III) and vanadium(II), **225**, 245
- mesoferriheme regeneration from its hemoprotein enzyme compound I analog: absorption and kinetic properties of one- and two-electron oxidation products, **225**, 255
- electron transfer 120. Reductions of the cuboidal mixed-valence molybdenum/sulfur cluster,  $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]$ , **225**, 297
- intramolecular electron transfer between Ru(I) and Ru(III) and the heme iron of cytochrome *c* labeled with ruthenium(II) polypyridine complexes, **226**, 129
- electron transfer reduction of cobalt tetraphenylporphyrin by hydroquinone dianions and alkylation with alkyl halides, **226**, 145
- dynamical investigations of the catalytic mechanisms of water oxidation by the  $[(\text{bpy})_2\text{Ru}(\text{OH}_2)]_2\text{O}^{4+}$  ion, **226**, 179
- the synthesis, X-ray structure and hydrolysis kinetics of chloro(2-aminomethylpyridine)(triamine)chromium(III) salts, **227**, 11
- substitution kinetics of some tetracarbonyl( $\eta^2$ -alkene)osmium complexes, **227**, 215
- thermal decomposition of cobalt carbonyl complexes in viscous media, **227**, 233
- kinetics and mechanism of the methanolysis of acetylcobalt tetracarbonyl, **227**, 297
- Kinetics of reduction**
- stereoselective electron transfer between Cu(II) superoxide dismutase and  $\Lambda$ - and  $\Delta$ -[Fe(pdta)], **225**, 151
- Lactone complexes**
- the influence of aluminum trichloride on a configuratively labile lactone-bridged biaryl: quantum chemical calculations and optical spectroscopy, **222**, 247
- synthesis, structure, dynamics and first atropisomer-selective cleavage of the chromium tricarbonyl complex of a lactone-bridged biaryl, **222**, 255
- Lanthanide complexes**
- systematics of some properties of hexamolybdotellurate(VI) salt hydrates with di- and trivalent metals,  $\text{M}_6[\text{TeMo}_6\text{O}_{24}] \cdot n\text{H}_2\text{O}$ , **217**, 93
- stability constants of the lanthanide(III)-1,4,7,10-tetraazacyclododecane-*N,N',N'',N'''*-tetraacetate complexes, **221**, 165
- synthesis, structure and stability of complexes of lanthanide nitrates with  $[(\text{MeO})_2\text{P}(\text{O})]_2\text{C}(\text{OH})\text{Ph}$  and the isomeric  $(\text{MeO})_2\text{P}(\text{O})\text{OCHPhP}(\text{O})(\text{OMe})_2$ , **223**, 43
- Lanthanide shift reagent**
- an europium(III) complex as an efficient catalyst for the Michael reaction of  $\alpha,\beta$ -enones with ketene silyl acetals: LIS-NMR analysis for the transition state through complexation

- between europium(III) catalyst and enones, **222**, 377
- Lead complexes**  
 synthesis and characterization of lead(II)–semicarbazone complexes. Crystal structures of dinitrato di-2-pyridylketone semicarbazone lead(II) and dichloro cyclohexane-1,2-dione bis(semicarbazone) dimethylformamide lead(II), **215**, 85  
 azacryptand host for lead: NMR and structural studies, **218**, 5  
 use of aluminium as an electron pool. Reductive coupling of iodobenzenes promoted by electron transfer in an Ni/Pb/Al multi redox system, **222**, 323
- Lead ions**  
 metal–metal interactions of aqueous lead(II) and tin(II) ions with the ground and excited states of tetrakis( $\mu$ -diphosphito)diplatinate(II), **225**, 201
- Ligand binding**  
 effect of aromatic–aromatic interaction on ligand binding to zinc porphyrins, **224**, 157
- Light-induced exchange**  
 light induced exchange energy transfer reactions of micelle forming Ru(II) diimine complexes, **225**, 275
- Liquid crystal**  
 synthesis of symmetric and unsymmetric 1,4-bis(*p*-R-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289
- Lithium complexes**  
 lithium coordination to amino acids and peptides. Synthesis, spectroscopic characterization and structure determination of lithium complexes of neutral and anionic glycine and diglycine, **218**, 121  
 structures of isomeric substituted 14-crown-4 ethers and their LiSCN complexes, **223**, 125
- Lithium compounds**  
 benzotriazole stabilized lithium intermediates as a route to the elaboration of N-substituents in amides, **220**, 67
- Loop-current effect**  
 a comparison of the loop-current effect of silicon phthalocyanine and silicon naphthalocyanine rings on their axial ligands, **218**, 41
- Low-valent complexes**  
 reaction of low-valent iron porphyrins with alkyl containing supporting electrolytes, **226**, 195
- Luminescence**  
 luminescent europium(III) and terbium(III) ions encapsulated in a 2-aminoethoxy or carbamoyloxy substituted calixarene host, **227**, 293
- Luminescence quenching**  
 oxidative quenching of luminescence from copper metallothionein, **226**, 275
- Macrocates**  
 photophysical properties of mono-, di- and tetranuclear copper(I)–polypyridine complexes, **225**, 251
- Macrocyclic complexes**  
 a novel sodium template approach for preparing tetraimine macrocycles of 2,6-diformyl-4-methylphenol and diamino derivatives, **223**, 169
- Macrocyclic ligand complexes**  
 redox reactions of an Mo(V) tetrasulphophthalocyanine, **215**, 27  
 spectroscopic studies on copper(II) complexes of chiral cyclens: [CuN<sub>4</sub>Cl] chromophores varying from square pyramidal to trigonal bipyramidal stereochemistry, **215**, 109  
 bulky substituent effects on the iron(III) complexation of 1,4,7-triazacyclononane, **216**, 89  
 synthesis, spectral properties and redox behaviour of oxovanadium(IV) complexes with tetraaza[14]annulenes, **216**, 217
- heterodinuclear Ni(II)M(II) (M=Pb, Mn, Fe, Co, Ni, Cu, Zn) complexes of a phenol-based dinucleating macrocycle with dissimilar 4- and 5-coordination sites, **218**, 27
- liquid matrix induced processes in fast atom bombardment mass spectrometry of polyazacrown transition metal complexes, **219**, 85
- solution studies of monovalent metal complex ions of the bi-brachial lariat ether 7,16-bis(2-methoxyethyl)-1,4,10,13-tetraoxa-7,16-diazacyclooctadecane, **219**, 217
- methane as the product of reaction of methyl-coenzyme-M with monovalent nickel complexes in aqueous solutions. A model for the in vivo activity of cofactor F<sub>430</sub>, **227**, 1
- equilibrium studies of 1,2-diaminobenzene and 3,4-diaminobenzoate anion with dinuclear cobalt(II) 1,4,7,13,16,19-hexaaza-10,22-dioxacyclotetracosane, **227**, 49
- outer-sphere coordination of polycyanometallate anions with polyammonium macrocycles: a spectrophotometric study, **227**, 71
- Macrocyclic pseudopeptide**  
 coordination abilities of amide oxygens in a 24-membered ring pseudopeptide toward several transition metal ions in acetonitrile, **223**, 121
- Macropolygons**  
 metal cluster topology 16. Macropolygons and macropolyhedra in coinage metal iron carbonyl cluster anions and related compounds, **227**, 207
- Macropolyhedra**  
 metal cluster topology 16. Macropolygons and macropolyhedra in coinage metal iron carbonyl cluster anions and related compounds, **227**, 207
- Magnesium complexes**  
 synthesis, crystal structure and thermal stability of the MgCl<sub>2</sub> formamide adduct, **216**, 261  
 synthesis, structure and solid state polymerization of diaqua-bis(propynoato)magnesium, **222**, 339
- Magnesium compounds**  
 magnesium bromide in Grignard reagent formation, **222**, 365
- Magnetism**  
 {[Ni(3,2,3-tet)( $\mu$ -NCO)]<sub>n</sub>}(ClO<sub>4</sub>)<sub>n</sub>: structural characterization of the first monodimensional nickel(II)–cyanato system with a description of its ferromagnetic coupling, **216**, 5  
 a polymeric two-dimensional mixed-metal network. Crystal structure and magnetic properties of {[P(Ph)<sub>4</sub>]-[MnCr(ox)<sub>3</sub>]}<sub>n</sub>, **216**, 65  
 a seven-spin cluster formed by an alkyl nitronyl nitroxide biradical and copper(II): crystal structure and magnetic properties, **217**, 7  
 preparation, crystal structures and magnetic properties of a *syn-anti* carboxylate-bridged dinuclear copper(II) complex and its precursor, a hydrogen-bonded polynuclear copper(II) complex, **218**, 109  
 crystal structure and magnetic properties of the first 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position: {catena-( $\mu_{N,S}$ -NCS)[Ni(en)<sub>2</sub>]}(PF<sub>6</sub>), **218**, 185  
 crystal structure and magnetic property of a 2D infinite layered copper(II) complex with bridging *trans*-oxamidate and pyrimidine ligands, **224**, 171
- Main group elements**  
 charge and electron transfer from metal-to-carbon bonds of main group organometallics MR<sub>n</sub> (M=Al, Ga, Zn) to aromatic N-heterocycles: colored precursor compounds and radical complex formation, **225**, 23
- Maleoyl complexes**  
 the development of a quinone  $\alpha$ -anion synthon. Utilization of a maleoylcobalt complex as a quinone surrogate and a dominant role for ligand effects, **222**, 235
- Manganese complexes**

- a polymeric two-dimensional mixed-metal network. Crystal structure and magnetic properties of  $\{[P(Ph)_4]_n[MnCr(ox)_3]_n\}$ , **216**, 65
- coordination chemistry of perhalogenated cyclopentadienes and alkynes XIV. Synthesis of dinuclear ring-bridged cyanotrenes with  $-SiMe_2-$  or  $-TiCp_2-$  bridges. Structures of  $[(OC)_3Mn(C_5Br_4-)]_2SiMe_2$  and  $[(OC)_3Mn(C_5Cl_4-)]_2TiCp_2$ , **218**, 173
- novel metal thiolates:  $[Mn_2(SC_6H_3\{SiMe_3\}_2)_4(C_4H_8O)]$ , a binuclear manganese thiolate complex with tetrahedral and trigonal-planar metal coordination, **219**, 1
- preparation, properties and coordination behaviour of planar or tridimensional compartmental Schiff bases, **219**, 135
- coordination polymers of Mn(II) with the ligand pyrazine-2,3,5,6-tetracarboxylic acid, **219**, 161
- variation of the exchange interaction in oximate-bridged  $Cu^{II}M^{II}$  dimers ( $M = Cu, Ni, Mn$ ). Crystal structure of  $[Cu(pdmg)Cu(bipy)(H_2O)_2](ClO_4)_2 \cdot H_2O$ , **219**, 179
- photochemical reactions of transition metal organyl complexes with olefins Part 10. Light-induced formal  $[5+2]$  cycloadditions at manganese, **220**, 77
- optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220**, 283
- the synthesis and reactions of A-ring aromatic steroid complexes of manganese tricarbonyl, **220**, 297
- the decomposition of peroxomonosulfate induced by manganese(II) in acid solution, **221**, 25
- an investigation of electron transfer properties of  $(Cp)Mn(CO)_3$  and substituted Cp derivatives, **221**, 85
- nickel(II) and manganese(II) complexes of substituted phenanthroline ligands, **221**, 99
- manganese and tungsten carbene complex mediated intramolecular Diels–Alder reactions, **222**, 291
- di-manganese carbonyl compounds containing bis(dicyclohexylphosphino)methane (dcpm) ligation, **224**, 105
- crystal and molecular structures of  $\{[exo\text{- and }endo\text{-}(RO)_2P(O)\eta^5\text{-}C_6H_6\}Mn(CO)_3\}$ , **224**, 147
- synthesis and electrochemical properties of oxo-bridged manganese dimers incorporating alkali and alkaline earth cations, **225**, 191
- ligand substitution at 19-electron organometallic centers. Electrocatalytic CO substitution reactions of (methylcyclopentadienyl) $Mn(CO)_2NO^+$  and (indenyl) $Mn(CO)_2NO$ , **226**, 53
- valence tautomerism for catechol/semiquinone complexes of the *trans*- $M(Bupy)_2(3,6\text{-}DBQ)_2$  ( $M = Mn, Fe, Co$ ) series, **226**, 91
- one-equivalent oxidative charge storage on an  $[Mn_2O_2]^{3+}$  core: synthesis, crystal structure and resonance Raman spectra of  $[Mn_2O_2(pbz)_4]^{n+}$  ( $n = 3, 4$ ), **227**, 33
- Mass spectrometry**
- an electrospray mass spectrometric study of methylmercury(II) nitrogen base derivatives and their ligand exchange reactions, **215**, 179
- complexes of tetradentate  $\beta$ -keto enamine Schiff bases: an electrospray mass spectrometric study, **216**, 75
- Mercuric ion complexes**
- interaction of cyanopentaamminechromium(III) with mercuric ion. Electrochemical, spectral and preparative study of adduct formation and linkage isomerism, **216**, 113
- Mercury complexes**
- an electrospray mass spectrometric study of methylmercury(II) nitrogen base derivatives and their ligand exchange reactions, **215**, 179
- mercury and platinum mixed cluster having a prismatic structure,  $[HgPt_6(2,6\text{-}Me_2C_6H_3NC)_2]$ , **217**, 121
- complexes of mercuric halides with the tripod ligand  $np_3$  ( $= N(CH_2CH_2PPh_2)_3$ ). Molecular structure of  $[(np_3)HgI] \cdot 0.5C_2H_5OH$ , **217**, 155
- photoredox chemistry of iodomercurate(II) complexes in acetonitrile, **221**, 79
- alkyne hydration using Pt(II) catalysts, **222**, 317
- positive and negative ion electrospray mass spectrometric studies of some amino acids and glutathione, and their interactions with alkali metal ions and methylmercury(II), **223**, 103
- multinuclear ( $^{31}P$ ,  $^{77}Se$ ,  $^{113}Cd$ ,  $^{199}Hg$ ) magnetic resonance, electrospray mass spectrometric and electrochemical studies on the reactions of  $M(S_2P\{OEt\}_2)_2$  ( $M = Cd, Hg$ ) with potentially bidentate mixed Group 15/Group 16 and Group 16 donor ligands, **224**, 137
- preparation, structural and physical studies of mercury(II) halide complexes with the  $N,N'$ -chelating biheteroaromatic ligand 2-(2'-pyridyl)quinoxaline, **227**, 129
- synthesis and structural analysis of *trans*- $Cp(CO)_2(PPh_3)WHgCl$ , **227**, 145
- Mesoferrheme regeneration**
- mesoferrheme regeneration from its hemoprotein enzyme compound I analog: absorption and kinetic properties of one- and two-electron oxidation products, **225**, 255
- Metal cluster topology**
- metal cluster topology 16. Macropolygons and macropolyhedra in coinage metal iron carbonyl cluster anions and related compounds, **227**, 207
- Metal complexes**
- principal component analysis of the structures of the 1,2-diaminobenzene derivatives chelated to metal ions: correlation between structural parameters and redox behavior, **215**, 219
- steric and electronic tuning of Michael addition reactions induced by pentacarbonyl metal fragments, **222**, 207
- stabilities of 1,2-diethyl-3-hydroxy-4-pyridinone chelates of divalent and trivalent metal ions, **223**, 21
- metal complexes of some 2-nitrosoanilides, **223**, 71
- charge and electron transfer from metal-to-carbon bonds of main group organometallics  $MR_n$  ( $M = Al, Ga, Zn$ ) to aromatic N-heterocycles: colored precursor compounds and radical complex formation, **225**, 23
- electron transfer in the electrochemistry of metals, metal compounds and metal complexes, **226**, 99
- quenching of a polypyridyl-ruthenium(II) chromophore by covalently attached  $\{ML(NO)Cl\}$  fragments ( $M = Mo, W$ ;  $L = \text{tris}(3,5\text{-dimethylpyrazolyl})\text{hydroborate}$ ), **226**, 171
- metal cluster topology 16. Macropolygons and macropolyhedra in coinage metal iron carbonyl cluster anions and related compounds, **227**, 207
- Metal compounds**
- electron transfer in the electrochemistry of metals, metal compounds and metal complexes, **226**, 99
- Metal hydrides**
- direct solution calorimetric measurements of enthalpies of proton and electron transfer reactions for transition metal complexes. Thermochemical study of metal–hydride and metal–metal bond energies, **227**, 285
- Metal ion complexes**
- the complexation of divalent metal ions by the cryptands 4,7,13,16-tetraoxa-1,10-diazabicyclo[8.8.2]eicosane, -[8.8.5]tricosane and -[8.8.8]hexacosane in aqueous solution, **215**, 79
- positive and negative ion electrospray mass spectrometric studies of some amino acids and glutathione, and their interactions with alkali metal ions and methylmercury(II), **223**, 103



- synthesis and characterization of metal complexes with *N,N'*-bis(2-hydroxybenzyl)-diethylenetriamine-*N,N',N''*-triacetic acid and the stabilities of its complexes with divalent and trivalent metal ions, **223**, 109
- stability of metal ion complexes of *N,N'*-bis(2-hydroxybenzyl)-ethylenediamine-*N,N'*-diacetic acid, **224**, 151
- Metallacyclophane complexes**
- preparation, properties and reactions of metal-containing heterocycles Part LXXXIX. Metalla[*n*]- and metalla[*m,n*]-orthocyclophanes as mediators for the synthesis of cyclic ketones, **220**, 107
- Metalloproteins**
- chemical and electrochemical approaches to the investigation of redox reactions of simple electron transfer metalloproteins, **226**, 293
- Metallosupramolecular chemistry**
- a new functionalised oligopyridine ligand containing ferrocene as a ball-bearing spacer for metallosupramolecular chemistry, **224**, 11
- Metal–metal bond energies**
- direct solution calorimetric measurements of enthalpies of proton and electron transfer reactions for transition metal complexes. Thermochemical study of metal–hydride and metal–metal bond energies, **227**, 285
- Metal–metal interactions**
- metal–metal interactions of aqueous lead(II) and tin(II) ions with the ground and excited states of tetrakis( $\mu$ -diphosphito)diplatinate(II), **225**, 201
- Metals**
- electron transfer in the electrochemistry of metals, metal compounds and metal complexes, **226**, 99
- Metal thiolates**
- novel metal thiolates:  $[\text{Mn}_2(\text{SC}_6\text{H}_3(\text{SiMe}_3)_2)_4(\text{C}_4\text{H}_8\text{O})]$ , a binuclear manganese thiolate complex with tetrahedral and trigonal-planar metal coordination, **219**, 1
- Methanolysis**
- kinetics and mechanism of the methanolysis of acetylcobalt tetracarbonyl, **227**, 297
- Methylacetoacetate complexes**
- intermediates in the palladium-catalysed reaction of 1,3-dienes Part 8. The reaction of palladium–butadiene complexes with ethyl methylacetoacetate, **222**, 179
- Methyl complexes**
- the equilibrium between acetyl- and methylcobalt tetracarbonyl, **227**, 211
- Micelle complexes**
- light induced exchange energy transfer reactions of micelle forming Ru(II) diimine complexes, **225**, 275
- Michael reactions**
- steric and electronic tuning of Michael addition reactions induced by pentacarbonyl metal fragments, **222**, 207
- an europium(III) complex as an efficient catalyst for the Michael reaction of  $\alpha,\beta$ -enones with ketene silyl acetals: LIS-NMR analysis for the transition state through complexation between europium(III) catalyst and enones, **222**, 377
- Mixed-ligand complexes**
- spectroscopic investigation of mixed-ligand nickel(II) chelates containing  $\beta$ -ketoenols and 1,2-diamines, **221**, 17
- Mixed-ligand tetrairon–sulfur cluster**
- syntheses and structures of mixed-ligand tetrairon–sulfur clusters,  $[\text{Cp}'_2(\text{Ph}_2\text{C}_2\text{S}_2)_2\text{Fe}_4\text{S}_4](\text{PF}_6)_n$  and  $[\text{Cp}'_3(\text{Ph}_2\text{C}_2\text{S}_2)\text{Fe}_4\text{S}_3](\text{PF}_6)_n$  ( $n=1$  and  $2$ ) ( $\text{Cp}' = \eta^5\text{-C}_5\text{Me}_5$ ). Structural change of the clusters accompanying electron transfer reactions, **225**, 229
- Mixed-metal carbonyl clusters**
- Solution structures of cobalt–rhodium clusters: the identification of a new hexanuclear cobalt–rhodium species, **227**, 187
- Mixed salt complexes**
- a mixed-metal salt comprised of metal–metal bonded dinuclear ions: structure and properties of  $[\text{Rh}_2(\text{OAc})_2(\text{MeCN})_6]\text{-}[\text{Re}_2\text{Cl}_8]$ , **217**, 79
- Mixed-valence complexes**
- synthesis and characterisation of cyanide bridged heterobinuclear complexes based on Group VI octacyanometalate(IV) and pentaammineosmium(III) ions: a comparison with Group VIII hexacyano analogues, **219**, 121
- properties of the mixed-valence binuclear complex ion,  $[(\text{NH}_3)_5\text{Ru}^{\text{III}}\text{-NC-OS}^{\text{I}}(\text{CN})_5]$ , **223**, 37
- mixed valence state associated with halogen migration in diruthenocymethane and its related compounds, **225**, 103
- solvent trapped valency in the [bis(pentaammineruthenium)-( $\mu$ -1,4-dicyanamidobenzene)] trication, **225**, 123
- electron transfer 120. Reductions of the cuboidal mixed-valence molybdenum/sulfur cluster,  $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]$ , **225**, 297
- synthesis, electrochemistry and photoexcited-state properties of dinuclear ruthenium complexes bridged by 2,6'-bis(2-pyridyl)-2,2':6,2''-thiazolo[4,5-*d*]-benzothiazole, **226**, 17
- solvent, electrolyte and solute shape effects on optical electron transfer in mixed-valence ruthenium ammine dimers, **226**, 137
- Mixed-valence systems**
- the influence of spin-forbidden monomer excitations on spin-allowed electron transfer and electron-localized states of mixed-valence and single-valence dimeric systems, **226**, 33
- MOCVD application**
- a novel route to the second-generation alkaline-earth metal precursors for metal-organic chemical vapour deposition: one-step synthesis of  $\text{M}(\text{hfa})_2 \cdot \text{tetraglyme}$  ( $\text{M} = \text{Ba}, \text{Sr}, \text{Ca}$  and  $\text{Hhfa} = 1,1,1,5,5,5\text{-hexafluoro-2,4-pentanedione}$ ), **224**, 203
- Molecular oxygen**
- optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220**, 283
- activation of molecular oxygen by iron nitrosyls in the presence of the bidentate phosphines 1,2-bis(diphenylphosphino)ethane and ethene, **221**, 133
- Molecular recognition**
- a novel amino acid-binding receptor based on weak non-covalent ligand–ligand interactions, **216**, 249
- Molecular structure**
- EXAFS studies of a chiral rhenium lactone complex and its precursors including multiple scattering calculations, **222**, 27
- Molybdate complexes**
- coordination study of D-xylo 5-hexulosonic acid with borate, molybdate and tungstate in aqueous solution by nuclear magnetic resonance spectroscopy, **221**, 69
- Molybdenum cluster metallates**
- ionic cluster dithiophosphinates  $[\text{Mo}_3\text{S}_4(\text{R}_2\text{PS}_2)_3\text{py}_3]^+ [\text{MI}_3]^-$  ( $\text{M} = \text{Zn} \cdot \text{py}, \text{Cd} \cdot \text{py}, \text{Hg}$ ), **216**, 177
- Molybdenum complexes**
- preparation and structural characterization of  $\text{Rh}_2(\text{O}_2\text{CCPh}_3)_4(\text{EtOH})_2$ ,  $\text{Ru}_2(\text{O}_2\text{CCPh}_3)_4(\text{H}_2\text{O}) \cdot (\text{EtOH}) \cdot 2\text{EtOH}$  and  $\text{Mo}_2(\text{O}_2\text{CCPh}_3)_4 \cdot 3\text{CH}_2\text{Cl}_2$ , **215**, 9
- redox reactions of a Mo(V) tetrasulfophthalocyanine, **215**, 27
- synthesis and characterization of Keggin derivatives containing an  $[\text{Mo}(\text{NO})]^{3+}$  unit:  $(n\text{-Bu}_4\text{N})_4[\text{PM}_{11}\text{O}_{39}\{\text{Mo}(\text{NO})\}]$  ( $\text{M} = \text{Mo}, \text{W}$ ), **215**, 61
- the unusual crystallographic disorder of  $\text{M}_2$  units in  $\text{M}_2\text{Cl}_4(\text{PEt}_3)_4$  ( $\text{M} = \text{Re}, \text{Mo}$ ), **215**, 103

- preparation and characterization of  $[(LL)Cl_2Mo]_2(\mu-SPr)_2$  compounds with  $Pr = n$ -propyl and  $LL = PrSCH_2CH_2SPr$  and  $Et_2PCH_2CH_2PEt_2$ . Further examples of C–S bond cleavage and oxidative addition across an Mo–Mo quadruple bond, **215**, 191
- oxo- and sulfido-bridged sandwich cubane-type molybdenum–indium mixed-metal cluster. Synthesis and X-ray structure of  $[(H_2O)_9Mo_3S_3OInOS_3Mo_3(H_2O)_9] \cdot (CH_3 \cdot C_6H_4 \cdot SO_3)_8 \cdot 30H_2O$ , **216**, 13
- spectroscopic investigation of dichloro and hydrazido(2–) derivatives of bis(dithiocarbamato)dioxomolybdenum(VI) compounds. Crystal and molecular structure of bis(diisobutylidithiocarbamato)(*N,N*-dimethylhydrazido(2–)-*N'*)-oxomolybdenum(VI), **217**, 61
- systematics of some properties of hexamolybdotellurate(VI) salt hydrates with di- and trivalent metals,  $M_4[TeMo_6O_{24}] \cdot nH_2O$ , **217**, 93
- molybdenum carbonyl derivatives containing eight- and nine-membered diphosphine chelating rings: the synthesis of  $[Mo(CO)_4(dppp)_2]$  ( $dppp_e = Ph_2P(CH_2)_5PPh_2$ ) and  $[Mo(CO)_4(dpph)]$  ( $dpph = Ph_2P(CH_2)_6PPh_2$ ) and the crystal structure of the latter, **218**, 9
- structural characterization of octahedral metal complexes of  $[MX_3(py)_3]$  type with three pyridines as revealed by  $^2H$  NMR spectra, **218**, 179
- synthesis and characterization of molybdenum and tungsten nitride complexes of the type  $[M(NO_2)_2(CO)_2(PPh_3)_2]$ , **218**, 195
- dimolybdenum(II) and diruthenium(II) complexes with 2-substituted 7-methyl-1,8-naphthyridines as bridging ligands, **219**, 23
- the study of isolobal displacement reactions. Synthesis and structures of the heterobimetallic  $\mu$ -alkyne cluster complexes  $\eta^5-R_2C_5H_4(CO)_2M(\mu-R^1C_2R^2)Co(CO)_3$  ( $M = Mo, W$ ;  $R = CH_3C(O), CH_3O_2C$ ), **219**, 93
- synthesis and characterisation of cyanide bridged heterobinuclear complexes based on Group VI octacyanometalate(IV) and pentaammineosmium(III) ions: a comparison with Group VIII hexacyano analogues, **219**, 121
- preparation of  $\eta^1$ -1,3-dienyl,  $\eta^1$ -1,2-dienyl and  $\eta^3$ -cyclobutene complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305
- $N^1, N^7$ -expeditious dialkylation of cyclen (1,4,7,10-tetraazacyclododecane). An astonishing reactivity of cyclen tricarbonyl molybdenum and chromium complexes, **220**, 347
- molybdenum complexes with sterically-hindered thio-carboxylate ligands. Electrochemical properties of the anionic complex bis(2,2-diphenyl-2-mercaptoethanoate)dioxomolybdate(VI) in protic solvents, **221**, 117
- synthesis and X-ray structural determination of pentamethylcyclopentadienyl dichloro oxo molybdenum, **221**, 147
- carbonyl carbene coupling on cationic  $\eta^2$ -carbene complexes of molybdenum and tungsten, **222**, 77
- extrusion of molecular clusters from solid-state materials: synthesis by application of  $\gamma$ -irradiation. Molecular and crystal structure of  $(H_9O_4)(Et_4N)[Mo_3S_7Br_6]$ , **223**, 5
- synthesis and characterization of the new peroxo complex aquaoxidiperoxoimidazole molybdenum(VI), **223**, 173
- phenylimido derivatives of  $[Mo_6O_{19}]^{2-}$ : syntheses, X-ray structures, vibrational, electrochemical,  $^{95}Mo$  and  $^{14}N$  NMR studies, **224**, 81
- effect of substituting sulfur for oxygen on the heterogeneous electron transfer kinetics of oxomolybdenum(V) hydrotris-(3,5-dimethyl-1-pyrazolyl)borate complexes, **225**, 1
- electron transfer reactivity of cuboidal heterometallic  $Mo_3MS_4$  clusters in aqueous acidic solutions, **225**, 157
- transition metal complexes with sulfur ligands Part CVII. Complexes of the  $[Mo(NO)(S_4)]$  fragment with 16, 17 and 18 valence electrons ( $S_4^{2-} = 1,2$ -bis(2-mercaptophenylthio)ethane(2–)), **225**, 163
- electron transfer 120. Reductions of the cuboidal mixed-valence molybdenum/sulfur cluster,  $[Mo_4S_4(H_2O)_{12}]$ , **225**, 297
- crystal structure and solid-state fluorescence of an oxomolybdenum(V)–benzylviologen electron donor–acceptor salt, **226**, 25
- $^{95}Mo$  NMR spectroscopy
- phenylimido derivatives of  $[Mo_6O_{19}]^{2-}$ : syntheses, X-ray structures, vibrational, electrochemical,  $^{95}Mo$  and  $^{14}N$  NMR studies, **224**, 81
- Mössbauer spectroscopy
- 'insulation' to lattice effects on quadrupole splitting in the Mössbauer spectra of salts of  $[Fe_4S_4(Cys)_4]^{2-}$ , a ferredoxin model, **217**, 3
- Mössbauer spectroscopy of iron dextran: comparison of solid and frozen solution, **223**, 9
- Multidentate ligand complexes
- an ethyl–cobalt complex with a buckled equatorial ligand, **221**, 35
- metal complexes of 5-acetyl-2,2'-pyridylthiophene arylhydrazones. Crystal structure of a cobalt(II) derivative, **223**, 97
- synthesis and characterization of metal complexes with *N,N'*-bis(2-hydroxybenzyl)-diethylenetriamine-*N,N',N''*-triacetic acid and the stabilities of its complexes with divalent and trivalent metal ions, **223**, 109
- fluorine-substituted tetraazaannulene complexes: syntheses and characterizations, **223**, 187
- synthesis, spectra and electrochemical behaviour of biomimetic copper(II) complexes with  $CuN_5$  and  $CuN_6$  chromophores, **224**, 19
- electron transfer properties of  $ReO(ECD)$ , **225**, 323
- Multi metal redox system
- use of aluminium as an electron pool. Reductive coupling of iodobenzenes promoted by electron transfer in an Ni/Pb/Al multi redox system, **222**, 323
- Multinuclear NMR
- multinuclear ( $^{31}P$ ,  $^{77}Se$ ,  $^{113}Cd$ ,  $^{199}Hg$ ) magnetic resonance, electrospray mass spectrometric and electrochemical studies on the reactions of  $M(S_2P(OEt)_2)_2$  ( $M = Cd, Hg$ ) with potentially bidentate mixed Group 15/Group 16 and Group 16 donor ligands, **224**, 137
- Neighbouring-group effect
- chele- and anticheleselectivity IV. Reaction-accelerating neighboring-group effects in the cross-coupling of vinylbromides with alkyl-transition metal reagents, **220**, 85
- N-heterocycle complexes
- synthesis of  $\eta^3$ -allyl–Pd complexes and subsequent formation of N-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- Nickel
- EXAFS investigation on nickel substituted Cu,Zn bovine superoxide dismutase, **216**, 121
- Nickel complexes
- $\{[Ni(3,2,3-tet)(\mu-NCO)]_n\}(ClO_4)_n$ : structural characterization of the first monodimensional nickel(II)–cyanato system with a description of its ferromagnetic coupling, **216**, 5
- octahedral  $\mu$ -oxalato–nickel(II) dinuclear complexes with water and tridentate amines as blocking ligands: magnetostructural correlations, **216**, 139
- preparation and properties of a metal dithiolene conductor, partially oxidized salt of tetra(*n*-butyl)ammonium bis(5,6-dithio-benzo[*d*]-1,3-dithiole-2-thione) nickelate,  $(n-Bu_4N)_{0.29}[Ni(dmbit)_2]$ , **216**, 147

- geometrical and optical isomers of the dinickel complexes of two chiral ligands of the  $N,N'$ -disubstituted oxamide type: 1D and 2D NMR study, **216**, 237
- synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129
- heterodinuclear Ni(II)M(II) ( $M = \text{Pb, Mn, Fe, Co, Ni, Cu, Zn}$ ) complexes of a phenol-based dinucleating macrocycle with dissimilar 4- and 5-coordination sites, **218**, 27
- crystal structure and magnetic properties of the first 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position: {*catena*-( $\mu_{N,S}$ -NCS)[Ni(en)<sub>2</sub>]}(PF<sub>6</sub>), **218**, 185
- electrochemical and structural studies of nickel(II) complexes with N<sub>2</sub>O<sub>2</sub> Schiff base ligands 2. Crystal and molecular structure of *N,N'*-1,2-ethane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II), *N,N'*-1,2-*cis*-cyclohexane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II) and *N,N'*-1,2-benzene-1,2-diyl-bis(3,5-dichlorosalicylideneimine)nickel(II), **219**, 43
- variation of the exchange interaction in oximate-bridged Cu<sup>II</sup>M<sup>II</sup> dimers ( $M = \text{Cu, Ni, Mn}$ ). Crystal structure of [Cu(pdmg)Cu(bipy)(H<sub>2</sub>O)<sub>2</sub>](ClO<sub>4</sub>)<sub>2</sub>·H<sub>2</sub>O, **219**, 179
- spectroscopic investigation of mixed-ligand nickel(II) chelates containing  $\beta$ -ketoenols and 1,2-diamines, **221**, 17
- nickel(II) and manganese(II) complexes of substituted phenanthroline ligands, **221**, 99
- the effects of counterion and solvent on the reactivity of nickel acylate complexes, **222**, 165
- use of aluminium as an electron pool. Reductive coupling of iodobenzenes promoted by electron transfer in an Ni/Pb/Al multi redox system, **222**, 323
- fluorine-substituted tetraazaannulene complexes: syntheses and characterizations, **223**, 187
- predicting the effect of oxidative doping on the conductivities of metal-tetrakis(dimethylphosphino)benzene coordination polymers from the electrochemical properties of their related bimetallic complexes, **225**, 15
- methane as the product of reaction of methyl-coenzyme-M with monovalent nickel complexes in aqueous solutions. A model for the *in vivo* activity of cofactor F<sub>430</sub>, **227**, 1
- synthesis, crystal structure and characterization of nickel(0) acrylonitrile complexes with a bidentate N-donor ligand, **227**, 137
- alkylation of 2-vinylpyridine and crystal structure of bis{(acetylacetonate)(1-pyridylbutyl)nickel(II)} complex, **227**, 149
- structures of a monomeric and a dimeric Ni(II) complex containing 2-acetylpyrrole, **227**, 159
- the synthesis and crystal structure of ethylenedinitrilotetraacetamidoxime nickel(II) sulfate trihydrate, **227**, 171
- Niobium complexes**
- synthesis of some dinuclear niobium chloro alkoxy carboxylates [Nb<sub>2</sub>Cl<sub>4</sub>(OEt)<sub>4</sub>(O<sub>2</sub>CR)<sub>2</sub>], and the X-ray crystal and molecular structure of the compound with R = Ph, **227**, 99
- a new type of face-sharing bioctahedral triply bonded diniobium cluster (PyH)[Nb<sub>2</sub>( $\mu$ -Cl)<sub>2</sub>( $\mu$ -THT)Cl<sub>3</sub>(Py)<sub>3</sub>](THF)-(C<sub>6</sub>H<sub>6</sub>)<sub>0.5</sub>, made by a designed synthesis, **227**, 191
- Nitration reaction**
- direct nitration of aryl groups  $\sigma$ -bound to ruthenium(II), **220**, 261
- Nitrido complexes**
- mixed-ligand complexes of rhenium II. Synthesis and characterization of [ReN(X)(Me<sub>2</sub>PhP)<sub>2</sub>(R<sub>2</sub>tcb)] complexes (X = I, N<sub>3</sub>, SCN, CN; R<sub>2</sub>tcb<sup>-</sup> = *N*-(*N,N*-dialkylthiocarbamoylbenzamidate)). X-ray crystal structure of [ReN(N<sub>3</sub>)-(Me<sub>2</sub>PhP)<sub>2</sub>(Et<sub>2</sub>tcb)], **215**, 159
- mixed-ligand complexes of rhenium III. Synthesis, characterization and structures of [ReN(Me<sub>2</sub>PhP)<sub>2</sub>(mnt)] and [ReN(Me<sub>2</sub>PhP)(2,6-DMTP)(morph tcb)] (Me<sub>2</sub>PhP = dimethylphenylphosphine; 2,6-DMTP<sup>-</sup> = 2,6-dimethylthiophenolate; morph tcb<sup>-</sup> = *N*-(morpholinylthiocarbonyl)benzamidate), **216**, 31
- isothiocyanato complexes of rhenium III. Synthesis and characterization of [ReN(NCS)<sub>2</sub>(Ph<sub>3</sub>P)<sub>2</sub>] and [ReN(NCS)<sub>2</sub>(CH<sub>3</sub>CN)(Ph<sub>3</sub>P)<sub>2</sub>]. X-ray crystal structure of [ReN(NCS)<sub>2</sub>(CH<sub>3</sub>CN)(Ph<sub>3</sub>P)<sub>2</sub>]·CH<sub>3</sub>CN, **224**, 193
- the redox behaviour of the cluster anion [Fe<sub>4</sub>N(CO)<sub>12</sub>]<sup>-</sup>. Electron transfer chain catalytic substitution reactions. Crystal structure of [Fe<sub>4</sub>N(CO)<sub>11</sub>(PPh<sub>3</sub>)], **226**, 1
- synthesis and characterization of neutral oxorhenium(V) and nitridotecnium(V) complexes with a tetradentate N<sub>2</sub>S<sub>2</sub> unsaturated ligand derived from dithiocarboxylic acid, **227**, 25
- Nitrile complexes**
- uranium hexafluoride: laboratory scale synthesis, coordination behaviour towards nitriles and Raman spectra, **215**, 5
- molecular and crystal structures of *cis*-Pt(DMSO)(C<sub>2</sub>H<sub>5</sub>CN)Cl<sub>2</sub> and Cl(DMSO)Pt( $\mu$ -N(CH<sub>3</sub>)CHO)<sub>2</sub>Pt(DMSO)Cl, **216**, 163
- new polypyrazolylborate chemistry of ruthenium(II) complexes with weakly coordinating organonitrile ligands, **219**, 3
- synthesis of 5,6-dihydro-4*H*-1,3-oxazines from neutral and cationic platinum(II) nitrile complexes. X-ray structure of *trans*-[Pt(CF<sub>3</sub>){N=C(Ph)OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>}(PPh<sub>3</sub>)<sub>2</sub>]BF<sub>4</sub>, **220**, 21
- a comparison of catalytic activity for imine hydrogenation using Ru ditertiary phosphine complexes, including chiral systems, **222**, 85
- reactions of aziridine with platinum(II) nitriles. Formation of (aziridino)amidines and 2-imidazolines and X-ray structure of *trans*-[PtCl<sub>2</sub>{N(H)=C(Ph)NCH<sub>2</sub>CH<sub>2</sub>}]<sub>2</sub>, **222**, 327
- Nitrite complexes**
- synthesis and characterization of molybdenum and tungsten nitrite complexes of the type [M(NO<sub>2</sub>)<sub>2</sub>(CO)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>], **218**, 195
- Nitro complexes**
- direct nitration of aryl groups  $\sigma$ -bound to ruthenium(II), **220**, 261
- ligand nature of coordinated NO<sub>2</sub><sup>-</sup> in the oxidation reaction of Ru(II) complexes, [Ru(NO<sub>2</sub>)(OH<sub>2</sub>)(py)<sub>4-2n</sub>(bpy)<sub>n</sub>]<sup>+</sup> ( $n = 0, 1, 2$ ), and their related complexes, **225**, 111
- Nitronyl complexes**
- a seven-spin cluster formed by an alkyl nitronyl nitroxide biradical and copper(II): crystal structure and magnetic properties, **217**, 7
- Nitrosoalkane complexes**
- nitrosoarene and nitrosoalkane insertion reactions of titanacyclobutenes, **222**, 305
- Nitrosoanilide complexes**
- metal complexes of some 2-nitrosoanilides, **223**, 71
- Nitrosoarene complexes**
- nitrosoarene and nitrosoalkane insertion reactions of titanacyclobutenes, **222**, 305
- Nitrosyl bending**
- synthesis, molecular structure and electrochemical investigation of the 1,2,5-thiadiazole-3,4-dithiolate-substituted complex Cp<sup>\*</sup>Ru(NO)(tdas). Evidence for nitrosyl bending during reduction, **226**, 267
- Nitrosyl complexes**
- synthesis and characterization of Keggin derivatives containing an [Mo(NO)]<sup>3+</sup> unit: (*n*-Bu<sub>4</sub>N)<sub>4</sub>[PM<sub>11</sub>O<sub>39</sub>{Mo(NO)}] ( $M = \text{Mo, W}$ ), **215**, 61
- transition metal complexes with sulfur ligands Part CII. Changes in configuration and reactivity caused by alkyl substituents R in [Fe(L)('N<sub>R</sub>S<sub>4</sub>')]<sup>+</sup> complexes (L = CO, NO<sup>+</sup>,

- NO,  $\text{PMe}_3$ ,  $\text{N}_2\text{H}_4$ ;  $\text{R} = \text{CH}_3$ ,  $\text{CH}_2\text{CH}_2\text{CO}_2\text{CH}_3$ ,  $\text{CH}_2\text{CH}_2\text{CO}_2^-$ ), **219**, 75
- activation of molecular oxygen by iron nitrosyls in the presence of the bidentate phosphines 1,2-bis(diphenylphosphino)ethane and ethene, **221**, 133
- transition metal complexes with sulfur ligands Part CVII. Complexes of the  $[\text{Mo}(\text{NO})(\text{S}_4)]$  fragment with 16, 17 and 18 valence electrons ( $\text{S}_4^{2-} = 1,2\text{-bis}(2\text{-mercaptophenylthio})\text{ethane}(2-)$ ), **225**, 163
- ligand substitution at 19-electron organometallic centers. Electrocatalytic CO substitution reactions of (methylcyclopentadienyl) $\text{Mn}(\text{CO})_2\text{NO}^+$  and (indenyl) $\text{Mn}(\text{CO})_2\text{NO}$ , **226**, 53
- quenching of a polypyridyl-ruthenium(II) chromophore by covalently attached  $\{\text{ML}(\text{NO})\text{Cl}\}$  fragments ( $\text{M} = \text{Mo}$ ,  $\text{W}$ ;  $\text{L} = \text{tris}(3,5\text{-dimethylpyrazolyl})\text{hydroborate}$ ), **226**, 171
- Nitroxide complexes**
- a seven-spin cluster formed by an alkyl nitronyl nitroxide biradical and copper(II): crystal structure and magnetic properties, **217**, 7
- NMR multinuclear**
- 1,1-organoboration of tri-1-alkynyltin compounds: novel triorganotin cations, stannoles, 3-stannolenes and 1-stanna-4-bora-2,5-cyclohexadienes, **220**, 161
- NMR spectroscopy**
- coordination study of D-xylo 5-hexulosonic acid with borate, molybdate and tungstate in aqueous solution by nuclear magnetic resonance spectroscopy, **221**, 69
- analysis of  $[(\text{dien})\text{Pd}]^{2+}$  binding to uracil and azauracils by proton NMR spectroscopy, **223**, 131
- effect of aromatic-aromatic interaction on ligand binding to zinc porphyrins, **224**, 157
- NMR study**
- azacryptand host for lead: NMR and structural studies, **218**, 5
- rotation and orientation of axially coordinated imidazoles in low-spin ferric porphyrin complexes, **224**, 113
- $^{15}\text{N}$  NMR**
- nitrogen-15 NMR coordination shifts of  $\beta$ -alanine and glycine in cobalt(III) complexes, **217**, 213
- $^{14}\text{N}$  NMR spectroscopy**
- phenylimido derivatives of  $[\text{Mo}_6\text{O}_{19}]^{2-}$ : syntheses, X-ray structures, vibrational, electrochemical,  $^{95}\text{Mo}$  and  $^{14}\text{N}$  NMR studies, **224**, 81
- Non-linear optics**
- synthesis of symmetric and unsymmetric 1,4-bis(*p*-R-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289
- Non-linear response effect**
- relationship between four-state delineation and two-state delineation in electron transfer reactions and excitation transfer, **225**, 185
- Non-stoichiometric compounds**
- electron transfer reactions in non-stoichiometric ceria and uranium, **225**, 305
- Nucleobase complexes**
- simultaneous binding of soft and hard metals to a pyrimidine nucleobase: *trans*- $\text{K}_2[\text{Pt}_2(1\text{-MeU})_2] \cdot 6\text{H}_2\text{O}$ , an anionic Pt-1-methyluracil (1-MeU) complex arranged in circles, **218**, 117
- simultaneous binding of soft and hard metals to a pyrimidine nucleobase: preparation and X-ray structure of  $[(\text{en})\text{Pd}(1\text{-MeT})_2\text{Na}_2](\text{NO}_3)_2 \cdot \text{H}_2\text{O}$  (1-MeT = 1-methylthymine anion), **227**, 5
- Nucleoside complexes**
- synthesis of water-soluble (tri(hydroxymethyl)phosphine)gold(I) complexes containing a nucleoside ligand, **217**, 201
- Nucleotide complexes**
- ternary Pt(II)-amino acid-nucleotide complexes: kinetics of formation, **218**, 11
- Octahedral complexes**
- structural characterization of octahedral metal complexes of  $[\text{MX}_3(\text{py})_3]$  type with three pyridines as revealed by  $^2\text{H}$  NMR spectra, **218**, 179
- the first isolated antineoplastic Ru(IV) complex: synthesis and structure of  $[\text{Cl}_2(1,2\text{-cyclohexanediaminotetraacetate})\text{Ru}] \cdot 2\text{H}_2\text{O}$ , **224**, 15
- Olefin complexes**
- three-coordinate Pt(0)  $\eta^2$ -complexes: electrophilic hydrogen attack through oxidative-addition of protic acids, **219**, 169
- photochemical reactions of transition metal organyl complexes with olefins Part 10. Light-induced formal [5+2] cycloadditions at manganese, **220**, 77
- stereochemical investigation of cationic olefin complexes of platinum(II) containing chiral diamines: a sequence of four asymmetric centres (N, C, C and N) in the chelate ring, **227**, 261
- Oligopyridine ligand complexes**
- a new functionalised oligopyridine ligand containing ferrocene as a ball-bearing spacer for metallosupramolecular chemistry, **224**, 11
- Optical electron transfer**
- solvent, electrolyte and solute shape effects on optical electron transfer in mixed-valence ruthenium ammine dimers, **226**, 137
- Optical spectroscopy**
- the influence of aluminum trichloride on a configuratively labile lactone-bridged biaryl: quantum chemical calculations and optical spectroscopy, **222**, 247
- Organoboration**
- 1,1-organoboration of tri-1-alkynyltin compounds: novel triorganotin cations, stannoles, 3-stannolenes and 1-stanna-4-bora-2,5-cyclohexadienes, **220**, 161
- Organometallic complexes**
- organometallic complexes of ruthenium(IV) with bidentate oxygen and nitrogen donor ligands, **218**, 81
- Organosilicon compounds**
- synthesis, properties and reactions of sila-functional  $\alpha$ -heteroatom-substituted organosilicon compounds, **220**, 129
- Organotin complexes**
- cyclic bis(amino)organotin cations, stabilized by  $\pi$ -coordination and new spirocyclic stannole derivatives, **216**, 51
- diorganotin(IV) derivatives of salicylaldehydethiosemicarbazone. The crystal structure of dimethyl- and diphenyl(salicylaldehydethiosemicarbazonato)tin(IV), **216**, 169
- synthesis and spectroscopic properties of diorganotin(IV) derivatives of 2,6-diacetylpyridine bis(thiosemicarbazone). Crystal structure of diphenyl{2,6-diacetylpyridine bis(thiosemicarbazonato)}tin(IV) bis(dimethylformamide) solvate, **221**, 61
- Organozinc reagents**
- influence of bulky substituents on the regioselective group-transfer reactions of diorganozinc compounds with *N,N'*-bis(2,6-diisopropylphenyl)-1,4-diaza-1,3-butadiene, **220**, 55
- Organyl complexes**
- photochemical reactions of transition metal organyl complexes with olefins Part 10. Light-induced formal [5+2] cycloadditions at manganese, **220**, 77
- Orotato complexes**
- synthesis, characterization and X-ray crystal structure of ethylenediamine(orotato)palladium(II), **221**, 173
- Orthometalation**
- orthometalation reactions in trifluoroacetate dirhodium(II) compounds. Molecular structure of

- $\text{Rh}_2(\text{O}_2\text{CCF}_3)_2[(\text{C}_6\text{H}_4)\text{PPh}_2]_2 \cdot (\text{PPh}_3)_2 \cdot 2(\text{C}_7\text{H}_8)$ , **218**, 189
- Orthophosphate complexes  
phosphate complexation of oxovanadium(IV). Evidence of bidentate chelation of orthophosphate, **215**, 225
- Osmium complexes  
synthesis and characterisation of cyanide bridged heterobinuclear complexes based on Group VI octacyanometalate(IV) and pentaammineosmium(III) ions: a comparison with Group VIII hexacyano analogues, **219**, 121  
regio- and stereoselective dimerization of 1-alkynes catalyzed by an Os(II) complex, **220**, 5  
preparation, properties and reactions of metal-containing heterocycles Part LXXXIX. Metalla[n]- and metalla[m,n]-orthocyclophanes as mediators for the synthesis of cyclic ketones, **220**, 107  
properties of the mixed-valence binuclear complex ion,  $[(\text{NH}_3)_5\text{Ru}^{\text{III}}-\text{NC}-\text{Os}^{\text{II}}(\text{CN})_5]$ , **223**, 37  
comparison of the rates of substitution in  $[\text{Ru}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  and  $[\text{Os}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  by hexacyano complexes: substitution coupled to electron transfer, **225**, 173  
inter-chromophore electronic interactions in ligand-bridged polynuclear complexes: a comparative study of various bridging ligands, **226**, 213  
substitution kinetics of some tetracarbonyl( $\eta^2$ -alkene)osmium complexes, **227**, 215
- Outer-sphere coordination  
outer-sphere coordination of polycyanometallate anions with polyammonium macrocycles: a spectrophotometric study, **227**, 71
- Oxalato complexes  
octahedral  $\mu$ -oxalato-nickel(II) dinuclear complexes with water and tridentate amines as blocking ligands: magnetostructural correlations, **216**, 139
- Oxamidate complexes  
crystal structure and magnetic property of a 2D infinite layered copper(II) complex with bridging *trans*-oxamidate and pyrimidine ligands, **224**, 171
- Oxazine complexes  
synthesis of 5,6-dihydro-4*H*-1,3-oxazines from neutral and cationic platinum(II) nitrile complexes. X-ray structure of *trans*- $[\text{Pt}(\text{CF}_3)\{\text{N}=\text{C}(\text{Ph})\text{OCH}_2\text{CH}_2\text{CH}_2\}(\text{PPh}_3)_2]\text{BF}_4$ , **220**, 21
- Oxidation  
optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220**, 283  
oxidation of *cis*- and *trans*-pinane with  $\text{RuO}_4$  generated in situ, **222**, 115  
the oxidation of  $[\text{Co}(\text{edta})]^{2-}$  by  $[\text{Co}(\text{phen})_3]$ , **225**, 75  
kinetics and mechanism of the ferrate oxidation of hydrazine and monomethylhydrazine, **225**, 145  
electron transfer reactivity of cuboidal heterometallic  $\text{Mo}_3\text{MS}_4$  clusters in aqueous acidic solutions, **225**, 157
- Oxidation potential  
synthesis, properties and reactions of sila-functional  $\alpha$ -heteroatom-substituted organosilicon compounds, **220**, 129
- Oxidation reaction  
ligand nature of coordinated  $\text{NO}_2^-$  in the oxidation reaction of Ru(II) complexes,  $[\text{Ru}(\text{NO}_2)(\text{OH}_2)(\text{py})_{4-2n}(\text{bpy})_n]^+$  ( $n=0, 1, 2$ ), and their related complexes, **225**, 111
- Oxidative quenching  
oxidative quenching of luminescence from copper metallothionein, **226**, 275
- Oxo complexes  
synthesis and characterization of Keggin derivatives containing an  $[\text{Mo}(\text{NO})]^{3+}$  unit:  $(n\text{-Bu}_4\text{N})_4[\text{PM}_{11}\text{O}_{39}\{\text{Mo}(\text{NO})\}]$  ( $M=\text{Mo}, \text{W}$ ), **215**, 61  
phosphate complexation of oxovanadium(IV). Evidence of bidentate chelation of orthophosphate, **215**, 225  
oxo- and sulfido-bridged sandwich cubane-type molybdenum-indium mixed-metal cluster. Synthesis and X-ray structure of  $[(\text{H}_2\text{O})_9\text{Mo}_3\text{S}_3\text{OInOS}_3\text{Mo}_3\text{-(H}_2\text{O)}_9]^{3-}(\text{CH}_3 \cdot \text{C}_6\text{H}_4 \cdot \text{SO}_3)_8 \cdot 30\text{H}_2\text{O}$ , **216**, 13  
binuclear oxo-bridged iron(III) complexes of *N,N'*-bis(2-pyridylmethyl)ethane-1,2-diamine (bispicen) and related ligands, **216**, 21  
synthesis, spectral properties and redox behaviour of oxovanadium(IV) complexes with tetraaza[14]annulenes, **216**, 217  
spectroscopic investigation of dichloro and hydrazido(2-) derivatives of bis(dithiocarbamato)dioxomolybdenum(VI) compounds. Crystal and molecular structure of bis(diisobutylthiocarbamato)(*N,N*-dimethylhydrazido(2-)-*N'*-oxomolybdenum(VI), **217**, 61  
systematics of some properties of hexamolybdotellurate(VI) salt hydrates with di- and trivalent metals,  $\text{M}_x[\text{TeMo}_6\text{O}_{24}] \cdot n\text{H}_2\text{O}$ , **217**, 93  
synthesis and characterization of symmetric and unsymmetric oxo-bridged trinuclear chromium benzoate complexes: crystal and molecular structure of  $[\text{Cr}_3\text{O}(\text{O}_2\text{CPh})_6(\text{py})_3]\text{ClO}_4$ , **217**, 171  
hydrothermal synthesis and crystal and molecular structure of a three-dimensional vanado-arsenate  $(\text{NH}_4)_3-[(\text{VO})(\text{V}_2\text{O}_3)(\text{AsO}_4)_2(\text{HAsO}_4)]$ , exhibiting both  $\{\text{V}-\text{O}-\text{V}\}_\infty$  chains and isolated  $\{\text{V}_2\text{O}_3\}$  units, **218**, 59  
synthesis and characterization of oxorhenium(V)-catecholate complexes. Crystal and molecular structures of  $(\text{CH}_3)_4\text{N}[\text{ReO}(\text{O}_2\text{C}_6\text{H}_4)_2]$  and  $(\text{CH}_3)_4\text{N}[\text{ReO}(\text{PPh}_3)(\text{O}_2\text{C}_6\text{H}_4)_2] \cdot \text{CH}_3\text{OH}$ , **218**, 157  
bis(2-mercaptopyridine-*N*-oxide)oxovanadium(IV): synthesis, structure and its conversion to the dichloro complex, **219**, 213  
molybdenum complexes with sterically-hindered thio-carboxylate ligands. Electrochemical properties of the anionic complex bis(2,2-diphenyl-2-mercaptoethanoate)dioxomolybdate(VI) in protic solvents, **221**, 117  
synthesis and X-ray structural determination of pentamethylcyclopentadienyl dichloro oxo molybdenum, **221**, 147  
oxidation of *cis*- and *trans*-pinane with  $\text{RuO}_4$  generated in situ, **222**, 115  
the reactions of thiolate ligands with rhenium hydrides Part 1. The syntheses and structures of  $[\text{ReO}(\text{SC}_6\text{H}_3\text{-2,6-Me}_2)_3(\text{PPh}_3)]$  and  $[\text{Re}(\text{SC}_6\text{H}_3\text{-2,6-Me}_2)_3(\text{PPh}_2\text{H})(\text{PPh}_3)]$ , **223**, 63  
thermodynamics of complexation of the uranyl(VI) ion with benzoate. The crystal structure of sodium tris(benzoate)dioxouranate dihydrate,  $\text{Na}[(\text{PhCO}_2)_3\text{UO}_2] \cdot 2\text{H}_2\text{O}$ , **223**, 151  
synthesis, characterization and crystal structures of two *cis*-dioxo vanadium(V) complexes of monoanionic tridentate Schiff base ligands, **223**, 193  
effect of substituting sulfur for oxygen on the heterogeneous electron transfer kinetics of oxomolybdenum(V) hydrotris(3,5-dimethyl-1-pyrazolyl)borate complexes, **225**, 1  
synthesis and electrochemical properties of oxo-bridged manganese dimers incorporating alkali and alkaline earth cations, **225**, 191  
electron transfer properties of  $\text{ReO}(\text{ECD})$ , **225**, 323  
crystal structure and solid-state fluorescence of an oxomolybdenum(V)-benzylviologen electron donor-acceptor salt, **226**, 25  
dynamical investigations of the catalytic mechanisms of water oxidation by the  $[(\text{bpy})_2\text{Ru}(\text{OH}_2)]_2\text{O}^{4+}$  ion, **226**, 179  
synthesis and characterization of neutral oxorhenium(V) and nitridotechnetium(V) complexes with a tetradentate  $\text{N}_2\text{S}_2$  unsaturated ligand derived from dithiocarboxylic acid, **227**, 25

- one-equivalent oxidative charge storage on an  $[\text{Mn}_2\text{O}_2]^{3+}$  core: synthesis, crystal structure and resonance Raman spectra of  $[\text{Mn}_2\text{O}_2(\text{pbz})_n]^{n+}$  ( $n=3, 4$ ), **227**, 33
- spectroscopic studies of the *trans*- $[\text{VOCl}_2(\text{H}_2\text{O})_2]$  chromophore in its hydrogen bonding adduct with 18-crown-6, **227**, 91
- Oxocyclophanes**
- preparation, properties and reactions of metal-containing heterocycles Part LXXXIX. Metalla[*n*]- and metalla[*m,n*]-orthocyclophanes as mediators for the synthesis of cyclic ketones, **220**, 107
- Oxomolybdenum complexes**
- redox reactions between oxomolybdenum(IV) and tetratolylporphinatoiron(III) complexes, **226**, 259
- Oxygenate compounds**
- rhodium-catalyzed synthesis of 2(5*H*)-furanones from terminal alkynes and non-substituted alkynes under water-gas shift reaction conditions, **220**, 45
- Oxygen carrier**
- a 'calix[4]arene' porphyrin as a new host and an oxygen carrier model, **224**, 1
- Palladium complexes**
- cyclometallation of *N,N*-dimethylbenzo[*b*]furan-2-carbothio (and seleno) amides with palladium(II), ruthenium(II) and rhodium(III), **215**, 91
- arylation of alkenes by the palladium(II) acetate–dialkyl sulfide system, **216**, 253
- synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129
- cyclopalladated *p*-azoxyanisole complexes. Synthesis and structure of  $\{[\text{bis}(2\text{-phenylpyridine})][4'-(\text{methoxyphenyl}-NNO\text{-azoxy}-N^2)-4\text{-methoxyphenyl-2-ato}]palladium(II)\}$  (tetrafluoroborate), **218**, 211
- spectroscopic studies on complex formation and non-covalent interactions in ternary palladium(II) systems involving spermidine and 2,3-diaminopropionate or 2,4-diaminobutyrate, **219**, 107
- asymmetric catalysis 87. Enantioselective allylation of 1,5-dimethylbarbituric acid with Pd catalysts and new optically active PN ligands, **220**, 63
- synthesis of  $\eta^3$ -allyl–Pd complexes and subsequent formation of *N*-heterocycles via the reaction of a cyclometallated compound with conjugated dienes, **220**, 115
- ultrasound-promoted palladium-catalyzed carbonyl allylation by allylic alcohols with tin(II) chloride in non-polar solvents, **220**, 155
- bite angle dependence of the rate of reductive elimination from diphosphine palladium complexes, **220**, 249
- synthesis of symmetric and unsymmetric 1,4-bis(*p*-*R*-phenylethynyl)benzenes via palladium/copper catalyzed cross-coupling and comments on the coupling of aryl halides with terminal alkynes, **220**, 289
- synthesis, characterization and X-ray crystal structure of ethylenediamine(oroato)palladium(II), **221**, 173
- palladium-assisted formation of carbon–carbon bonds Part 2. Stoichiometric synthesis of spirocyclic compounds. X-ray structure of a  $\pi$ -allylic palladium intermediate, **222**, 1
- formation of an allyl–tin bond in  $\eta^3$ -allyl(trichlorostannyl)palladium(II) complexes, **222**, 21
- strain, silyl and steric effects on the regioselectivity of palladium(0) catalyzed allyl esters reduction as alternative to the Wittig reaction, **222**, 37
- intermediates in the palladium-catalysed reaction of 1,3-dienes Part 8. The reaction of palladium–butadiene complexes with ethyl methylacetoacetate, **222**, 179
- synthesis and structure of new chiral ferrocenylphosphines for asymmetric catalysis, **222**, 213
- palladium complex catalysed synthesis of sulfinic acids, sulfinic acid esters, sulfonic acids and *S*-alkyl alkanethiosulfonates, **222**, 381
- analysis of  $[(\text{dien})\text{Pd}]^{2+}$  binding to uracil and azauracils by proton NMR spectroscopy, **223**, 131
- pyrazolyl carbon and proton NMR assignments for transition-metal poly(1-pyrazolyl)borates. Syntheses and NMR spectral studies of the ( $\eta^2$ - $\text{BR}_2\text{Pz}_2$ ) (Pz=1-pyrazolyl) palladium and platinum complexes, **224**, 131
- predicting the effect of oxidative doping on the conductivities of metal–tetrakis(dimethylphosphino)benzene coordination polymers from the electrochemical properties of their related bimetallic complexes, **225**, 15
- simultaneous binding of soft and hard metals to a pyrimidine nucleobase: preparation and X-ray structure of  $[(\text{en})\text{Pd}(1\text{-MeT})_2\text{Na}_2](\text{NO}_3)_2 \cdot \text{H}_2\text{O}$  (1-MeT=1-methylthymine anion), **227**, 5
- synthesis and characterization of a bidentate chelating diisonitrile ligand with two thioetheral sulfurs in a bridging group and its copper(I), rhodium(I) and palladium(II) complexes, **227**, 113
- Pb(II)Au(I) dicyanide complex**
- light-induced electron transfer in lead(II)gold(I) dicyanide, **226**, 345
- Peltier effect**
- thermometric determination of electrochemical Peltier heat (thermal effect associated with electron transfer) of some redox couples, **226**, 69
- Pentacoordinate complexes**
- the synthesis, structure and hydrolytic activity of a functional pentacoordinate zinc complex, **219**, 205
- Peptide complexes**
- bis(2-diphenylphosphinoethyl)phenylphosphineruthenium(II) complexes of amino acids and dipeptides, **217**, 51
- lithium coordination to amino acids and peptides. Synthesis, spectroscopic characterization and structure determination of lithium complexes of neutral and anionic glycine and diglycine, **218**, 121
- ternary complexes of *cis*- and *trans*- $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$  (*cis*-, *trans*-DDP) with 9-methylguanine (9-MeG), the dipeptides glycyglycine (glygly), glycyl-L-alanine (glyala), glycyl-L-2-aminobutyric acid (gly-2-aba), glycyl-L-norvaline (glynval), glycyl-L-norleucine (glynleu), and of *trans*- $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$  with Na-L-acetylhistidine, **227**, 17
- Peptide hydrolysis**
- hydrolytic cleavage of peptides by palladium(II) complexes is enhanced as coordination of peptide nitrogen to palladium(II) is suppressed, **217**, 21
- Perfluoroaryl complexes**
- synthesis of bis- or tris(pentafluorophenyl) thallium(III) complexes with bidentate ligands. X-ray crystal structure of  $[\text{Tl}(\text{C}_6\text{F}_5)_2(\text{acac})\{(\text{O})\text{Ph}_2\text{PCH}_2\text{PPh}_2(\text{O})\}]$ , **215**, 229
- Perhalocyclopentadienyl complexes**
- coordination chemistry of perhalogenated cyclopentadienes and alkynes XIV. Synthesis of dinuclear ring-bridged cyantrienes with  $-\text{SiMe}_2-$  or  $-\text{TiCp}_2-$  bridges. Structures of  $[(\text{OC})_3\text{Mn}(\text{C}_5\text{Br}_4-)]_2\text{SiMe}_2$  and  $[(\text{OC})_3\text{Mn}(\text{C}_5\text{Cl}_4-)]_2\text{TiCp}_2$ , **218**, 173
- Peroxo complexes**
- unusually stable peroxocopper complexes. Stoichiometry, products and kinetics of oxidation of the dimeric copper(I) complex  $[\text{LCuBr}]_2$  ( $L=N,N'$ -diethylethylenediamine) by dioxygen in methylene chloride from  $-51$  to  $30$  °C, **217**, 109

- synthesis and characterization of the new peroxo complex aquaoxodiperoxoimidazole molybdenum(VI), **223**, 173
- Peroxo sulfate complexes**  
the decomposition of peroxomonosulfate induced by manganese(II) in acid solution, **221**, 25
- Phenanthroline complexes**  
nickel(II) and manganese(II) complexes of substituted phenanthroline ligands, **221**, 99
- Phenolate complexes**  
synthesis, crystal and molecular structures, UV–Vis spectroscopy and electrochemical properties of two iron(III) phenolate complexes, **215**, 123
- Phenolic ligands**  
stability of metal ion complexes of *N,N'*-bis(2-hydroxybenzyl)ethylenediamine-*N,N'*-diacetic acid, **224**, 151
- Phosphaalkyne complexes**  
cyclodimerisation and addition reactions of Bu'C≡P at a cobalt(I) centre, **222**, 13
- Phosphate complexes**  
crystal and molecular structures of [*exo*- and *endo*-(RO)<sub>2</sub>P(O)-η<sup>5</sup>-C<sub>6</sub>H<sub>6</sub>]<sub>2</sub>Mn(CO)<sub>3</sub>, **224**, 147
- Phosphido complexes**  
electrochemical investigation of the electron-poor/precise (*n* = 5/6) complexes [Fe<sub>2</sub>(CO)<sub>*n*</sub>(μ-PR<sub>2</sub>)(μ-PR'<sub>2</sub>)] (*n* = 5, R = R' = Bu'; *n* = 6, R = R' = Ph; R = Bu', R' = Ph; R = Bu', R' = Cy). EPR study of the radical anion [Fe<sub>2</sub>(μ-PBu'<sub>2</sub>)<sub>2</sub>(CO)<sub>5</sub>], **217**, 29
- Phosphine complexes**  
phosphine and phosphine oxide adducts of vanadium(IV) chloride: synthesis and structural studies, **215**, 47  
reactions of phosphine–boron trihalide adducts with carbonyl-bis(triethylphosphine)rhodium(I) halides, **215**, 67  
Raman and infrared spectroscopic studies of tris triphenyl phosphine chloride complexes of transition metals, **216**, 191  
polyfunctional phosphine ligands II. Iridium(I) complexes of the 7-diphenylphosphino-2,4-dimethyl-1,8-naphthyridine (dpnapy) ligand. X-ray crystal structure of [Ir(cod)Cl]<sub>2</sub>(μ-dpnapy)], **216**, 209  
synthesis of water-soluble (tri(hydroxymethyl)phosphine)gold(I) complexes containing a nucleoside ligand, **217**, 201  
bromo derivatives of amine and phosphine complexes of cyanodihydroborane. Synthesis and reactivity, **218**, 21  
preparation and crystal structure of *anti*-(η<sup>3</sup>:η<sup>5</sup>-azulene)-pentakis(trimethylphosphine)dnicobalt. Reactions with carbon monoxide and solution dynamics, **218**, 143  
some new but-2-yne phosphine complexes of tungsten(II), **219**, 99  
gold in organic synthesis Part 2. Preparation of benzyl-alkyl and -arylketones via C–C coupling, **220**, 1  
the importance of phosphine-to-rhodium ratios in enantioselective hydroborations, **220**, 93  
a comparison of catalytic activity for imine hydrogenation using Ru ditertiary phosphine complexes, including chiral systems, **222**, 85  
transition metal complexes with sulfur ligands CVI. Phosphine and azido [Fe(L)(N<sub>3</sub>S<sub>4</sub>')] complexes; structural characterization of the key complex [Fe(N<sub>3</sub>S<sub>4</sub>')]·THF (L = PMe<sub>3</sub>, PBu<sub>3</sub>, PMe<sub>2</sub>Ph, PMePh<sub>2</sub>, N<sub>3</sub><sup>−</sup>; N<sub>3</sub>S<sub>4</sub>'<sup>2−</sup> = dianion of 2,2'-bis(2-mercaptophenylthio)diethylamine), **224**, 61
- Phosphinogallane complexes**  
trimethylgallium based phosphinogallanes. Synthesis and molecular structure of [Me<sub>2</sub>Ga–P(SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub>, **216**, 267
- Phosphite complexes**  
synthesis and X-ray structure of [acenaphthene–chromium(0)-dicarbonyl-tri(2,3,5-trimethylphenyl)phosphite], **219**, 209  
conversion of cationic phosphite complexes into phosphonate complexes with selective deamination on the coordinating phosphorus. Reaction of [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)L<sub>2</sub>Fe{P(NC<sub>4</sub>H<sub>8</sub>)<sub>*n*</sub>(OMe)<sub>3−*n*</sub>}]<sup>+</sup> (L = CO, PMe<sub>3</sub>) with KOH to afford (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)L<sub>2</sub>Fe-{P(O)(NC<sub>4</sub>H<sub>8</sub>)<sub>*n*−1</sub>(OMe)<sub>3−*n*</sub>}, **222**, 123
- Phosphonate complexes**  
conversion of cationic phosphite complexes into phosphonate complexes with selective deamination on the coordinating phosphorus. Reaction of [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)L<sub>2</sub>Fe{P(NC<sub>4</sub>H<sub>8</sub>)<sub>*n*</sub>(OMe)<sub>3−*n*</sub>}]<sup>+</sup> (L = CO, PMe<sub>3</sub>) with KOH to afford (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)L<sub>2</sub>Fe-{P(O)(NC<sub>4</sub>H<sub>8</sub>)<sub>*n*−1</sub>(OMe)<sub>3−*n*</sub>}, **222**, 123
- Phosphoryl complexes**  
synthesis, structure and stability of complexes of lanthanide nitrates with [(MeO)<sub>2</sub>P(O)]<sub>2</sub>C(OH)Ph and the isomeric (MeO)<sub>2</sub>P(O)OCHPhP(O)(OMe)<sub>2</sub>, **223**, 43
- Photochemical reaction**  
light-induced electron transfer in lead(II)gold(I) dicyanide, **226**, 345
- Photochemistry**  
photochemical reactions of transition metal organyl complexes with olefins Part 10. Light-induced formal [5 + 2] cycloadditions at manganese, **220**, 77  
reactivity of allene at phosphine-bridged di-iron centres: X-ray crystal structures of [Fe<sub>2</sub>(CO)<sub>5</sub>{μ-σ,η<sup>3</sup>-C(O)C(CH<sub>2</sub>)<sub>2</sub>}(μ-dppm)] and [Fe<sub>2</sub>(CO)<sub>4</sub>{μ-η<sup>3</sup>,η<sup>3</sup>-(CH<sub>2</sub>)<sub>2</sub>C<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>}(μ-dppm)]·Et<sub>2</sub>O, **220**, 201  
ruthenium chromophores containing terpyridine and a series of polyazine bridging ligands, **221**, 55  
photoredox chemistry of iodomercurate(II) complexes in acetonitrile, **221**, 79  
synthesis of α-hydroxy acid derivatives by the photochemical reaction of *O*-silylchromium(0) carbene complexes and nucleophiles, **222**, 261  
the synthesis, and UV–Vis, electrochemical, photochemical and intervalence characterization of tetraammine and tris chelated ruthenium(II) complexes bound to the bridging ligand dipyrido(2,3-*a*;2',3'-*h*)phenazine (dpop), **224**, 27  
cage escape yields for photoinduced bimolecular electron transfer reactions of Re(I) complexes, **225**, 41  
electrochemical, photophysical and photochemical properties of a hexanuclear ruthenium(II) bis(2,2'-bipyridyl) complex containing a tris(pyrazyltriazole)benzene ligand, **225**, 67  
bimolecular electron transfer at high driving forces. Photoreactions of gold(I) and iridium(I) complexes with pyridinium derivatives, **225**, 319  
photochemical and spectroscopic studies of complexes of iron(III) with citric acid and other carboxylic acids, **226**, 117  
spectroscopy and photoredox chemistry of ion-paired cobalt(III) am(m)ine complexes, **226**, 285
- Photoinduced electron transfer**  
photoinduced electron transfer reaction between [Cu(dmp)P<sub>2</sub>]<sup>+or−</sup> (P = PPh<sub>3</sub> or PPh<sub>2</sub>(*m*-C<sub>6</sub>H<sub>4</sub>SO<sub>3</sub><sup>−</sup>) and violon derivatives, **225**, 261  
electron transfer quenching of the excited MLCT state of Ru(II) in the double complex salts [Ru(bpy)<sub>3</sub>]<sub>2</sub>[M(CN)<sub>6</sub>]Cl·8H<sub>2</sub>O (M = Co and Fe), **226**, 203  
the construction of molecular systems capable of photoinduced multielectron charge transfer: a photochemical analysis of the reactivity of [(CN)<sub>5</sub>Fe<sup>II</sup>-(CN)-Pt<sup>IV</sup>(L)<sub>4</sub>-(NC)-Fe<sup>II</sup>(CN)<sub>5</sub>]<sup>4−</sup> (L = NH<sub>3</sub>, ethylenediamine), **226**, 251
- Photoinduced intramolecular electron transfer**  
photoinduced intramolecular electron transfer in bimetallic donor–acceptor assemblies, **226**, 79
- Photoluminescence studies**  
light-induced electron transfer in lead(II)gold(I) dicyanide, **226**, 345
- Photooxidation**

- simultaneous photooxidation of  $[\text{Ru}(\text{bpy})_2\text{Cl}_2]$  and photoreduction of  $[\text{Ru}(\text{bpy})_2\text{Cl}_2]^+$  in several solvent systems, **226**, 247
- Photophysics**
- electrochemical, photophysical and photochemical properties of a hexanuclear ruthenium(II) bis(2,2'-bipyridyl) complex containing a tris(pyrazyltriazole)benzene ligand, **225**, 67
- photophysical properties of mono-, di- and tetranuclear copper(I)-polypyridine complexes, **225**, 251
- Photoredox**
- spectroscopy and photoredox chemistry of ion-paired cobalt(III) am(m)ine complexes, **226**, 285
- Photoredox reactions**
- photoredox behavior of a trinuclear Ru(III),Ru(II),Co(III) complex induced by metal-to-metal charge transfer excitations, **225**, 327
- Photoreduction**
- photoinduced electron transfer reaction between  $[\text{Cu}(\text{dmp})\text{P}_2]^{+or-}$  ( $\text{P} = \text{PPh}_3$  or  $\text{PPh}_2(m\text{-C}_6\text{H}_4\text{SO}_3^-)$ ) and viologen derivatives, **225**, 261
- simultaneous photooxidation of  $[\text{Ru}(\text{bpy})_2\text{Cl}_2]$  and photoreduction of  $[\text{Ru}(\text{bpy})_2\text{Cl}_2]^+$  in several solvent systems, **226**, 247
- Phthalato complexes**
- antiferromagnetism tuned by the topology of the intervening bridging ligand. Crystal structure and magnetic properties of the  $(\mu\text{-phthalato})\{\{\text{bis}(\text{piperidine})\}\text{Cu}(\text{II})\}$  chain complex, **218**, 167
- Phthalocyanine complexes**
- redox reactions of an Mo(V) tetrasulfophthalocyanine, **215**, 27
- a comparison of the loop-current effect of silicon phthalocyanine and silicon naphthalocyanine rings on their axial ligands, **218**, 41
- equilibrium and kinetic study of piperidine binding to phthalocyaninatoiron(II) in dimethyl sulfoxide, **219**, 199
- tBuPcCrCl<sub>2</sub>H a sparingly soluble tetra-*t*-butyl-phthalocyaninato-chloro-chromium complex, **224**, 185
- electron transfer in surfactant films on electrodes; copper phthalocyaninetetrasulfonate-didodecyldimethylammonium bromide, **226**, 159
- Picrate complexes**
- synthesis and molecular structure of two crystalline forms of hydrated cerium(III) picrate, **221**, 169
- Piperidine binding**
- equilibrium and kinetic study of piperidine binding to phthalocyaninatoiron(II) in dimethyl sulfoxide, **219**, 199
- Piperidine complexes**
- antiferromagnetism tuned by the topology of the intervening bridging ligand. Crystal structure and magnetic properties of the  $(\mu\text{-phthalato})\{\{\text{bis}(\text{piperidine})\}\text{Cu}(\text{II})\}$  chain complex, **218**, 167
- Platinaborane complexes**
- ten-vertex polyhedral mixed diheteromonometalaborane chemistry. Directed synthesis of *arachno*-9,6,8-platinathiacarbodecaboranes, **218**, 1
- Platinum blue complexes**
- aggregation of platinum-uridine blue as a function of concentration in aqueous solution by small angle X-ray scattering, **216**, 37
- Platinum complexes**
- cyclometallated derivatives of platinum(II) derived from 1,4-benzodiazepin-2-ones. Crystal and molecular structure of  $\text{Pt}(\text{L})(\text{HL})\text{Cl}$  ( $\text{HL} = 7\text{-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin-2-one}$ , DIAZEPAM), a molecule containing a neutral and a deprotonated 1,4-benzodiazepin-2-one, **215**, 17
- metal complexes of benzodiazepines. Kinetic properties of bromazepam toward square planar dimethyl sulfoxide complexes of platinum(II) and palladium(II), **215**, 41
- synthesis and characterization of platinum(II)-thiolate complexes. Crystal and molecular structures of *cis*- $\text{Pt}(\text{Ph}_3\text{P})_2(\text{SC}_6\text{H}_2\text{-2,4,6-iPr}_3)\text{Cl}$  and *cis*- $\text{Pt}(\text{Ph}_3\text{P})_2(\text{SC}_6\text{H}_2\text{-2,4,6-iPr}_3)_2$ , **216**, 83
- molecular and crystal structures of *cis*- $\text{Pt}(\text{DMSO})(\text{C}_2\text{H}_5\text{CN})\text{Cl}_2$  and  $\text{Cl}(\text{DMSO})\text{Pt}(\mu\text{-N}(\text{CH}_3)\text{CHO})_2\text{Pt}(\text{DMSO})\text{Cl}$ , **216**, 163
- mercury and platinum mixed cluster having a prismatic structure,  $[\text{HgPt}_6(2,6\text{-Me}_2\text{C}_6\text{H}_3\text{NC})_{12}]$ , **217**, 121
- synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129
- molecular mechanical and quantum chemical study on the species involved in the hydrolysis of *cis*-diamminedichloroplatinum(II) and substituted bis(ethylenediamine)-dichloroplatinum(II) complexes Part I. Reactants and products, **217**, 159
- insertion reaction of dimethylacetylenedicarboxylate with the head-to-tail isomer of  $\text{Pt}_2\text{I}_2(\mu\text{-Ppy}_3)_2$ ; Ppy<sub>3</sub> = tris(2-pyridyl)phosphine, **217**, 209
- ternary Pt(II)-amino acid-nucleotide complexes: kinetics of formation, **218**, 11
- simultaneous binding of soft and hard metals to a pyrimidine nucleobase: *trans*- $\text{K}_2[\text{PtI}_2(1\text{-MeU})_2] \cdot 6\text{H}_2\text{O}$ , an anionic Pt-1-methyluracil (1-MeU) complex arranged in circles, **218**, 117
- preparation and structure of  $[\text{PhPt}(\text{OCO}_2\text{H})(\text{PEt}_3)_2]$  and  $[\text{PhPt}(\text{OH}_2)(\text{PEt}_3)_2]\text{BF}_4$ , **218**, 199
- new methods for synthesis of platinum(II) dimethyl sulfide complexes, **219**, 155
- three-coordinate Pt(0)  $\eta^2$ -complexes: electrophilic hydrogen attack through oxidative-addition of protic acids, **219**, 169
- synthesis and X-ray crystal structure of *cis*-1,4-diaminocyclohexanetetrachloroplatinum(IV): a new antitumor agent, **219**, 193
- synthesis of 5,6-dihydro-4H-1,3-oxazines from neutral and cationic platinum(II) nitrile complexes. X-ray structure of *trans*- $[\text{Pt}(\text{CF}_3)_2\{\text{N}=\text{C}(\text{Ph})\text{OCH}_2\text{CH}_2\text{CH}_2\}(\text{PPh}_3)_2]\text{BF}_4$ , **220**, 21
- non-isothermal spectrophotometric kinetics applied to inorganic reactions, **221**, 31
- binuclear complexes from 1,8-substituted anthracene derivatives. The crystal structure of  $\{\{\mu\text{-}[1,8\text{-anthracenedicarboxaldehydebis}(\text{dimethylhydrazone})]\}\text{tetrachlorobis}(\eta^2\text{-ethylene})\text{di-platinum}\}$ , **221**, 151
- trimetallic complexes of the tripod ligand 1,1,1-tris(diphenylphosphinomethyl)ethane Part I. Homotrimetallic *cis,cis,cis* and *cis,trans,cis* complexes of the type  $[\{\text{PtX}_2\text{-}\eta\text{-tripod}\}_2\text{PtX}_2]$  ( $\text{X} = \text{Cl, I and CH}_3$ ), **221**, 187
- alkyne hydration using Pt(II) catalysts, **222**, 317
- reactions of aziridine with platinum(II) nitriles. Formation of (aziridino)amidines and 2-imidazolines and X-ray structure of *trans*- $[\text{PtCl}_2\{\text{N}(\text{H})=\text{C}(\text{Ph})\text{NCH}_2\text{CH}_2\}_2]$ , **222**, 327
- chemistry of  $\pi$ -complexes derived from  $[(\text{Ph}_3\text{P})_2\text{Pt}(\eta^3\text{-CH}_2\text{CCPh})](\text{O}_3\text{SCF}_3)$ . Interconversion between  $\eta^3$ -alkoxoallyl and  $\eta^3$ -oxoallyl complexes and between  $\eta^3$ -aminoallyl and  $\eta^3$ -iminoallyl complexes, **223**, 1
- binuclear platinum(II) complexes of 1,1,4,7,10,10-hexaphenyl-1,4,7,10-tetraphosphadecane (P4) containing various bridging ligands as spacers between the P4 coordination units, **223**, 13
- pyrazolyl carbon and proton NMR assignments for transition-metal poly(1-pyrazolyl)borates. Syntheses and NMR spectral studies of the  $(\eta^2\text{-BR}_2\text{Pz}_2)$  ( $\text{Pz} = 1\text{-pyrazolyl}$ ) palladium and platinum complexes, **224**, 131
- predicting the effect of oxidative doping on the conductivities of metal-tetrakis(dimethylphosphino)benzene coordination polymers from the electrochemical properties of their related bimetallic complexes, **225**, 15



- electron transfer in *trans*-[Pt(PPh<sub>3</sub>)<sub>2</sub>(-C≡C-Fc)] and related compounds, **225**, 35
- the use of pressure-tuning spectroscopy to distinguish between one- and two-electron transfer processes, **225**, 89
- metal–metal interactions of aqueous lead(II) and tin(II) ions with the ground and excited states of tetrakis(μ-diphosphito)diplatinate(II), **225**, 201
- the construction of molecular systems capable of photoinduced multielectron charge transfer: a photochemical analysis of the reactivity of [(CN)<sub>5</sub>Fe<sup>II</sup>-(CN)-Pt<sup>IV</sup>(L)<sub>4</sub>-(NC)-Fe<sup>II</sup>(CN)<sub>5</sub>]<sup>4-</sup> (L = NH<sub>3</sub>, ethylenediamine), **226**, 251
- ternary complexes of *cis*- and *trans*-Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (*cis*-, *trans*-DDP) with 9-methylguanine (9-MeG), the dipeptides glycylglycine (glygly), glycyl-L-alanine (glyala), glycyl-L-2-aminobutyric acid (gly-2-aba), glycyl-L-norvaline (glynval), glycyl-L-norleucine (glynleu), and of *trans*-Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> with Na-L-acetylhistidine, **227**, 17
- organoplatinum(II) complexes containing disubstituted dithioamides: evidence for an S,S' Pt coordinated neutral dithioamide acting as an anion binding agent, **227**, 63
- thiopyridyl triazine derivatives and their platinum complexes: a new class of potential antitumor agents, **227**, 175
- stereochemical investigation of cationic olefin complexes of platinum(II) containing chiral diamines: a sequence of four asymmetric centres (N, C, C and N) in the chelate ring, **227**, 261
- <sup>31</sup>P NMR spectroscopy
- ionic cluster dithiophosphinates [Mo<sub>3</sub>S<sub>4</sub>(R<sub>2</sub>PS<sub>2</sub>)<sub>3</sub>py<sub>3</sub>]<sup>+</sup>[MI<sub>3</sub>]<sup>-</sup> (M = Zn·py, Cd·py, Hg), **216**, 177
- Poly(1-oxo-2-phenyltrimethylene)
- labeling studies of some carbonylation reactions of styrene with cationic palladium complexes, **220**, 273
- Polyacetylene
- ruthenium complex catalyzed polymerization of OH or COOH group containing alkynes to give functionalized poly(acetylene)s, **220**, 35
- Polyazacrown ligand complexes
- liquid matrix induced processes in fast atom bombardment mass spectrometry of polyazacrown transition metal complexes, **219**, 85
- Polychalcogenide complexes
- synthesis, structure and properties of the polychalcogenides [M<sub>4</sub>Te<sub>12</sub>]<sup>4-</sup> (M = Cd, Hg), **224**, 163
- Polycyanometallate complexes
- outer-sphere coordination of polycyanometallate anions with polyammonium macrocycles: a spectrophotometric study, **227**, 71
- Polydentate ligand complexes
- binuclear oxo-bridged iron(III) complexes of *N,N'*-bis(2-pyridylmethyl)ethane-1,2-diamine (bispicen) and related ligands, **216**, 21
- nucleophilic addition of water to coordinated di-(2-pyridyl)ketone (DPK) in rhenium(V) complexes. Synthesis and crystal structure of the adduct (DPK·H)<sup>+</sup>·[ReOCl<sub>2</sub>(DPK·OH)]<sub>2</sub>Cl, **217**, 141
- complexes of mercuric halides with the tripod ligand np<sub>3</sub> (= N(CH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>3</sub>). Molecular structure of [(np<sub>3</sub>)HgI]·0.5C<sub>2</sub>H<sub>5</sub>OH, **217**, 155
- stereoselectivity in reactions of metal complexes Part XVIII. Kinetics and stereoselectivity in electron-transfer reactions between [Co(L)H<sub>2</sub>O]<sup>+</sup> (L = *N,N'*-[(pyridine-2,6-diyl)bis(methylene)]bis[proline] (promp) and *N,N'*-[(4-methoxy-pyridine-2,6-diyl)bis(methylene)]bis[proline] (MeO-promp) and optically active iron(II) complexes, **218**, 73
- new heptacoordinated tin complexes of 2,6-diacetylpyridine-bis(thiosemicarbazone), H<sub>2</sub>daptsc, and of 2,6-diacetylpyridine-bis(semicarbazone), H<sub>2</sub>dapsc. Crystal and molecular structures of [MeSnCl(Hdaptsc)]Cl·MeOH and [MeSnCl(H<sub>2</sub>dapsc)]Cl<sub>2</sub>·2H<sub>2</sub>O, **218**, 139
- organometallic chelating ligands. Synthesis of a new anionic tripod like ligand with an N,N',O-donor set. X-ray crystal structure of [(η<sup>5</sup>-C<sub>6</sub>Me<sub>6</sub>)Ru{μ-P(O)(OMe)<sub>2</sub>}(μ-Pz)<sub>2</sub>Re(CO)<sub>3</sub>] (Me = methyl, Pz = pyrazolate), **219**, 67
- reactions of the polynuclear chain complex [Cu(L)(H<sub>2</sub>O)]<sub>n</sub> (H<sub>2</sub>L is 5-[(2'-carboxyphenyl)azo]-1,3-dimethylbarbituric acid) with imidazole. Crystal structures of the dinuclear and mononuclear derivatives and EPR study of the dimer–monomer equilibrium in solution, **219**, 127
- the synthesis, structure and hydrolytic activity of a functional pentacoordinate zinc complex, **219**, 205
- ruthenium chromophores containing terpyridine and a series of polyazine bridging ligands, **221**, 55
- trimetallic complexes of the tripod ligand 1,1,1-tris-(diphenylphosphinomethyl)ethane Part I. Homotrimetallic *cis,cis,cis* and *cis,trans,cis* complexes of the type [(PtX<sub>2</sub>-η-tripod)<sub>2</sub>PtX<sub>2</sub>] (X = Cl, I and CH<sub>3</sub>), **221**, 187
- novel didentate 2,2':6',2''-terpyridine complexes of ruthenium(II), **223**, 177
- electron transfer 122. Reductions of an N<sub>4</sub>-coordinated complex of silver(III), **225**, 83
- Polydentate phosphine ligand complexes
- binuclear platinum(II) complexes of 1,1,4,7,10,10-hexaphenyl-1,4,7,10-tetraphosphadecane (P4) containing various bridging ligands as spacers between the P4 coordination units, **223**, 13
- Polydentate sulfur ligand complexes
- transition metal complexes with sulfur ligands Part CVII. Complexes of the [Mo(NO)(S<sub>4</sub>')] fragment with 16, 17 and 18 valence electrons (S<sub>4</sub>'<sup>2-</sup> = 1,2-bis(2-mercaptophenylthio)ethane(2-)), **225**, 163
- Polyfunctional ligand complexes
- polyfunctional phosphine ligands II. Iridium(I) complexes of the 7-diphenylphosphino-2,4-dimethyl-1,8-naphthyridine (dpnapy) ligand. X-ray crystal structure of [Ir(cod)Cl]<sub>2</sub>(μ-dpnapy)], **216**, 209
- synthesis, properties and molecular structure of copper(II) and cobalt(II) 1,2-bis(pyrazol-1-ylmethyl)benzene complexes, **223**, 139
- Polyhaloaryl complexes
- polyhalophenylsilver(I) complexes, **223**, 161
- Polymer complexes
- coordination polymers of Mn(II) with the ligand pyrazine-2,3,5,6-tetracarboxylic acid, **219**, 161
- Polymeric compounds
- [Rh<sub>2</sub>(O<sub>2</sub>CCH<sub>3</sub>)<sub>4</sub>TCNE]<sub>x</sub> polymers: isomers with either 1,1- or *trans*-1,2-TCNE bridges, **221**, 1
- Polymerization
- ruthenium complex catalyzed polymerization of OH or COOH group containing alkynes to give functionalized poly(acetylene)s, **220**, 35
- Polymolybdate complexes
- synthesis and characterization of Keggin derivatives containing an [Mo(NO)]<sup>3+</sup> unit: (n-Bu<sub>4</sub>N)<sub>4</sub>[PM<sub>11</sub>O<sub>39</sub>{Mo(NO)}] (M = Mo, W), **215**, 61
- Polynuclear complexes
- reactions of the polynuclear chain complex [Cu(L)(H<sub>2</sub>O)]<sub>n</sub> (H<sub>2</sub>L is 5-[(2'-carboxyphenyl)azo]-1,3-dimethylbarbituric acid) with imidazole. Crystal structures of the dinuclear and mononuclear derivatives and EPR study of the dimer–monomer equilibrium in solution, **219**, 127
- photophysical properties of mono-, di- and tetranuclear copper(I)–polypyridine complexes, **225**, 251
- inter-chromophore electronic interactions in ligand-bridged polynuclear complexes: a comparative study of various bridging

- ing ligands, **226**, 213
- Polyoxomolybdate complexes  
phenylimido derivatives of  $[\text{Mo}_6\text{O}_{19}]^{2-}$ : syntheses, X-ray structures, vibrational, electrochemical,  $^{95}\text{Mo}$  and  $^{14}\text{N}$  NMR studies, **224**, 81
- Polypyrazolylborate complexes  
new polypyrazolylborate chemistry of ruthenium(II) complexes with weakly coordinating organonitrile ligands, **219**, 3
- Polypyridine complexes  
photophysical properties of mono-, di- and tetranuclear copper(I)–polypyridine complexes, **225**, 251  
intramolecular electron transfer between Ru(I) and Ru(III) and the heme iron of cytochrome *c* labeled with ruthenium(II) polypyridine complexes, **226**, 129
- Polypyridyl complexes  
quenching of a polypyridyl–ruthenium(II) chromophore by covalently attached  $\{\text{ML}(\text{NO})\text{Cl}\}$  fragments ( $\text{M}=\text{Mo}$ ,  $\text{W}$ ;  $\text{L}=\text{tris}(3,5\text{-dimethylpyrazolyl})\text{hydroborate}$ ), **226**, 171
- Porphindione complexes  
electrochemistry of nitrite reductase model compounds 5.  
Electrochemistry and spectroelectrochemistry of iron porphyrone, porphindione and isobacteriochlorin complexes, **225**, 57
- Porphinone complexes  
electrochemistry of nitrite reductase model compounds 5.  
Electrochemistry and spectroelectrochemistry of iron porphyrone, porphindione and isobacteriochlorin complexes, **225**, 57
- Porphyrin  
a ‘calix[4]arene’ porphyrin as a new host and an oxygen carrier model, **224**, 1
- Porphyrin complexes  
preparation and dioxygen binding property of a new zeolite encapsulated cobalt tetramethylporphyrin complex, **217**, 135  
synthesis, characterization and X-ray crystal structure of the benzenedithiolatohafnium(IV) porphyrin complex, **221**, 51  
rotation and orientation of axially coordinated imidazoles in low-spin ferric porphyrin complexes, **224**, 113  
effect of aromatic–aromatic interaction on ligand binding to zinc porphyrins, **224**, 157  
electrocatalysts for the four-electron reduction of dioxygen based on adsorbed cobalt tetrapyrrolylporphyrin molecules linked by aquaammine complexes of ruthenium(II), **225**, 215  
electron transfer reduction of cobalt tetraphenylporphyrin by hydroquinone dianions and alkylation with alkyl halides, **226**, 145  
reaction of low-valent iron porphyrins with alkyl containing supporting electrolytes, **226**, 195  
free radical-mediated electron transfer in organometallic complexes: homolysis and alkyl group crossover reactions for alkyliron(II) porphyrins, **226**, 231
- Potentiometric titration  
a potentiometric study on the complexation of Ag(I) with various N-methyl and C-methyl substituted ethylenediamines in 1 M  $\text{NO}_3^-$ , **224**, 97
- Pressure-tuning spectroscopy  
the use of pressure-tuning spectroscopy to distinguish between one- and two-electron transfer processes, **225**, 89
- Propynoato complexes  
synthesis, structure and solid state polymerization of di-aquabis(propynoato)magnesium, **222**, 339
- Protonation  
transition metal complexes with sulfur ligands CIV. Protonation and alkylation of thiolate donors of  $[\text{Fe}(\text{CO})_2(\text{S}_4)]$  and  $[\text{Fe}(\text{CO})(\text{S}_5)]$ : syntheses and properties of Fe(II) complexes with open-chain thioether ligands ( $\text{S}_4^{2-} = 1,2\text{-bis}(2\text{-mercaptophenylthio})\text{ethane}(2-)$ ;  $\text{S}_5^{2-} = 2,2'\text{-bis}(2\text{-mercaptophenylthio})\text{diethylsulfide}(2-)$ ), **224**, 45
- Proton transfer reactions  
direct solution calorimetric measurements of enthalpies of proton and electron transfer reactions for transition metal complexes. Thermochemical study of metal–hydride and metal–metal bond energies, **227**, 285
- Pseudopeptide complexes  
coordination abilities of amide oxygens in a 24-membered ring pseudopeptide toward several transition metal ions in acetonitrile, **223**, 121
- Pulse radiolysis  
reduction of hexaazacyclophane complexes of Co(II) and Cu(II) by  $\text{CH}_2\text{OH}$  and  $e_{\text{sol}}^-$ ; a pulse radiolysis study, **226**, 151
- Purine complexes  
dirhenium(III) complexes with *N,N*-dimethyladenine and 7-azaindole, **216**, 131  
synthesis and crystal structure of the  $\mu_2$ - $\{\text{tris}[\text{bis}(\text{diphenylphosphine})\text{methane}]\}-\mu_3\text{-3-methyl-8-ethylxanthine-}N,O,O\}$ - $\mu_3$ -chloro-triangle-tricopper(I)-water complex exhibiting a novel oxopurine coordination mode, **227**, 43
- Pyrazine complexes  
the electronic structure of pentaammine(pyrazine) ruthenium(II) and (III): the metal–ligand  $\pi$  conjugation and its implications in electron delocalization, **226**, 43
- Pyrazolylborate  
pyrazolyl carbon and proton NMR assignments for transition-metal poly(1-pyrazolyl)borates. Syntheses and NMR spectral studies of the  $(\eta^2\text{-BR}_2\text{Pz}_2)$  ( $\text{Pz} = 1\text{-pyrazolyl}$ ) palladium and platinum complexes, **224**, 131
- Pyridine complexes  
 $\text{tTBuPcCrCl}_2\text{H}$  a sparingly soluble tetra-*t*-butyl-phthalocyaninato-chloro-chromium complex, **224**, 185
- Pyridine derivative complexes  
crystal and molecular structure of the solvate of bis(2,6-bis(1'-methylbenzimidazol-2'-yl)pyridine)cobalt(II) dihexafluorophosphate with 4-methyl-1,3-dioxolan-2-on, **219**, 11
- Pyridinium derivatives  
bimolecular electron transfer at high driving forces. Photoreactions of gold(I) and iridium(I) complexes with pyridinium derivatives, **225**, 319
- Pyrimidine complexes  
crystal structure and magnetic property of a 2D infinite layered copper(II) complex with bridging *trans*-oxamidate and pyrimidine ligands, **224**, 171
- Quantum chemical calculations  
the influence of aluminum trichloride on a configuratively labile lactone-bridged biaryl: quantum chemical calculations and optical spectroscopy, **222**, 247
- Quantum mechanical calculations  
hemocyanin model compounds; a quantum mechanical calculation of geometry-dependent exchange interaction, **221**, 141
- Quantum mechanical study  
molecular mechanical and quantum chemical study on the species involved in the hydrolysis of *cis*-diamminedichloro-platinum(II) and substituted bis(ethylenediamine)dichloroplatinum(II) complexes Part I. Reactants and products, **217**, 159
- Quinone complexes  
the development of a quinone  $\alpha$ -anion synthon. Utilization of a maleoylcobalt complex as a quinone surrogate and a dominant role for ligand effects, **222**, 235
- Radical complexes

- a seven-spin cluster formed by an alkyl nitronyl nitroxide biradical and copper(II): crystal structure and magnetic properties, **217**, 7
- Raman spectroscopy  
 raman and infrared spectroscopic studies of tris triphenyl phosphine chloride complexes of transition metals, **216**, 191
- Reagent dependent  
 chemoselective functionalization of zirconacyclopentenes, **220**, 319
- Redox couples  
 thermometric determination of electrochemical Peltier heat (thermal effect associated with electron transfer) of some redox couples, **226**, 69
- Redox potentials  
 a spectroelectrochemical method for evaluating factors which regulate the redox potential of hemoglobins, **226**, 187
- Redox properties  
 synthesis, spectral and redox properties of a new series of aqua complexes of ruthenium(II), **221**, 125
- Redox reactions  
 photoredox chemistry of iodomercurate(II) complexes in acetonitrile, **221**, 79  
 transition metal complexes with sulfur ligands Part CVII. Complexes of the  $[\text{Mo}(\text{NO})(\text{S}_4)]$  fragment with 16, 17 and 18 valence electrons ( $\text{S}_4^{2-} = 1,2\text{-bis}(2\text{-mercaptophenylthio})\text{ethane}(2-)$ ), **225**, 163  
 redox reactions between oxomolybdenum(IV) and tetratolylporphyrinatoiron(III) complexes, **226**, 259  
 chemical and electrochemical approaches to the investigation of redox reactions of simple electron transfer metalloproteins, **226**, 293
- Reduction  
 the reduction of pentaammine(phenanthroline-2-carboxamido-*N*)cobalt(III) by titanium(III) and vanadium(II), **225**, 245  
 electron transfer reduction of cobalt tetraphenylporphyrin by hydroquinone dianions and alkylation with alkyl halides, **226**, 145  
 reduction of hexaazacyclophane complexes of Co(II) and Cu(II) by  $\text{CH}_2\text{OH}$  and  $e_{\text{sol}}^-$ ; a pulse radiolysis study, **226**, 151
- Reduction of carbon dioxide  
 electrocatalytic reduction of  $\text{CO}_2$  and  $\text{O}_2$  with electropolymerized films of vinyl-terpyridine complexes of Fe, Ni and Co, **225**, 285
- Reduction of oxygen  
 electrocatalytic reduction of  $\text{CO}_2$  and  $\text{O}_2$  with electropolymerized films of vinyl-terpyridine complexes of Fe, Ni and Co, **225**, 285
- Reductive cleavage  
 strain, silyl and steric effects on the regioselectivity of palladium(0) catalyzed allyl esters reduction as alternative to the Wittig reaction, **222**, 37
- Reductive coupling of iodobenzenes  
 use of aluminium as an electron pool. Reductive coupling of iodobenzenes promoted by electron transfer in an Ni/Pb/Al multi redox system, **222**, 323
- Reductive elimination  
 bite angle dependence of the rate of reductive elimination from diphosphine palladium complexes, **220**, 249
- Regioselection  
 ultrasound-promoted palladium-catalyzed carbonyl allylation by allylic alcohols with tin(II) chloride in non-polar solvents, **220**, 155
- Regioselective group transfer  
 influence of bulky substituents on the regioselective group-transfer reactions of diorganozinc compounds with *N,N'*-bis(2,6-di-isopropylphenyl)-1,4-diaza-1,3-butadiene, **220**, 55
- Regioselectivity  
 alkyne hydration using Pt(II) catalysts, **222**, 317
- Rhenium complexes  
 the unusual crystallographic disorder of  $\text{M}_2$  units in  $\text{M}_2\text{Cl}_4(\text{PEt}_3)_4$  ( $\text{M} = \text{Re}, \text{Mo}$ ), **215**, 103  
 mixed-ligand complexes of rhenium II. Synthesis and characterization of  $[\text{ReN}(\text{X})(\text{Me}_2\text{PhP})_2(\text{R}_2\text{tcb})]$  complexes ( $\text{X} = \text{I}, \text{N}_3, \text{SCN}, \text{CN}$ ;  $\text{R}_2\text{tcb}^- = N\text{-}(N,N\text{-dialkylthiocarbamoylbenzamidinate})$ ). X-ray crystal structure of  $[\text{ReN}(\text{N}_3)(\text{Me}_2\text{PhP})_2(\text{Et}_2\text{tcb})]$ , **215**, 159  
 mixed-ligand complexes of rhenium III. Synthesis, characterization and structures of  $[\text{ReN}(\text{Me}_2\text{PhP})_2(\text{mnt})]$  and  $[\text{ReN}(\text{Me}_2\text{PhP})(2,6\text{-DMTP})(\text{morph tcb})]$  ( $\text{Me}_2\text{PhP} = \text{dimethylphenylphosphine}$ ;  $2,6\text{-DMTP}^- = 2,6\text{-dimethylthiophenolate}$ ;  $\text{morph tcb}^- = N\text{-}(morpholinylthiocarbonyl)benzamidinate$ ), **216**, 31  
 dirhenium(III) complexes with *N,N*-dimethyladenine and 7-azaindole, **216**, 131  
 isothiocyanato complexes of rhenium II. Synthesis, characterization and structures of  $\text{ReN}(\text{NCS})_2(\text{Me}_2\text{PhP})_3$  and  $\text{Re}(\text{NS})(\text{NCS})_2(\text{Me}_2\text{PhP})_3$ , **216**, 223  
 a mixed-metal salt comprised of metal-metal bonded dinuclear ions: structure and properties of  $[\text{Rh}_2(\text{OAc})_2(\text{MeCN})_6]_n[\text{Re}_2\text{Cl}_8]$ , **217**, 79  
 nucleophilic addition of water to coordinated di-(2-pyridyl)ketone (DPK) in rhenium(V) complexes. Synthesis and crystal structure of the adduct  $(\text{DPK}\cdot\text{H})^+ \cdot \{\{\text{ReOCl}_2(\text{DPK}\cdot\text{OH})_2\text{Cl}\}\}_n$ , **217**, 141  
 synthesis and characterization of oxorhenium(V)-catecholate complexes. Crystal and molecular structures of  $(\text{CH}_3)_4\text{N}[\text{ReO}(\text{O}_2\text{C}_6\text{H}_4)_2]$  and  $(\text{CH}_3)_4\text{N}[\text{ReO}(\text{PPh}_3)(\text{O}_2\text{C}_6\text{H}_4)_2] \cdot \text{CH}_3\text{OH}$ , **218**, 157  
 exchange of azine ligands in *fac*- $\text{Re}(\text{CO})_3(4\text{-cyanopyridine})_2\text{Cl}$  and *fac*- $\text{Re}(\text{CO})_3(\text{quinoline})_2\text{Cl}$ , **221**, 161  
 EXAFS studies of a chiral rhenium lactone complex and its precursors including multiple scattering calculations, **222**, 27  
 insertion of NH into the rhenium-carbon bonds of a Fischer-type rhenacyclobutadiene. Preparation and characterization of azarhenacyclic complexes, **222**, 199  
 the reactions of thiolate ligands with rhenium hydrides Part 1. The syntheses and structures of  $[\text{ReO}(\text{SC}_6\text{H}_3\text{-}2,6\text{-Me}_2)_3(\text{PPh}_3)]$  and  $[\text{Re}(\text{SC}_6\text{H}_3\text{-}2,6\text{-Me}_2)_3(\text{PPh}_2\text{H})(\text{PPh}_3)]$ , **223**, 63  
 isothiocyanato complexes of rhenium III. Synthesis and characterization of  $[\text{ReN}(\text{NCS})_2(\text{Ph}_3\text{P})_2]$  and  $[\text{ReN}(\text{NCS})_2(\text{CH}_3\text{CN})(\text{Ph}_3\text{P})_2]$ . X-ray crystal structure of  $[\text{ReN}(\text{NCS})_2(\text{CH}_3\text{CN})(\text{Ph}_3\text{P})_2] \cdot \text{CH}_3\text{CN}$ , **224**, 193  
 cage escape yields for photoinduced bimolecular electron transfer reactions of Re(I) complexes, **225**, 41  
 electron transfer properties of  $\text{ReO}(\text{ECD})$ , **225**, 323  
 electron-transfer activation of the aminocarbene and the hydrogen isocyanide complexes *trans*- $[\text{ReCl}(\text{CNH}_n)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2][\text{BF}_4]_{n-1}$  ( $n = 2$  or  $1$ ). Interconversion of coordinated  $\text{CNH}_2$  and  $\text{CNH}$ , **226**, 9  
 photoinduced intramolecular electron transfer in bimetallic donor-acceptor assemblies, **226**, 79  
 synthesis and characterization of neutral oxorhenium(V) and nitridotechnetium(V) complexes with a tetradentate  $\text{N}_2\text{S}_2$  unsaturated ligand derived from dithiocarboxylic acid, **227**, 25
- Rhodium  
 solution structures of cobalt-rhodium clusters: the identification of a new hexanuclear cobalt-rhodium species, **227**, 187
- Rhodium carboxamidate complexes  
 dirhodium(II) tetrakis[*N,N*-dimethyl-2-pyrrolidone-5(*S*)-carboxamide]. Structural effects on enantioselection in metal car-

- bene transformations, **220**, 193
- Rhodium complexes
- preparation and structural characterization of  $\text{Rh}_2(\text{O}_2\text{CCPh}_3)_4(\text{EtOH})_2$ ,  $\text{Ru}_2(\text{O}_2\text{CCPh}_3)_4 \cdot (\text{H}_2\text{O})(\text{EtOH}) \cdot 2\text{EtOH}$  and  $\text{Mo}_2(\text{O}_2\text{CCPh}_3)_4 \cdot 3\text{CH}_2\text{Cl}_2$ , **215**, 9
- reactions of phosphine–boron trihalide adducts with carbonyl-bis(triethylphosphine)rhodium(I) halides, **215**, 67
- cyclometallation of *N,N*-dimethylbenzo[*b*]furan-2-carbothio (and seleno) amides with palladium(II), ruthenium(II) and rhodium(III), **215**, 91
- evidence of the interaction between steric and electronic influence in rhodoximes and cobaloximes. Synthesis of  $\text{pyRh}(\text{DH})_2\text{I}$  and X-ray structure of  $\text{pyRh}(\text{DH})_2\text{Cl}$ ,  $\text{pyCo}(\text{DH})_2\text{Cl}$  and  $\text{pyRh}(\text{DH})_2\text{I}$ , **216**, 125
- pressure and concentration dependent formation of oligomers of tetrakis(*p*-methylphenylisocyanide)rhodium(I), **216**, 245
- a mixed-metal salt comprised of metal–metal bonded dinuclear ions: structure and properties of  $[\text{Rh}_2(\text{OAc})_2(\text{MeCN})_6][\text{Re}_2\text{Cl}_8]$ , **217**, 79
- a mechanistic study of the homogeneous catalytic hydroformylation of formaldehyde: synthesis and characterization of model intermediates, **218**, 89
- orthometallation reactions in trifluoroacetate dirhodium(II) compounds. Molecular structure of  $\text{Rh}_2(\text{O}_2\text{CCF}_3)_2[(\text{C}_6\text{H}_4)\text{PPh}_2]_2 \cdot (\text{PPh}_3)_2 \cdot 2(\text{C}_7\text{H}_8)$ , **218**, 189
- rhodium-catalyzed synthesis of 2(*5H*)-furanones from terminal alkynes and non-substituted alkynes under water-gas shift reaction conditions, **220**, 45
- the importance of phosphine-to-rhodium ratios in enantioselective hydroborations, **220**, 93
- hydroformylation of alkenes and alkynes using a heterobinuclear Rh–W catalyst, **220**, 187
- synthesis and structural characterization of a new series of ruthenium(II) complexes containing the short bite  $\text{Ph}_2\text{PPy}$  ligand. Cooperative effect between the anionic rhodium and cationic ruthenium species in the catalytic hydroformylation of styrene by  $[\text{Ru}(\text{Ph}_2\text{PPy})_3\text{Cl}][\text{Rh}(\text{CO})_2\text{Cl}_2]$ , **221**, 109
- reaction of primary silanes with dinuclear rhodium hydride complexes: silane coupling reactions, **222**, 345
- synthesis and X-ray analysis of the dirhodium(II,II) carboxylate complex  $\text{Rh}_2(\text{O}_2\text{CCH}_3)_4(\text{formH})_2$  (formH = *N,N'*-di-*p*-tolylformamidine), **223**, 145
- infrared spectroelectrochemistry of  $[\text{Rh}(\text{TM}_4)_4\text{M}(\text{CO})_5]^{2+}$  (TM4 = 2,5-diisocyano-2,5-dimethylhexane; M = Re, Mn): electron-transfer-promoted metal–metal bond cleavage and formation, **226**, 61
- synthesis and characterization of a bidentate chelating diisonitrile ligand with two thioetheral sulfurs in a bridging group and its copper(I), rhodium(I) and palladium(II) complexes, **227**, 113
- Rhodium compounds
- $[\text{Rh}_2(\text{O}_2\text{CCH}_3)_4\text{TCNE}]_x$  polymers: isomers with either 1,1- or *trans*-1,2-TCNE bridges, **221**, 1
- Rhodoxime complexes
- evidence of the interaction between steric and electronic influence in rhodoximes and cobaloximes. Synthesis of  $\text{pyRh}(\text{DH})_2\text{I}$  and X-ray structure of  $\text{pyRh}(\text{DH})_2\text{Cl}$ ,  $\text{pyCo}(\text{DH})_2\text{Cl}$  and  $\text{pyRh}(\text{DH})_2\text{I}$ , **216**, 125
- Ring opening
- reactions of benzocyclobutene chromium complexes with carbon, nitrogen and oxygen nucleophiles: nucleophilic addition and unexpected proximal ring opening reactions, **220**, 175
- Ruthenium amine complexes
- a spectroelectrochemical method for evaluating factors which regulate the redox potential of hemoglobins, **226**, 187
- Ruthenium complexes
- cyclometallation of *N,N*-dimethylbenzo[*b*]furan-2-carbothio (and seleno) amides with palladium(II), ruthenium(II) and rhodium(III), **215**, 91
- synthesis and fluxional behaviour of (cyclopentadienyl) and (pentamethylcyclopentadienyl)ruthenium(II) complexes with hemilabile ether-phosphines.  $^{31}\text{P}$  DNMR spectroscopic studies and line shape analysis, **215**, 165
- pyrazine-mediated antiferromagnetic intermolecular exchange in mixed-valent diruthenium tetracarboxylates, **215**, 203
- bis(2-diphenylphosphinoethyl)phenylphosphineruthenium(II) complexes of amino acids and dipeptides, **217**, 51
- solid-state isomerization of ruthenium(II) isocyanide complexes, **217**, 181
- synthesis and characterization of phosphane-substituted hydroxyruthenocenes: crystal structure of  $[\text{Ru}(\eta^5\text{-C}_5\text{H}_4\text{PPh}_3)(\eta^5\text{-C}_5\text{H}_4\text{OH})]\text{PF}_6$ , **217**, 203
- electrochemistry of organometallic halide complexes III. Oxidation of  $[\text{C}_6\text{Me}_6\text{RuCl}_2]_2$ , **218**, 33
- organometallic complexes of ruthenium(IV) with bidentate oxygen and nitrogen donor ligands, **218**, 81
- new polypyrazolylborate chemistry of ruthenium(II) complexes with weakly coordinating organonitrile ligands, **219**, 3
- dimolybdenum(II) and diruthenium(II) complexes with 2-substituted 7-methyl-1,8-naphthyridines as bridging ligands, **219**, 23
- organometallic chelating ligands. Synthesis of a new anionic tripod like ligand with an *N,N'*O-donor set. X-ray crystal structure of  $[(\eta^6\text{-C}_6\text{Me}_6)\text{Ru}\{\mu\text{-P}(\text{O})(\text{OMe})_2\}(\mu\text{-Pz})_2\text{Re}(\text{CO})_3]$  (Me = methyl, Pz = pyrazolate), **219**, 67
- ruthenium complex catalyzed polymerization of OH or COOH group containing alkynes to give functionalized poly(acetylene)s, **220**, 35
- preparation, properties and reactions of metal-containing heterocycles Part LXXXIX. Metalla[*n*]- and metalla[*m,n*]-orthocyclophanes as mediators for the synthesis of cyclic ketones, **220**, 107
- direct nitration of aryl groups  $\sigma$ -bound to ruthenium(II), **220**, 261
- $\pi$ -facial selectivities in the coordination of an optically pure fused cyclopentadienyl ligand to arene ruthenium moieties. NMR spectral and structural characterization of a single diastereomer of  $[(\text{PCp})\text{Ru}(\text{Me}_6\text{C}_6)]\text{PF}_6$ , **221**, 41
- ruthenium chromophores containing terpyridine and a series of polyazine bridging ligands, **221**, 55
- synthesis and structural characterization of a new series of ruthenium(II) complexes containing the short bite  $\text{Ph}_2\text{PPy}$  ligand. Cooperative effect between the anionic rhodium and cationic ruthenium species in the catalytic hydroformylation of styrene by  $[\text{Ru}(\text{Ph}_2\text{PPy})_3\text{Cl}][\text{Rh}(\text{CO})_2\text{Cl}_2]$ , **221**, 109
- synthesis, spectral and redox properties of a new series of aqua complexes of ruthenium(II), **221**, 125
- the reaction of  $\text{CpRu}(\text{PPh}_3)_2^+$  with trimethylenesulfide. Synthesis and molecular and crystal structure of  $[\text{CpRu}(\text{PPh}_3)_2(\text{SC}_3\text{H}_6)]\text{CF}_3\text{SO}_3$ , **221**, 157
- a comparison of catalytic activity for imine hydrogenation using Ru ditertiary phosphine complexes, including chiral systems, **222**, 85
- oxidation of *cis*- and *trans*-pinane with  $\text{RuO}_4$  generated in situ, **222**, 115
- selective transformations of alkynols catalyzed by ruthenium complexes, **222**, 155
- hydration of conjugated dienes to produce ketones catalyzed by ruthenium complexes, **222**, 225
- arene migration in ruthenium clusters: a kinetic study of the isomerisation of  $\text{Ru}_5\text{C}(\text{CO})_{12}(\mu_3\text{-}\eta^2\text{-}\eta^2\text{-C}_6\text{H}_6)$  to  $\text{Ru}_5\text{C}(\text{CO})_{12}(\eta^6\text{-C}_6\text{H}_6)$ , **222**, 299

- properties of the mixed-valence binuclear complex ion,  $[(\text{NH}_3)_5\text{Ru}^{\text{III}}-\text{NC}-\text{Os}^{\text{II}}(\text{CN})_5]$ , **223**, 37
- an improved synthesis of Ru(BINAP) type complexes, **223**, 165
- novel didentate 2,2':6',2''-terpyridine complexes of ruthenium(II), **223**, 177
- the first isolated antineoplastic Ru(IV) complex: synthesis and structure of  $[\text{Cl}_2(1,2\text{-cyclohexanediaminotetraacetate})\text{Ru}]\cdot 2\text{H}_2\text{O}$ , **224**, 15
- the synthesis, and UV–Vis, electrochemical, photochemical and intervalence characterization of tetraammine and tris chelated ruthenium(II) complexes bound to the bridging ligand dipyrrodo(2,3-*a*;2',3'-*h*)phenazine (dpop), **224**, 27
- electrochemical behavior and X-ray crystal structure of the unusual dimeric ruthenium(cyclooctadiene) complex, (tri- $\mu$ -chloro)(chloro)(acetonitrile)bis( $\eta^4$ -cyclooctadiene)-diruthenium(II), **224**, 125
- charge type effects on the temperature dependence of intervalence transfer in ferrocyanide substituted ruthenium(III) amine and aminopolycarboxylate compounds, **224**, 189
- electrochemical, photophysical and photochemical properties of a hexanuclear ruthenium(II) bis(2,2'-bipyridyl) complex containing a tris(pyrazyltriazole)benzene ligand, **225**, 67
- mixed valence state associated with halogen migration in diruthenocenylmethane and its related compounds, **225**, 103
- ligand nature of coordinated  $\text{NO}_2^-$  in the oxidation reaction of Ru(II) complexes,  $[\text{Ru}(\text{NO}_2)(\text{OH}_2)(\text{py})_{4-2n}(\text{bpy})_n]^+$  ( $n=0, 1, 2$ ), and their related complexes, **225**, 111
- solvent trapped valency in the [bis(pentaammineruthenium)-( $\mu$ -1,4-dicyanamidobenzene)] trication, **225**, 123
- intramolecular electron transfer from  $\text{Ru}^{\text{II}}(\text{EDTA})^{2-}$  to  $\text{Co}^{\text{III}}(\text{NH}_3)_5^{3+}$  via nitrogen heterocycle bridging ligands, **225**, 129
- comparison of the rates of substitution in  $[\text{Ru}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  and  $[\text{Os}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}$  by hexacyano complexes: substitution coupled to electron transfer, **225**, 173
- electrocatalysts for the four-electron reduction of dioxygen based on adsorbed cobalt tetrapyrrolylporphyrin molecules linked by aquaammine complexes of ruthenium(II), **225**, 215
- light induced exchange energy transfer reactions of micelle forming Ru(II) diimine complexes, **225**, 275
- photoredox behavior of a trinuclear Ru(III), Ru(II), Co(III) complex induced by metal-to-metal charge transfer excitations, **225**, 327
- synthesis, electrochemistry and photoexcited-state properties of dinuclear ruthenium complexes bridged by 2,6'-bis(2-pyridyl)-2,2':6,2''-thiazolo[4,5-*d*]-benzothiazole, **226**, 17
- the electronic structure of pentaammine(pyrazine) ruthenium(II) and (III): the metal–ligand  $\pi$  conjugation and its implications in electron delocalization, **226**, 43
- photoinduced intramolecular electron transfer in bimetallic donor–acceptor assemblies, **226**, 79
- forbidden  $\pi$ - $\sigma$  donor–acceptor electronic coupling in linked transition metal complexes: contrasting behavior of ruthenium(II) donors with ruthenium(III) and cobalt(III) acceptors, **226**, 109
- intramolecular electron transfer between Ru(I) and Ru(III) and the heme iron of cytochrome *c* labeled with ruthenium(II) polypyridine complexes, **226**, 129
- solvent, electrolyte and solute shape effects on optical electron transfer in mixed-valence ruthenium ammine dimers, **226**, 137
- quenching of a polypyridyl–ruthenium(II) chromophore by covalently attached  $\{\text{ML}(\text{NO})\text{Cl}\}$  fragments ( $\text{M}=\text{Mo}, \text{W}$ ;  $\text{L}=\text{tris}(3,5\text{-dimethylpyrazolyl})\text{hydroborate}$ ), **226**, 171
- dynamical investigations of the catalytic mechanisms of water oxidation by the  $[(\text{bpy})_2\text{Ru}(\text{OH}_2)]_2\text{O}^{4+}$  ion, **226**, 179
- electron transfer quenching of the excited MLCT state of Ru(II) in the double complex salts  $[\text{Ru}(\text{bpy})_3]_2[\text{M}(\text{CN})_6]\text{Cl}\cdot 8\text{H}_2\text{O}$  ( $\text{M}=\text{Co}$  and  $\text{Fe}$ ), **226**, 203
- inter-chromophore electronic interactions in ligand-bridged polynuclear complexes: a comparative study of various bridging ligands, **226**, 213
- simultaneous photooxidation of  $[\text{Ru}(\text{bpy})_2\text{Cl}_2]$  and photoreduction of  $[\text{Ru}(\text{bpy})_2\text{Cl}_2]^+$  in several solvent systems, **226**, 247
- syntheses of ruthenium–iridium heterobimetallic complexes using tridentate phosphine ligands, **227**, 79
- the role of  $\text{H}_2\text{Ru}(\text{CO})_2(\text{PBu}_3)_2$  in the activation of alkynes and alkenes, **227**, 247
- Ruthenocenylmethane complexes
- mixed valence state associated with halogen migration in diruthenocenylmethane and its related compounds, **225**, 103
- Salt complexes
- the preparation, properties and crystal structure of a salt containing the hexabromochromate(III) anion, **227**, 167
- Schiff base complexes
- complexes of tetradentate  $\beta$ -keto enamine Schiff bases: an electrospray mass spectrometric study, **216**, 75
- preparation, properties and coordination behaviour of planar or tridimensional compartmental Schiff bases, **219**, 135
- direct electrochemical synthesis of tin(IV) complexes of 2-(2-mercaptoethyl)iminomethyl-phenols; the molecular structure of bis(2-(2-mercaptoethyl)iminomethyl-5-bromo-phenoxy)-tin(IV), **223**, 87
- synthesis, characterization and crystal structures of two *cis*-dioxo vanadium(V) complexes of monoanionic tridentate Schiff base ligands, **223**, 193
- structures of a monomeric and a dimeric Ni(II) complex containing 2-acetylpyrrole, **227**, 159
- Schiff base ligand complexes
- electrochemical and structural studies of nickel(II) complexes with  $\text{N}_2\text{O}_2$  Schiff base ligands 2. Crystal and molecular structure of *N,N'*-1,2-ethane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II), *N,N'*-1,2-*cis*-cyclohexane-1,2-diyl-bis(2-hydroxyacetophenonylideneimine)nickel(II) and *N,N'*-1,2-benzene-1,2-diyl-bis(3,5-dichlorosalicylideneimine)nickel(II), **219**, 43
- Selenido complexes
- synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129
- Semicarbazone complexes
- synthesis and characterization of lead(II)–semicarbazone complexes. Crystal structures of dinitrato di-2-pyridylketone semicarbazone lead(II) and dichloro cyclohexane-1,2-dione bis(semicarbazone) dimethylformamide lead(II), **215**, 85
- cobalt(III) complexes with 2-acetyl- and 2-formylpyridine *4N*-methylthiosemicarbazones, **216**, 229
- Semiquinone complexes
- valence tautomerism for catechol/semiquinone complexes of the *trans*-M(Bupy)<sub>2</sub>(3,6-DBQ)<sub>2</sub> ( $\text{M}=\text{Mn}, \text{Fe}, \text{Co}$ ) series, **226**, 91
- Silane complexes
- reactions of  $\text{HCCO}_3(\text{CO})_9$  with silanes; synthesis and electrochemistry of  $\text{X}[\text{SiMe}_2\text{CCO}_3(\text{CO})_9]_2$  ( $\text{X}=\text{O}, 1,4\text{-C}_6\text{H}_4$ ), **224**, 73
- Silane coupling reactions
- reaction of primary silanes with dinuclear rhodium hydride complexes: silane coupling reactions, **222**, 345
- Silicon complexes

- two silicon-bridged cyclopentadienyl  $\text{Fe}_2\text{S}_4$  cluster units connected through an  $\text{Fe(III)S}_6$  unit. Crystal and molecular structure of  $[\{(\text{CH}_3)_2\text{SiCp}^*_2\}_2\text{Fe}_5(\mu_3\text{-S}_2)_2(\mu_4\text{-S}_2)(\mu_4\text{-S}_4)]\text{FeCl}_4$ , **216**, 1
- a comparison of the loop-current effect of silicon phthalocyanine and silicon naphthalocyanine rings on their axial ligands, **218**, 41
- Silicon compounds
- electrophilic addition to styrylsilanes: sequential carbon-carbon bond forming reactions, **220**, 145
- Silver complexes
- the crystal structure of a chelate silver(I) complex:  $\{\text{Ag}[\text{Ph}_2\text{P}(\text{CH}_2)_2\text{SCH}_2\text{CH}_3]_2\}\text{ClO}_4$ , **215**, 199
- a novel tetranuclear iodoargentate(I) ion,  $[\text{Ag}_4\text{I}_8]^{4-}$ , with a cubane-type core, **217**, 15
- solution studies of monovalent metal complex ions of the bibrachial lariat ether 7,16-bis(2-methoxyethyl)-1,4,10,13-tetraoxa-7,16-diazacyclooctadecane, **219**, 217
- $\pi$ -bonded alkene and arene complexes of silver(I): an electrospray mass spectrometric study, **220**, 99
- polyhalophenylsilver(I) complexes, **223**, 161
- syntheses, spectroscopy and crystal structure of several binuclear silver(I) cryptates, **223**, 181
- a potentiometric study on the complexation of  $\text{Ag(I)}$  with various N-methyl and C-methyl substituted ethylenediamines in 1 M  $\text{NO}_3$ , **224**, 97
- electron transfer 122. Reductions of an  $\text{N}_4$ -coordinated complex of silver(III), **225**, 83
- Silyl complexes
- coordination chemistry of perhalogenated cyclopentadienes and alkynes XIV. Synthesis of dinuclear ring-bridged cymantrones with  $-\text{SiMe}_2-$  or  $-\text{TiCp}_2-$  bridges. Structures of  $[(\text{OC})_3\text{Mn}(\text{C}_5\text{Br}_4-)]_2\text{SiMe}_2$  and  $[(\text{OC})_3\text{Mn}(\text{C}_5\text{Cl}_4-)]_2\text{TiCp}_2$ , **218**, 173
- silicon-containing carbene complexes 16. Intramolecular agostic stabilization versus nucleophilic addition to the 16-electron carbene complex  $(\text{CO})_4(\text{L})\text{W}=\text{C}(\text{NMe}_2)\text{SiPh}_2\text{Me}$ , **220**, 73
- transition-metal silyl complexes 51. Preparation of the mixed silyl/stannyl complexes  $(\pi\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})(\text{SnPh}_3)(\text{SiR}_3)\text{H}$ , **224**, 177
- Single-valence systems
- the influence of spin-forbidden monomer excitations on spin-allowed electron transfer and electron-localized states of mixed-valence and single-valence dimeric systems, **226**, 33
- Solid-state fluorescence
- crystal structure and solid-state fluorescence of an oxomolybdenum(V)-benzylviologen electron donor-acceptor salt, **226**, 25
- Solid state polymerization
- synthesis, structure and solid state polymerization of diaqua-bis(propynoato)magnesium, **222**, 339
- Solution study
- Zn(II) complexes with thiopyrimidine derivatives: solution study, synthesis and crystal structure of a zig-zag chain zinc(II) complex with the ligand 4,6-dimethyl-2-thiopyrimidine, **221**, 177
- Solvato complexes
- non-isothermal spectrophotometric kinetics applied to inorganic reactions, **221**, 31
- Spectral overlap law
- relationship between four-state delineation and two-state delineation in electron transfer reactions and excitation transfer, **225**, 185
- Spectroelectrochemistry
- synthesis, electrochemistry and photoexcited-state properties of dinuclear ruthenium complexes bridged by 2,6'-bis(2-pyridyl)-2,2':6,2''-thiazolo[4,5-d]-benzothiazole, **226**, 17
- infrared spectroelectrochemistry of  $[\text{Rh}(\text{TM4})_4\text{M}(\text{CO})_5]_2^{2+}$  ( $\text{TM4} = 2,5$ -diisocyno-2,5-dimethylhexane;  $\text{M} = \text{Re}, \text{Mn}$ ): electron-transfer-promoted metal-metal bond cleavage and formation, **226**, 61
- a spectroelectrochemical method for evaluating factors which regulate the redox potential of hemoglobins, **226**, 187
- Spectrophotometric titration
- outer-sphere coordination of polycyanometallate anions with polyammonium macrocycles: a spectrophotometric study, **227**, 71
- Spectroscopic studies
- main group metal halide complexes with sterically hindered thioureas XIV. A reinvestigation of coordination sensitive modes in the solid state vibrational spectrum of 1,3-dimethyl-2(3*H*)-imidazolethione and related compounds, **218**, 133
- spectroscopic studies on complex formation and non-covalent interactions in ternary palladium(II) systems involving spermidine and 2,3-diaminopropionate or 2,4-diaminobutyrate, **219**, 107
- spectroscopic studies of the *trans*- $[\text{VOCl}_2(\text{H}_2\text{O})_2]$  chromophore in its hydrogen bonding adduct with 18-crown-6, **227**, 91
- Spectroscopy
- a study of the reactivity and structure of cyclic  $\alpha,\beta$ -unsaturated Fischer-type carbene complexes, **220**, 233
- photochemical and spectroscopic studies of complexes of iron(III) with citric acid and other carboxylic acids, **226**, 117
- Spermidine complexes
- spectroscopic studies on complex formation and non-covalent interactions in ternary palladium(II) systems involving spermidine and 2,3-diaminopropionate or 2,4-diaminobutyrate, **219**, 107
- Spin-allowed electron transfer
- the influence of spin-forbidden monomer excitations on spin-allowed electron transfer and electron-localized states of mixed-valence and single-valence dimeric systems, **226**, 33
- Spin-forbidden excitation
- the influence of spin-forbidden monomer excitations on spin-allowed electron transfer and electron-localized states of mixed-valence and single-valence dimeric systems, **226**, 33
- Spontaneous self-organization
- spontaneous formation of a 1D-coordination polymer with helix structure starting from (S)-pyroglutamic acid and copper(II) acetate, **222**, 5
- Stability constants
- solution studies of monovalent metal complex ions of the bibrachial lariat ether 7,16-bis(2-methoxyethyl)-1,4,10,13-tetraoxa-7,16-diazacyclooctadecane, **219**, 217
- stability constants of the lanthanide(III)-1,4,7,10-tetraazacyclododecane-*N,N',N'',N'''*-tetraacetate complexes, **221**, 165
- stabilities of 1,2-diethyl-3-hydroxy-4-pyridinone chelates of divalent and trivalent metal ions, **223**, 21
- coordination abilities of amide oxygens in a 24-membered ring pseudopeptide toward several transition metal ions in acetonitrile, **223**, 121
- a potentiometric study on the complexation of  $\text{Ag(I)}$  with various N-methyl and C-methyl substituted ethylenediamines in 1 M  $\text{NO}_3$ , **224**, 97
- stability of metal ion complexes of *N,N'*-bis(2-hydroxybenzyl)-ethylenediamine-*N,N'*-diacetic acid, **224**, 151
- Stannyl complexes
- transition-metal silyl complexes 51. Preparation of the mixed silyl/stannyl complexes  $(\pi\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})(\text{SnPh}_3)(\text{SiR}_3)\text{H}$ , **224**, 177
- Stereochemistry

- a unified view of inorganic stereochemistry and stereochemical changes, **219**, 187
- stereochemical investigation of cationic olefin complexes of platinum(II) containing chiral diamines: a sequence of four asymmetric centres (N, C, C and N) in the chelate ring, **227**, 261
- Stereoselective synthesis
- synthesis, structure, dynamics and first atropisomer-selective cleavage of the chromium tricarbonyl complex of a lactone-bridged biaryl, **222**, 255
- Stereoselectivity
- stereoselectivity in reactions of metal complexes Part XVIII. Kinetics and stereoselectivity in electron-transfer reactions between  $[\text{Co}(\text{L})\text{H}_2\text{O}]^+$  ( $\text{L} = \text{N}, \text{N}'$ -[(pyridine-2,6-diyl)bis(methylene)]bis[proline] (pomp) and  $\text{N}, \text{N}'$ -[(4-methoxy-pyridine-2,6-diyl)bis(methylene)]bis[proline] (MeO-pomp) and optically active iron(II) complexes, **218**, 73
  - stereoselective aldol addition reactions of Fischer carbene complexes via electronic tuning of the metal center for enolate reactivity, **220**, 215
- Steric effects
- influence of bulky substituents on the regioselective group-transfer reactions of diorganozinc compounds with  $\text{N}, \text{N}'$ -bis(2,6-di-isopropylphenyl)-1,4-diaza-1,3-butadiene, **220**, 55
  - the steric effect on the ring opening process of the decarboxylation of *cis*-carbonato-bis(diamine)cobalt(III) ions, **221**, 9
  - strain, silyl and steric effects on the regioselectivity of palladium(0) catalyzed allyl esters reduction as alternative to the Wittig reaction, **222**, 37
- Steroid complexes
- the synthesis and reactions of A-ring aromatic steroid complexes of manganese tricarbonyl, **220**, 297
- Styryl compounds
- electrophilic addition to styrylsilanes: sequential carbon-carbon bond forming reactions, **220**, 145
- Substituent dependent
- chemoselective functionalization of zirconacyclopentenes, **220**, 319
- Substituent independent
- chemoselective functionalization of zirconacyclopentenes, **220**, 319
- Substituted ruthenocene complexes
- synthesis and characterization of phosphane-substituted hydroxyruthenocenes: crystal structure of  $[\text{Ru}(\eta^5\text{-C}_5\text{H}_4\text{PPh}_3)(\eta^5\text{-C}_5\text{H}_4\text{OH})]\text{PF}_6$ , **217**, 203
- Succinate
- labeling studies of some carbonylation reactions of styrene with cationic palladium complexes, **220**, 273
- Sulfide complexes
- preparation and characterization of  $[(\text{LL})\text{Cl}_2\text{Mo}]_2(\mu\text{-SPr})_2$  compounds with  $\text{Pr} = \text{n-propyl}$  and  $\text{LL} = \text{PrSCH}_2\text{CH}_2\text{SPr}$  and  $\text{Et}_2\text{PCH}_2\text{CH}_2\text{PEt}_2$ . Further examples of C-S bond cleavage and oxidative addition across an Mo-Mo quadruple bond, **215**, 191
  - the reaction of  $\text{CpRu}(\text{PPh}_3)_2^+$  with trimethylenesulfide. Synthesis and molecular and crystal structure of  $[\text{CpRu}(\text{PPh}_3)_2(\text{SC}_3\text{H}_6)]\text{CF}_3\text{SO}_3$ , **221**, 157
- Sulfido complexes
- oxo- and sulfido-bridged sandwich cubane-type molybdenum-indium mixed-metal cluster. Synthesis and X-ray structure of  $[(\text{H}_2\text{O})_9\text{Mo}_3\text{S}_3\text{OInOS}_3\text{Mo}_3(\text{H}_2\text{O})_9](\text{CH}_3\cdot\text{C}_6\text{H}_4\cdot\text{SO}_3)_8\cdot 30\text{H}_2\text{O}$ , **216**, 13
- Sulfonate complexes
- structures of layered metal sulfonate salts: trends in coordination behavior of alkali, alkaline earth and transition metals, **216**, 101
- Sulfonium-methylidene complexes
- synthesis and reactions of the novel sulfonium-methylidene complex  $[\text{Fe}_2(\text{CO})_2(\text{cp})_2(\mu\text{-CO})\{\mu\text{-C}(\text{H})(\text{SMe}_2)\}]\text{SO}_3\text{CF}_3$ , **223**, 31
- Sulfur complexes
- 'insulation' to lattice effects on quadrupole splitting in the Mössbauer spectra of salts of  $[\text{Fe}_4\text{S}_4(\text{Cys})_4]^{2-}$ , a ferredoxin model, **217**, 3
  - electron transfer reactivity of cuboidal heterometallic  $\text{Mo}_3\text{MS}_4$  clusters in aqueous acidic solutions, **225**, 157
  - syntheses and structures of mixed-ligand tetrairon-sulfur clusters,  $[\text{Cp}'_2(\text{Ph}_2\text{C}_2\text{S}_2)_2\text{Fe}_4\text{S}_4](\text{PF}_6)$  and  $[\text{Cp}'_3(\text{Ph}_2\text{C}_2\text{S}_2)\text{Fe}_4\text{S}_3](\text{PF}_6)_n$  ( $n = 1$  and  $2$ ) ( $\text{Cp}' = \eta^5\text{-C}_5\text{Me}_5$ ). Structural change of the clusters accompanying electron transfer reactions, **225**, 229
  - electron transfer 120. Reductions of the cuboidal mixed-valence molybdenum/sulfur cluster,  $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]$ , **225**, 297
  - heteronuclear metal cluster complexes with a new coordination mode of the  $\text{Fe}_2\text{S}_2(\text{CO})_6$  unit. Synthesis, structure and possible reaction pathway of the first Fe-Cu-S cluster compound,  $[\text{Fe}_6\text{Cu}_5\text{S}_6(\text{CO})_{18}(\text{PPh}_3)_2]$ , **227**, 119
- Sulfur ligand complexes
- transition metal complexes with sulfur ligands CIII.  $[(\mu\text{-S}_2)\{\text{Fe}(\mu\text{-S}_4')\}_2]$ , a novel  $[2\text{Fe}-2\text{S}]$  cluster with octahedral low-spin  $[\text{Fe}^{\text{II}}\text{S}_6]$  centers ( $\text{S}_4'^{2-} = 1,2$ -bis(2-mercaptophenylthio)ethane(2-)), **224**, 35
  - transition metal complexes with sulfur ligands CVI. Phosphine and azido  $[\text{Fc}(\text{L})(\text{N}_3\text{S}_4)]$  complexes; structural characterization of the key complex  $[\text{Fe}(\text{N}_3\text{S}_4)] \cdot \text{THF}$  ( $\text{L} = \text{PMe}_3, \text{PBu}_3, \text{PMe}_2\text{Ph}, \text{PMePh}_2, \text{N}_3^-$ ;  $\text{N}_3\text{S}_4'^{2-} = \text{dianion of } 2,2'$ -bis(2-mercaptophenylthio)diethylamine), **224**, 61
- Superoxide dismutase
- EXAFS investigation on nickel substituted Cu,Zn bovine superoxide dismutase, **216**, 121
  - stereoselective electron transfer between Cu(II) superoxide dismutase and  $\Lambda$ - and  $\Delta$ - $[\text{Fe}(\text{pdta})]$ , **225**, 151
- Surfactant film
- electron transfer in surfactant films on electrodes; copper phthalocyaninetetrasulfonate-didodecyldimethylammonium bromide, **226**, 159
- Synthetic estrogens
- enantioselective synthesis of ferrocene analogs of hexestrol and estradiol; recognition towards estradiol receptors, **222**, 51
- Technetium complexes
- reactions of the technetium(V) phenylimido complex  $[\text{TcX}_3(\text{NPh})(\text{PPh}_3)_2]$  with aromatic thiolate ligands (where  $\text{X} = \text{Cl}, \text{Br}$ ), **218**, 97
  - synthesis and characterization of neutral oxorhenium(V) and nitridotechnetium(V) complexes with a tetradentate  $\text{N}_2\text{S}_2$  unsaturated ligand derived from dithiocarboxylic acid, **227**, 25
- Tellurido complexes
- synthesis and cyclic voltammetry of trinuclear selenido and tellurido clusters of nickel(II), palladium(II) and platinum(II) with 1,2-bis(diphenylphosphino)ethane and crystal structure of the tellurido cluster of platinum(II), **217**, 129
- Tellurium complexes
- systematics of some properties of hexamolybdotellurate(VI) salt hydrates with di- and trivalent metals,  $\text{M}_x[\text{TeMo}_6\text{O}_{24}] \cdot n\text{H}_2\text{O}$ , **217**, 93
  - synthesis, structure and properties of the polychalcogenides  $[\text{M}_4\text{Te}_{12}]^{4-}$  ( $\text{M} = \text{Cd}, \text{Hg}$ ), **224**, 163
- Terbium complexes
- luminescent europium(III) and terbium(III) ions encapsulated in a 2-aminoethoxy or carbamoyloxy substituted calixarene host, **227**, 293

## Terpyridine complexes

novel didentate 2,2':6',2''-terpyridine complexes of ruthenium(II), **223**, 177

volume profile analysis for intermolecular electron transfer between cytochrome *c* and Co(terpy)<sub>2</sub>, **225**, 95

## Tertiary phosphine complexes

the unusual crystallographic disorder of M<sub>2</sub> units in M<sub>2</sub>Cl<sub>4</sub>(PEt<sub>3</sub>)<sub>4</sub> (M = Re, Mo), **215**, 103

## Tetraazaannulene complexes

fluorine-substituted tetraazaannulene complexes: syntheses and characterizations, **223**, 187

## Tetraazamacrocyclic complexes

N<sup>1</sup>,N<sup>7</sup>-expeditious dialkylation of cyclen (1,4,7,10-tetraazacyclododecane). An astonishing reactivity of cyclen tricarbonyl molybdenum and chromium complexes, **220**, 347

## Tetracyanoethylene compounds

[Rh<sub>2</sub>(O<sub>2</sub>CCH<sub>3</sub>)<sub>4</sub>TCNE]<sub>x</sub> polymers: isomers with either 1,1- or *trans*-1,2-TCNE bridges, **221**, 1

## Tetranuclear complexes

di- and tetranuclear gold(II) complexes with dithiocarbamate and related ligands. X-ray structure of [Au<sub>2</sub>(μ-CH<sub>2</sub>PPh<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>(S<sub>2</sub>CNMe<sub>2</sub>)<sub>2</sub>], **223**, 55

## Tetratolylporphyrinatoiron(III) complexes

redox reactions between oxomolybdenum(IV) and tetratolylporphyrinatoiron(III) complexes, **226**, 259

## Thallium complexes

synthesis of bis- or tris(pentafluorophenyl) thallium(III) complexes with bidentate ligands. X-ray crystal structure of [Tl(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>(acac){(O)Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub>(O)}], **215**, 229

dithiocarbamates of dimethylthallium(III). Crystal structure of dimethyl(di-*n*-propyldithiocarbamate)thallium(III), **216**, 15

## Thermal decomposition

thermal decomposition of cobalt carbonyl complexes in viscous media, **227**, 233

## Thermodynamics

equilibrium and kinetic study of piperidine binding to phthalocyaninatoiron(II) in dimethyl sulfoxide, **219**, 199

thermodynamics of complexation of the uranyl(VI) ion with benzoate. The crystal structure of sodium tris(benzoate) dioxouranate dihydrate, Na[(PhCO<sub>2</sub>)<sub>3</sub>UO<sub>2</sub>]·2H<sub>2</sub>O, **223**, 151

volume profile analysis for intermolecular electron transfer between cytochrome *c* and Co(terpy)<sub>2</sub>, **225**, 95

## Thermometry

thermometric determination of electrochemical Peltier heat (thermal effect associated with electron transfer) of some redox couples, **226**, 69

## Thiocarboxylato complexes

molybdenum complexes with sterically-hindered thio-carboxylate ligands. Electrochemical properties of the anionic complex bis(2,2-diphenyl-2-mercaptoethanoate)dioxomolybdate(VI) in protic solvents, **221**, 117

## Thiocyanato complexes

crystal structure and magnetic properties of the first 1D nickel(II) complex with bridging thiocyanato ligands in *cis* position: {catena-(μ<sub>N,S</sub>-NCS)}[Ni(en)<sub>2</sub>](PF<sub>6</sub>), **218**, 185

## Thioether complexes

transition metal complexes with sulfur ligands CIV. Protonation and alkylation of thiolate donors of [Fe(CO)<sub>2</sub>(S<sub>4</sub><sup>2-</sup>)] and [Fe(CO)(S<sub>5</sub><sup>2-</sup>)]: syntheses and properties of Fe(II) complexes with open-chain thioether ligands (S<sub>4</sub><sup>2-</sup> = 1,2-bis(2-mercaptophenylthio)ethane(2-); S<sub>5</sub><sup>2-</sup> = 2,2'-bis(2-mercaptophenylthio)diethylsulfide(2-)), **224**, 45

## Thiolate complexes

synthesis and characterization of platinum(II)-thiolate complexes. Crystal and molecular structures of *cis*-Pt(Ph<sub>3</sub>P)<sub>2</sub>(SC<sub>6</sub>H<sub>2</sub>-2,4,6-*i*Pr<sub>3</sub>)Cl and *cis*-Pt(Ph<sub>3</sub>P)<sub>2</sub>(SC<sub>6</sub>H<sub>2</sub>-2,4,6-*i*Pr<sub>3</sub>)<sub>2</sub>, **216**, 83

reactions of the technetium(V) phenylimido complex

[TcX<sub>3</sub>(NPh)(PPh<sub>3</sub>)<sub>2</sub>] with aromatic thiolate ligands (where X = Cl, Br), **218**, 97

## Thiolato complexes

the reactions of thiolate ligands with rhenium hydrides Part 1.

The syntheses and structures of [ReO(SC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>)<sub>3</sub>(PPh<sub>3</sub>)] and [Re(SC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>)<sub>3</sub>(PPh<sub>2</sub>H)(PPh<sub>3</sub>)], **223**, 63

## Thione complexes

main group metal halide complexes with sterically hindered thioureas XIV. A reinvestigation of coordination sensitive modes in the solid state vibrational spectrum of 1,3-dimethyl-2(3*H*)-imidazoethione and related compounds, **218**, 133

## Thiopyridyl triazine complexes

thiopyridyl triazine derivatives and their platinum complexes: a new class of potential antitumor agents, **227**, 175

## Thiopyrimidine complexes

Zn(II) complexes with thiopyrimidine derivatives: solution study, synthesis and crystal structure of a zig-zag chain zinc(II) complex with the ligand 4,6-dimethyl-2-thiopyrimidine, **221**, 177

## Thiosemicarbazone complexes

diorganotin(IV) derivatives of salicylaldehydethiosemicarbazone. The crystal structure of dimethyl- and diphenyl-(salicylaldehydethiosemicarbazonato)tin(IV), **216**, 169

new heptacoordinated tin complexes of 2,6-diacetylpyridine-bis(thiosemicarbazone), H<sub>2</sub>daptsc, and of 2,6-diacetylpyridine-bis(semicarbazone), H<sub>2</sub>dapsc. Crystal and molecular structures of [MeSnCl(Hdaptsc)]Cl·MeOH and [MeSnCl(H<sub>2</sub>dapsc)]Cl<sub>2</sub>·2H<sub>2</sub>O, **218**, 139

synthesis and spectroscopic properties of diorganotin(IV) derivatives of 2,6-diacetylpyridine bis(thiosemicarbazone). Crystal structure of diphenyl[2,6-diacetylpyridine bis(thiosemicarbazonato)}tin(IV) bis(dimethylformamide) solvate, **221**, 61

synthesis, spectroscopic and structural characterization of methyl pyruvate- and pyridoxal-hydrazinopyruvylthiosemicarbazones, **223**, 77

## Thiourea compounds

main group metal halide complexes with sterically hindered thioureas XIV. A reinvestigation of coordination sensitive modes in the solid state vibrational spectrum of 1,3-dimethyl-2(3*H*)-imidazoethione and related compounds, **218**, 133

## Thiourea-functionalized xerogels

formation of an anchored copper(II) complex on a thiourea-functionalized silica gel by *in situ* modification of the tethered ligating function, **221**, 183

## Tin alkyne complexes

hydrozirconation/transmetalation of acetylenic stannanes. New 1,1-dimetallo reagents, **220**, 41

## Tin complexes

cyclic bis(amino)organotin cations, stabilized by π-coordination and new spirocyclic stannole derivatives, **216**, 51

diorganotin(IV) derivatives of salicylaldehydethiosemicarbazone. The crystal structure of dimethyl- and diphenyl-(salicylaldehydethiosemicarbazonato)tin(IV), **216**, 169

synthesis, characterization and crystal structure of (C<sub>3</sub>H<sub>5</sub>N<sub>2</sub>)<sub>2</sub>[Sn<sub>2</sub>(NO<sub>3</sub>)<sub>4</sub>(μ-OH)<sub>2</sub>Me<sub>4</sub>], a di-μ-hydroxo-bridged compound with heptacoordinated tin(IV), **216**, 257

equilibrium and spectroscopic studies of diethyltin(IV) complexes formed with carbohydrate derivatives of thiazolidine-4-carboxylic acid, **218**, 65

new heptacoordinated tin complexes of 2,6-diacetylpyridine-bis(thiosemicarbazone), H<sub>2</sub>daptsc, and of 2,6-diacetylpyridine-bis(semicarbazone), H<sub>2</sub>dapsc. Crystal and molecular structures of [MeSnCl(Hdaptsc)]Cl·MeOH and



- [MeSnCl(H<sub>2</sub>dapsc)]Cl<sub>2</sub>·2H<sub>2</sub>O, **218**, 139
- ultrasound-promoted palladium-catalyzed carbonyl allylation by allylic alcohols with tin(II) chloride in non-polar solvents, **220**, 155
- 1,1-organoboration of tri-1-alkynyltin compounds: novel triorganotin cations, stannoles, 3-stannolenes and 1-stanna-4-bora-2,5-cyclohexadienes, **220**, 161
- synthesis and spectroscopic properties of diorganotin(IV) derivatives of 2,6-diacetylpyridine bis(thiosemicarbazone). Crystal structure of diphenyl{2,6-diacetylpyridine bis(thiosemicarbazonato)}tin(IV) bis(dimethylformamide) solvate, **221**, 61
- formation of an allyl–tin bond in  $\eta^3$ -allyl(trichlorostannyl)palladium(II) complexes, **222**, 21
- direct electrochemical synthesis of tin(IV) complexes of 2-(2-mercaptoethyl)iminomethyl-phenols; the molecular structure of bis{2-(2-mercaptoethyl)iminomethyl-5-bromo-phenoxy}tin(IV), **223**, 87
- Tin ions**
- metal–metal interactions of aqueous lead(II) and tin(II) ions with the ground and excited states of tetrakis( $\mu$ -diphosphito)diplatinate(II), **225**, 201
- Titanacyclobutene complexes**
- nitrosoarene and nitrosoalkane insertion reactions of titanacyclobutenes, **222**, 305
- Titanium complexes**
- designed binaphthyl-derived titanium complexes: a new type of asymmetric catalyst for the carbonyl-ene reaction with glyoxylate, **222**, 71
- the preparation of chiral titanium reagents: a comparison of new and known ligands, **222**, 95
- carbon dioxide fixation by Cp<sub>2</sub>( $\eta^3$ -allyl)Ti complexes generated from various dienes, **222**, 145
- nitrosoarene and nitrosoalkane insertion reactions of titanacyclobutenes, **222**, 305
- early/late bimetallic complexes: MCo<sub>2</sub> (M = Ti, Zr) derivatives, **227**, 201
- Transition metal complexes**
- complexes of tetradentate  $\beta$ -keto enamine Schiff bases: an electropray mass spectrometric study, **216**, 75
- structures of layered metal sulfonate salts: trends in coordination behavior of alkali, alkaline earth and transition metals, **216**, 101
- Raman and infrared spectroscopic studies of tris triphenyl phosphine chloride complexes of transition metals, **216**, 191
- transmetalation of targets ( $\mu$ -Y)N<sub>4</sub>Cu<sub>2</sub>Cu<sub>2</sub><sup>II</sup>Cl<sub>4</sub> and ( $\mu$ -Y, $\mu$ -O)N<sub>4</sub>Cu<sub>4</sub>Cl<sub>4</sub> (Y = 3,4,5,6-tetrachlorocatecholate) with M(NS)<sub>2</sub> reagents, **217**, 85
- estrogen derivatives of transition metal complexes for analytical detection enhancement. Part II, **218**, 207
- influence of pendant cyanoalkyl side-arms on the stabilities of metal ion complexes of monoaza- and diaza-18-crown-6 ethers in methanol, **219**, 31
- liquid matrix induced processes in fast atom bombardment mass spectrometry of polyazacrown transition metal complexes, **219**, 85
- a novel reaction of the cyano ligand opening an organometallic route to 4-amino imidazoles: four component condensation (4CC) with isocyanide, aldehyde and amine, **222**, 275
- electron transfer reactivity of cuboidal heterometallic Mo<sub>3</sub>MS<sub>4</sub> clusters in aqueous acidic solutions, **225**, 157
- Transition metal ion complexes**
- coordination abilities of amide oxygens in a 24-membered ring pseudopeptide toward several transition metal ions in acetonitrile, **223**, 121
- Transmetalation**
- transmetalation of targets ( $\mu$ -Y)N<sub>4</sub>Cu<sub>2</sub>Cu<sub>2</sub><sup>II</sup>Cl<sub>4</sub> and ( $\mu$ -Y, $\mu$ -O)N<sub>4</sub>Cu<sub>4</sub>Cl<sub>4</sub> (Y = 3,4,5,6-tetrachlorocatecholate) with M(NS)<sub>2</sub> reagents, **217**, 85
- hydrozirconation/transmetalation of acetylenic stannanes. New 1,1-dimetallo reagents, **220**, 41
- Triazole ligand complexes**
- synthesis and X-ray structure of the dinuclear  $\mu$ -5,5'-diamino-3,3'-bis-1,2,4-triazolato(1-)-bis(diethylenetriaminocopper(II)) triperchlorate, [{Cu(dien)}<sub>2</sub>Hdabt](ClO<sub>4</sub>)<sub>3</sub>, **227**, 181
- Tridentate ligand complexes**
- neutral transition metal complexes with the tridentate ligand 4-(5-methyl-4-imidazolyl)-3-thiabutanoic acid. A series of isostructural octahedral compounds, **215**, 151
- transition metal compounds of the tridentate pyrazole substituted amine ligand bis(2-(3,5-dimethyl-1-pyrazolyl)ethyl)ethylamine (ddae). X-ray structures of [Co(ddae)(NO<sub>3</sub>)<sub>2</sub>], [Cu(ddae)(NO<sub>3</sub>)(H<sub>2</sub>O)](NO<sub>3</sub>) and [Cu(ddae)(Cl)<sub>2</sub>]·C<sub>2</sub>H<sub>5</sub>OH, **216**, 43
- zinc(II) and cadmium(II) complexes of the potentially tridentate ligands 2-[[2-pyridinylmethyl]amino]methyl]phenol (HSALAMP) and 2-[[[2-(2-pyridinyl)ethyl]amino]methyl]phenol (HSALAEF). X-ray crystal structures of the [Zn(SALAMP)(NO<sub>3</sub>)<sub>2</sub>] and the [Cd(SALAEF)(NO<sub>3</sub>)<sub>2</sub>] dimers, **219**, 55
- Tridentate phosphine complexes**
- syntheses of ruthenium–iridium heterobimetallic complexes using tridentate phosphine ligands, **227**, 79
- Trifluoroacetate complexes**
- orthometalation reactions in trifluoroacetate dirhodium(II) compounds. Molecular structure of Rh<sub>2</sub>(O<sub>2</sub>CCF<sub>3</sub>)<sub>2</sub>·[(C<sub>6</sub>H<sub>4</sub>)PPh<sub>2</sub>]<sub>2</sub>·(PPh<sub>3</sub>)<sub>2</sub>·2(C<sub>7</sub>H<sub>8</sub>), **218**, 189
- Triiodide complexes**
- the structure of (CH<sub>3</sub>)<sub>3</sub>Si<sub>3</sub>. Comparison between the structure in the solid and liquid state, **215**, 33
- Trimetallic complexes**
- trimetallic complexes of the tripod ligand 1,1,1-tris(diphenylphosphinomethyl)ethane Part I. Homotrimetallic *cis,cis,cis* and *cis,trans,cis* complexes of the type [(PtX<sub>2</sub>- $\eta$ -tripod)<sub>2</sub>PtX<sub>2</sub>] (X = Cl, I and CH<sub>3</sub>), **221**, 187
- Trinuclear complexes**
- synthesis and characterization of symmetric and unsymmetric oxo-bridged trinuclear chromium benzoate complexes: crystal and molecular structure of [Cr<sub>3</sub>O(O<sub>2</sub>CPh)<sub>6</sub>(py)<sub>3</sub>]ClO<sub>4</sub>, **217**, 171
- synthesis and crystal structure of the  $\mu_2$ -[tris[bis-(diphenylphosphine)methane]]- $\mu_3$ -3-methyl-8-ethylxanthine-N,O,O]- $\mu_3$ -chloro-triangle-tricopper(I)-water complex exhibiting a novel oxopurine coordination mode, **227**, 43
- Triply bonded complexes**
- a new type of face-sharing bioctahedral triply bonded dinobium cluster (PyH)[Nb<sub>2</sub>( $\mu$ -Cl)<sub>2</sub>( $\mu$ -THT)Cl<sub>3</sub>(Py)<sub>3</sub>](THF)(C<sub>6</sub>H<sub>6</sub>)<sub>0.5</sub>, made by a designed synthesis, **227**, 191
- Tripodal ligand complexes**
- crystal structure of a binuclear vanadium(III) complex with a new tripodal ligand, [V<sub>2</sub>Cl<sub>4</sub>(tped)(EtOH)<sub>2</sub>]·2EtOH, **224**, 199
- Tripod ligand complexes**
- organometallic chelating ligands. Synthesis of a new anionic tripod like ligand with an N,N',O-donor set. X-ray crystal structure of [( $\eta^6$ -C<sub>6</sub>Me<sub>6</sub>)Ru( $\mu$ -P(O)(OMe)<sub>2</sub>)( $\mu$ -Pz)<sub>2</sub>Re(CO)<sub>3</sub>](Me = methyl, Pz = pyrazolate), **219**, 67
- Trivalent ion complexes**
- stabilities of 1,2-diethyl-3-hydroxy-4-pyridinone chelates of divalent and trivalent metal ions, **223**, 21
- Tungstate complexes**

- coordination study of *D*-xylo 5-hexulosonic acid with borate, molybdate and tungstate in aqueous solution by nuclear magnetic resonance spectroscopy, **221**, 69
- Tungsten complexes**
- evidence of the  $W(O)(S_2)_2$  core as an intermediate in the acidification of  $WS_4^{2-}$ . Structural characterization of  $W(O)(S_2)_2(\text{bpy})$  and  $W(O)(S_2)_2(\text{phen})$ , **216**, 201
- synthesis and characterization of molybdenum and tungsten nitrite complexes of the type  $[M(\text{NO}_2)_2(\text{CO})_2(\text{PPh}_3)_2]$ , **218**, 195
- the study of isolobal displacement reactions. Synthesis and structures of the heterobimetallic  $\mu$ -alkyne cluster complexes  $\eta^5\text{-RC}_3\text{H}_4(\text{CO})_2\text{M}(\mu\text{-R}^1\text{C}_2\text{R}^2)\text{Co}(\text{CO})_3$  ( $\text{M} = \text{Mo}, \text{W}$ ;  $\text{R} = \text{CH}_3\text{C}(\text{O}), \text{CH}_3\text{O}_2\text{C}$ ), **219**, 93
- some new but-2-yne phosphine complexes of tungsten(II), **219**, 99
- synthesis and characterisation of cyanide bridged heterobinuclear complexes based on Group VI octacyanometalate(IV) and pentaammineosmium(III) ions: a comparison with Group VIII hexacyano analogues, **219**, 121
- silicon-containing carbene complexes 16. Intramolecular agostic stabilization versus nucleophilic addition to the 16-electron carbene complex  $(\text{CO})_4(\text{L})\text{W}=\text{C}(\text{NMe}_2)\text{SiPh}_2\text{Me}$ , **220**, 73
- hydroformylation of alkenes and alkynes using a heterobinuclear Rh–W catalyst, **220**, 187
- preparation of  $\eta^1$ -1,3-dienyl,  $\eta^1$ -1,2-dienyl and  $\eta^3$ -cyclobutenone complexes via reactions of transition-metal carbonyl containing anions with allenic electrophiles, **220**, 305
- bicyclic carbo- and heterocycles from an allylidene complex and cyclic 1,3-dienes by tandem cyclopropanation/Cope rearrangement, **220**, 327
- investigation of the dithiocarbamate-induced coupling of allylidene and carbonyl ligands on a tungsten center, **220**, 335
- carbonyl carbene coupling on cationic  $\eta^2$ -carbene complexes of molybdenum and tungsten, **222**, 77
- manganese and tungsten carbene complex mediated intramolecular Diels–Alder reactions, **222**, 291
- synthesis and structural analysis of *trans*- $\text{Cp}(\text{CO})_2(\text{PPh}_3)\text{WHgCl}$ , **227**, 145
- Two-state delineation**
- relationship between four-state delineation and two-state delineation in electron transfer reactions and excitation transfer, **225**, 185
- Ultrasound**
- ultrasound-promoted palladium-catalyzed carbonyl allylation by allylic alcohols with tin(II) chloride in non-polar solvents, **220**, 155
- Unsaturated  $N_2S_2$  ligand**
- synthesis and characterization of neutral oxorhenium(V) and nitridotechnetium(V) complexes with a tetradentate  $N_2S_2$  unsaturated ligand derived from dithiocarboxylic acid, **227**, 25
- Uracil complexes**
- simultaneous binding of soft and hard metals to a pyrimidine nucleobase: *trans*- $\text{K}_2[\text{PtI}_2(1\text{-MeU})_2]\cdot 6\text{H}_2\text{O}$ , an anionic Pt-1-methyluracil (1-MeU) complex arranged in circles, **218**, 117
- analysis of  $[(\text{dien})\text{Pd}]^{2+}$  binding to uracil and azauracils by proton NMR spectroscopy, **223**, 131
- Urania compounds**
- electron transfer reactions in non-stoichiometric ceria and urania, **225**, 305
- Uranium complexes**
- uranium hexafluoride: laboratory scale synthesis, coordination behaviour towards nitriles and Raman spectra, **215**, 5
- synthesis and characterization of uranium triiodide tetrakis(acetonitrile), **218**, 203
- uranium compounds as catalysts for aldolisation reactions, **222**, 91
- thermodynamics of complexation of the uranyl(VI) ion with benzoate. The crystal structure of sodium tris(benzoato) dioxouranate dihydrate,  $\text{Na}[(\text{PhCO}_2)_3\text{UO}_2]\cdot 2\text{H}_2\text{O}$ , **223**, 151
- Uridine complexes**
- aggregation of platinum–uridine blue as a function of concentration in aqueous solution by small angle X-ray scattering, **216**, 37
- Valence tautomerism**
- valence tautomerism for catechol/semiquinone complexes of the *trans*- $\text{M}(\text{Bupy})_2(3,6\text{-DBQ})_2$  ( $\text{M} = \text{Mn}, \text{Fe}, \text{Co}$ ) series, **226**, 91
- Vanadium complexes**
- phosphine and phosphine oxide adducts of vanadium(IV) chloride: synthesis and structural studies, **215**, 47
- phosphate complexation of oxovanadium(IV). Evidence of bidentate chelation of orthophosphate, **215**, 225
- synthesis, spectral properties and redox behaviour of oxovanadium(IV) complexes with tetraaza[14]annulenes, **216**, 217
- hydrothermal synthesis and crystal and molecular structure of a three-dimensional vanado-arsenate  $(\text{NH}_4)_3[(\text{VO})(\text{V}_2\text{O}_3)(\text{AsO}_4)_2(\text{HASO}_4)]$ , exhibiting both  $\{\text{V}-\text{O}-\text{V}\}_\infty$  chains and isolated  $\{\text{V}_2\text{O}_3\}$  units, **218**, 59
- bis(2-mercaptopyridine-*N*-oxide)oxovanadium(IV): synthesis, structure and its conversion to the dichloro complex, **219**, 213
- synthesis, characterization and crystal structures of two *cis*-dioxo vanadium(V) complexes of monoanionic tridentate Schiff base ligands, **223**, 193
- crystal structure of a binuclear vanadium(III) complex with a new tripodal ligand,  $[\text{V}_2\text{Cl}_4(\text{tped})(\text{EtOH})_2]\cdot 2\text{EtOH}$ , **224**, 199
- spectroscopic studies of the *trans*- $[\text{VOCl}_2(\text{H}_2\text{O})_2]$  chromophore in its hydrogen bonding adduct with 18-crown-6, **227**, 91
- Vinylbromides**
- chelo- and anticheleselectivity IV. Reaction-accelerating neighboring-group effects in the cross-coupling of vinylbromides with alkyl-transition metal reagents, **220**, 85
- Vinylterpyridine complexes**
- electrocatalytic reduction of  $\text{CO}_2$  and  $\text{O}_2$  with electropolymerized films of vinyl-terpyridine complexes of Fe, Ni and Co, **225**, 285
- Viologen derivatives**
- photoinduced electron transfer reaction between  $[\text{Cu}(\text{dmp})\text{P}_2]^{+\alpha-}$  ( $\text{P} = \text{PPh}_3$  or  $\text{PPh}_2(m\text{-C}_6\text{H}_4\text{SO}_3^-)$ ) and viologen derivatives, **225**, 261
- Voltammetric techniques**
- synthesis, molecular structure and electrochemical investigation of the 1,2,5-thiadiazole-3,4-dithiolate-substituted complex  $\text{Cp}^*\text{Ru}(\text{NO})(\text{tdas})$ . Evidence for nitrosyl bending during reduction, **226**, 267
- Volume of activation**
- volumes of activation for  $\beta$ -elimination and related reactions of chromium(III)–alkyl complexes in aqueous solutions, **227**, 57
- Water oxidation**
- dynamical investigations of the catalytic mechanisms of water oxidation by the  $[(\text{bpy})_2\text{Ru}(\text{OH}_2)_2]\text{O}^{4+}$  ion, **226**, 179
- X-ray diffraction**
- optically active  $\beta$ -ketoiminato manganese(III) complexes as efficient catalysts in enantioselective aerobic epoxidation of unfunctionalized olefins, **220**, 283
- X-ray diffraction analysis

- synthesis, molecular structure and electrochemical investigation of the 1,2,5-thiadiazole-3,4-dithiolate-substituted complex  $\text{Cp}^*\text{Ru}(\text{NO})(\text{tdas})$ . Evidence for nitrosyl bending during reduction, **226**, 267
- X-ray scattering  
aggregation of platinum–uridine blue as a function of concentration in aqueous solution by small angle X-ray scattering, **216**, 37
- Zeolite complexes  
preparation and dioxygen binding property of a new zeolite encapsulated cobalt tetramethylporphyrin complex, **217**, 135
- Zinc  
stereoselective electron transfer between  $\text{Cu}(\text{II})$  superoxide dismutase and  $\Lambda$ - and  $\Delta$ -[Fe(pdta)], **225**, 151
- Zinc complexes  
the synthesis and crystal structure of  $[\text{Cr}(\text{acacen})\text{py}_2][\text{ZnCl}_3\text{py}]$ , **217**, 195  
main group metal halide complexes with sterically hindered thioureas XIV. A reinvestigation of coordination sensitive modes in the solid state vibrational spectrum of 1,3-dimethyl-2(3*H*)-imidazolethione and related compounds, **218**, 133  
zinc(II) and cadmium(II) complexes of the potentially tridentate ligands 2-[[2-pyridinylmethyl]amino]methylphenol (HSALAMP) and 2-[[[2-(2-pyridinyl)ethyl]amino]methyl]phenol (HSALAEF). X-ray crystal structures of the  $[\text{Zn}(\text{SALAMP})(\text{NO}_3)_2]$  and the  $[\text{Cd}(\text{SALAEF})(\text{NO}_3)_2]$  dimers, **219**, 55  
the synthesis, structure and hydrolytic activity of a functional pentacoordinate zinc complex, **219**, 205
- Zn(II) complexes with thiopyrimidine derivatives: solution study, synthesis and crystal structure of a zig-zag chain zinc(II) complex with the ligand 4,6-dimethyl-2-thiopyrimidine, **221**, 177  
effect of aromatic–aromatic interaction on ligand binding to zinc porphyrins, **224**, 157
- Zirconacyclopentene complexes  
chemoselective functionalization of zirconacyclopentenes, **220**, 319
- Zirconium complexes  
a tris-( $\mu_2$ -I) face-sharing dizirconium(III) complex,  $[\text{1,4-Zr}_2\text{I}_7(\text{PEt}_3)_2]^-$ ; the first such dizirconium species, **216**, 9  
synthesis and structure of  $\text{Zr}_2\text{Cl}_4(\eta^6\text{-C}_6\text{H}_5\text{PMe}_2)_2(\text{PMe}_2\text{Ph})_2$ , an arene complex of zirconium(II), **217**, 187  
alkyl for hydride exchange between alane-trimethylamine and Group IVB metal alkyls, **218**, 47  
hydrozirconation/transmetalation of acetylenic stannanes. New 1,1-dimetallo reagents, **220**, 41  
chemoselective functionalization of zirconacyclopentenes, **220**, 319  
synthesis of stable tail-to-tail-conjugated primary enamines containing hetarene substituents, **222**, 193  
the  $\text{Zr}_6\text{Cl}_{12}(\text{PMe}_2\text{Ph})_6$  molecule revisited: a directed synthesis, **223**, 93  
early/late bimetallic complexes:  $\text{MCo}_2$  (M=Ti, Zr) derivatives, **227**, 201
- Zirconocene complexes  
synthesis of stable tail-to-tail-conjugated primary enamines containing hetarene substituents, **222**, 193
- Zwitterion alkynyl complexes  
1,1-organoboration of tri-1-alkynyltin compounds: novel tri-organotin cations, stannoles, 3-stannolenes and 1-stanna-4-bora-2,5-cyclohexadienes, **220**, 161