

Corrigendum

Inorganica Chimica Acta, 158 (1989) 217–220

Structure of Bis(2,6-dimethylpyridine)copper(I) Perchlorate. The Relationship Between the Cu–N(pyridine) Distance and Cu···CH₃(methyl group) ContactsMEGUMU MUNAKATA, SUSUMU KITAGAWA, HISAO SHIMONO (*Osaka, Japan*) and HIDEKI MASUDA (*Okazaki, Japan*)

Page 217, Table 2, should read as follows:

TABLE 2. Final Atomic Coordinates for [Cu(2,6-Me₂py)₂]ClO₄ (polymorph 2)^a

Atom	<i>x</i>	<i>y</i>	<i>z</i>
Cu	0.7513(0)	0.1296(3)	1.0013(0)
Cl	0.7498(9)	0.6154(5)	1.0022(8)
C(1)	0.8956(11)	0.1663(19)	0.9391(9)
C(2)	0.9269(11)	0.1549(24)	0.8751(12)
C(3)	0.8663(16)	0.0781(33)	0.7894(15)
C(4)	0.7688(16)	0.0193(27)	0.7633(12)
C(5)	0.7329(12)	0.0444(24)	0.8229(11)
C(6)	0.9614(14)	0.2577(22)	1.0334(14)
C(7)	0.6245(16)	-0.0282(26)	0.7933(13)
C(8)	0.6110(14)	0.1788(24)	1.0650(15)
C(9)	0.5682(15)	0.1549(29)	1.1212(13)
C(10)	0.6408(14)	0.0816(25)	1.2149(12)
C(11)	0.7386(17)	0.0219(30)	1.2411(13)
C(12)	0.7759(15)	0.0369(23)	1.1821(12)
C(13)	0.5395(14)	0.2615(33)	0.9630(13)
C(14)	0.8850(15)	-0.0033(32)	1.2060(14)
N(1)	0.7954(9)	0.1180(14)	0.9140(8)
N(2)	0.7091(12)	0.1048(24)	1.0942(11)
O(1)	0.8213(18)	0.6994(33)	0.9877(22)
O(2)	0.7119(17)	0.7253(31)	1.0317(15)
O(3)	0.7812(19)	0.4738(24)	1.0507(14)
O(4)	0.6477(15)	0.5813(30)	0.9035(12)

^ae.s.d.s. given in parentheses.