

is based on the three-variable expansion of the Mori equations [4, 5], adapted to anisotropic systems, where the generalized spherical harmonics are no longer independent variables.

The Fourier transforms  $J_{MN}(\omega)$  of the correlation functions  $G_{MN}(t) = (\delta D^M(t), \delta D^N(0)^x)$  are found to be:

$$J_{MN}(\omega) = \sum_{Ki} \sum_{Mi} X_{Mi} X_{iK}^{-1} G_{KN}(0) \times \\ \times \{i\omega + \lambda_i K_0 [i\omega + K_1 / (i\omega + \gamma)]^{-1}\}^{-1}$$

where  $X$  and  $\{\lambda\}$  are eigenfunctions and eigenvalues of the anisotropic diffusion operator. The expansion parameters  $K_0, K_1$  are related to mean-square angular velocity and (total) torque  $N$ , and  $1/\gamma$  to the torque relaxation time. The diffusion equation results are recovered under 'strong anisotropic interaction limit' (SAIL) conditions [6],  $K_1/K_0 = N/kT \gg 1$ .

#### References

- 1 P. L. Nordio and U. Segre, 'Rotational Dynamics', in *The Molecular Physics of Liquid Crystals*, G. W. Gray and G. R. Luckhurst, Eds., Academic Press, London (1979).
- 2 A. De Vries, *J. Chem. Phys.*, 71, 25 (1972).
- 3 F. Volino, A. J. Dianoux and H. Hervet, *Mol. Cryst. Liq. Cryst.*, 38, 25 (1977).
- 4 D. Kivelson and T. Keyes, *J. Chem. Phys.*, 57, 4599 (1972).
- 5 G. J. Evans and M. W. Evans, *J. Chem. Soc. Faraday II*, 73, 285 (1977).
- 6 G. Moro and P. L. Nordio, *Chem. Phys.*, 43, 303 (1979).

### The Effect of Solute Structure on the Nematic-Isotropic Transition in Binary Mixtures

D. E. MARTIRE\*, G. A. OWEIMREEN and F. DOWELL

Department of Chemistry, Georgetown University, Washington, D.C., 20057 U.S.A.

Systematic thermodynamic and statistical-mechanical studies of nematic-isotropic (NI) phase equilibria in binary mixtures have provided significant information on the effects of molecular size, shape and flexibility on the orientational order and stability of nematic mesophases [1-3].

The addition of solute to a nematogenic solvent either depresses or elevates the NI transition temperature ( $T_{NI}$ ) of the pure solvent and gives a two-phase region. The phase diagram in the  $T^*-x_2$  plane  $T^* = T/T_{NI}$ ;  $x_2 =$  solute mole fraction) yields coexistence curves that are virtually linear for  $x_2 < \sim 0.10$ . Of interest is the negative of the slope of the lower phase-boundary line (nematic/nematic + isotropic),  $\beta_N = -(dT^*/dx_2)_N$ , which is a measure of the order-destroying (positive  $\beta_N$ ) or order-enhancing (negative  $\beta_N$ ) ability of a solute.

Experimental  $\beta_N$  results are presented for mixtures of quasispherical (tetra-n-alkyl tins), chain-like (n-

alkanes) and rodlike (*p*-polyphenyls) solutes dissolved in nematogenic solvents (MBBA and 5CB). These results are compared with the predictions of lattice models, which stress the predominant role of repulsive interactions.

#### References

- 1 D. E. Martire, 'The Molecular Physics of Liquid Crystals', G. R. Luckhurst and G. W. Gray, eds., Academic Press, London (1979); chaps. 10 and 11.
- 2 D. E. Martire and F. Dowell, *J. Chem. Phys.*, 70, 5914 (1979), and references therein.
- 3 G. A. Oweimreen and D. E. Martire, *J. Chem. Phys.*, 72, (1980), and references therein.

### Properties of Amphiphilic Nematic Systems†

A. SAUPE, T. HAVEN and L. J. YU

Liquid Crystal Institute, Kent State University, Kent, Ohio 44242, U.S.A.

Nematic liquid crystalline states formed in mixtures of surfactant and water correspond to anisotropic micelle solutions with anisometric surfactant aggregates of a finite size. Addition of salts and co-surfactants stabilize in general the nematic state. An addition of chiral compounds leads to the formation of cholesteric states.

There are two different nematic states and direct transitions between the states are possible. The transition is weakly first order. The temperature dependence of the nematic order parameter can be positive or negative. It differs in general significantly from that found in thermotropic nematics. Curvature elastic and viscous properties seems to be qualitatively the same as in thermotropic nematics. Results on curvature elasticity coefficients obtained on the decylammoniumchloride/ $\text{NH}_4\text{Cl}$ /water system will be discussed and the methods of measurements described.

### Incorporation and Transport of Solutes in Lipid Bilayers

E. SACKMANN\* and D. RÜPPEL

Abteilung für Biophysik, Universität Ulm, D-7900 Ulm, Oberer Eselsberg, F.R.G.

Lipid bilayers are excellent solvents for amphiphatic, hydrophobic and even charged molecules. This property is essential for the role of the lipid bilayer as fundamental building unit of biological membranes where a large class of molecules ranging from small substrates to large enzyme complexes must be organized in well defined structures. This property is closely related to the liquid-crystalline structure and the two-dimensionality of the bilayer. The molecular organization of bilayers and mono-

†Research supported under NSF Grant #DMR-7907789.