

wave functions of all particles involved are mutually disturbed. Therefore, the interaction energy is not pair-additive. Accordingly, we attempted to evaluate the contribution of the non-additivity to the pair correlation function and thermodynamic functions of a fluid in the simplest case of triplet interactions.

The Hamiltonian is as follows:

$$H = \frac{1}{2} \sum_{i,j} \psi_{ij} + \frac{1}{3!} \sum_{i,j,k} \psi_{ijk}^*$$

where ψ_{ij} is the pair (undisturbed by the presence of other particles) potential, ψ_{ijk}^* is the non-additive part (perturbation) of the triple interaction potential. The solution of the problem was achieved in the framework of the BBGKY method and using the functional expansion of the Hamiltonian in non-additive three-particle potentials for diluted solutions both neutral and charged particles.

Theoretical Study on Solvation of a Neutral Molecule: N₂. Two Different Approaches

P. F. DEMONTIS, R. ERCOLI, G. B. SUFFRITTI and A. GAMBA

Istituto di Chimica Fisica dell'Università di Sassari, Italy

A wide investigation has been performed in view of a theoretical study of the solvation of neutral molecules. N₂ as a test molecule and HF, H₂O, NH₃, CH₄ as solvents have been chosen.

In the former approach, the contour maps of interaction energy between N₂ and one solvent molecule have been evaluated, with fast semiempirical methods (Extended Hückel and CNDO), and compared with each other, in order to find a reliable but not too expansive calculation method. For N₂–H₂O interactions, a map has been computed also *ab initio*. The maps are shown, and qualitative and quantitative differences are discussed.

The latter approach rests on building up clusters with one molecule of N₂ surrounded by a number (2 to 8) of solvent molecules and finding the structure of such clusters by energy minimization.

From the reported structures it results that they are determined mainly by steric factors, so that clusters optimized by means of different methods are quite similar.

Solvent Extrusion: A Pathway from Solvent Separated Ion Pairs to Tight Ion Pairs

BERNARD BIGOT

Laboratoire de Chimie Organique Théorique, Université Pierre et Marie Curie, 4 Place Jussieu, Paris Cedex 05, 75230, France

and LIONEL SALEM

Laboratoire de Chimie Théorique, Université d'Orsay, Orsay 91405, France

A simple model of two oppositely charged ions, embedded in a sea of hard sphere solvent dipoles (whose centers are fixed on a lattice) is implemented to study the recombination of ions in solution. The first step is the passage from 'solvent separated' ion pair to 'intimate' or 'tight' ion pair. A preliminary two dimensional calculation (restricted to purely electrostatic terms) of the potential energy surface for such a recombination reveals two minima, as required by the experimental observations of Winstein and others.

References

S. Winstein, P. E. Klinedinst Jr. and G. C. Robinson, *J. Am. Chem. Soc.*, **83**, 885 (1961), and earlier papers cited therein; S. Winstein, P. Baker and S. G. Smith, *ibid.*, **86**, 2072 (1964); A. F. Diaz and S. Winstein, *ibid.*, **88**, 1318 (1966).

I.R. Experimental Studies of Dynamical Aspects of Water Structure: Effects of H–D Substitution and of Small Solutes

M. B. PALMA-VITTORELLI, G. ANDALORO, P. CHIRICÒ
Gruppo Nazionale Struttura della Materia, 90123 Palermo, Italy

G. GUZZIO and M. LEONE

Istituto di Fisica dell'Università di Palermo, 90123 Palermo, Italy

Results on i.r. absorption spectra of H₂O, D₂O and electrolytic solutions and their thermal modifications are presented. From the data, some parameters can be obtained which are suitable for a description of dynamical aspects of water structure. Solute induced changes of these parameters provide a quantitative, although phenomenologic description of the water–solute interaction.