

Aggregation of Tri-n-Octylammonium Chloride (TOAHCl), Tri-n-Octylammonium Bromide (TOAHBr) and Tri-n-Octylammonium Iodide (TOAHI) in Benzene Solutions

CVETO KLOFUTAR* and SPELA PALJK

J. Stefan Institute, University of Ljubljana, 61000 Ljubljana, Yugoslavia

In non-polar solvents tertiary n-alkylammonium salts exist as hydrogen bonded ion-pairs, which with increasing concentration aggregate to quadrupoles and higher oligomers, while at highest dilution ($<10^{-5}$ mol kg $^{-1}$) species in the form of positive and negative triple ions can be produced. These species may be further solvated. The extent and degree of association depend on the magnitude of the dipole moment, as well as the size and shape of the solute molecules and, additionally, on the nature of the solvent [1, 2].

The aggregation of the investigated solutes in benzene solutions is studied by cryoscopic method over the concentration range 0.05–0.25 mol kg $^{-1}$. We show, that the non-ideal behaviour of these systems as a consequence of a series of association equilibria, due to the dipole–dipole interactions of simple ion-pairs, $\text{IBHX} \rightleftharpoons (\text{BHX})_1$, with the self-association constant β_1 , can be described with two independent parameters: the dimerization constant β_2 and constant K , which are related to the self-association constant via the relation

$$\beta_1 = \beta_2 \frac{K^{1-2}}{1-1}, \quad (l \geq 2)$$

and to the stepwise association constant through

$$K_1 = \frac{1-2}{1-1} K, \quad (l \geq 3).$$

From the hypothesis used [3], it follows, that the first stepwise associations occur less readily than the rest. The values of $\beta_2 \times 10^{-3} = 5.5$ (TOAHCl); 3.0 (TOAHBr); 2.5 (TOAHI), $K \times 10^{-2} = 1.75$ (TOAHCl); 2.25 (TOAHBr); 3.00 (TOAHI) and free monomer concentration $n_{10}^* \times 10^3 = 1.0$ (TOAHCl); 1.1 (TOAHBr); 1.1 (TOAHI) are determined by curve-fitting method [3, 4]. The results indicate, that with increasing size of anion of tri-n-octylammonium halides in benzene solutions the dimerization decreases, while the presence of higher oligomers increases.

References

- 1 C. A. Kraus, *J. Phys. Chem.*, **60**, 129 (1956).
- 2 C. Klofutar and Š. Paljk, *J. Chem. Soc. Faraday Trans. I*, **75**, 825 (1979).

- 3 F. J. C. Rossotti and H. Rossotti, *J. Phys. Chem.*, **65**, 1376 (1961).
- 4 F. J. C. Rossotti and H. Rossotti, *J. Phys. Chem.*, **63**, 1041 (1959).

Interactions in the Systems $\text{Et}_3\text{N}/\text{AgNO}_3/\text{C}_6\text{H}_6$ and $\text{Et}_3\text{N}/\text{AgNO}_3/\text{CH}_3\text{COCH}_3$. Their Effect on the Catalysed Quaternization of Et_3N with EtI

ESTER F. G. BARBOSA and ISABEL M. S. LAMPREIA

Centor de Electroquímica e Cinética da Universidade de Lisboa, Faculdade de Ciências, Rua da Escola Politécnica, 58, 1200 Lisboa Codex, Portugal

A strong catalytic effect having been observed due to the presence of AgNO_3 in a Menschutkin reaction mixture in benzene [1] and acetone [2], it has been considered of interest to study more deeply this effect. Studies of interactions in the systems $\text{Et}_3\text{N}/\text{AgNO}_3/\text{solvent}$ are required to understand the mechanism and rate law of the proposed catalysed reaction.

The studies started with benzene and acetone as solvents. Density measurements in the binary mixtures $\text{Et}_3\text{N}/\text{C}_6\text{H}_6$ and $\text{Et}_3\text{N}/\text{CH}_3\text{COCH}_3$ as well as in the ternary mixtures $\text{Et}_3\text{N}/\text{AgNO}_3/\text{C}_6\text{H}_6$ and $\text{Et}_3\text{N}/\text{AgNO}_3/\text{CH}_3\text{COCH}_3$ have been undertaken in this laboratory using a Digital Anton Paar DMA O2D Density Meter. Computation of partial molal volumes were made.

Discussion of the foregoing results together with infrared spectroscopic information and the analysis of data found in the literature [3–7] led us to propose structural models for the investigated mixtures and to make correlations with the experimental rates measured for the catalysed reactions.

References

- 1 E. F. G. Barbosa and M. Spiro, *J. Chem. Soc., Chem. Comm.*, 423 (1977).
- 2 This Laboratory.
- 3 B. G. Cox, A. J. Parker and W. E. Waghorne, *J. Phys. Chem.*, **78**, 1731 (1974).
- 4 J. Edward, P. Farrell and F. Shahidi, *J. Chem. Soc., Faraday I*, **73**, 705 (1977).
- 5 F. Shahidi, P. Farrell and J. Edward, *J. Chem. Soc., Faraday I*, **73** (1977).
- 6 S. Cabani, G. Conti and L. Lepori, *J. Phys. Chem.*, **78**, 1030 (1974).
- 7 Š. Paljk, C. Klofutar and M. Zumer, *J. Inorg. Nucl. Chem.*, **38**, 293 (1976).