

The role of the keto acid and metal ion in this pathway, are significant and it has been shown that oxygen uptake is reduced by excess keto acid and varies, as expected, with the choice of metal ion. With copper further oxidation occurs yielding other carbon fragments including carbon dioxide and acetic acid.

The reaction is also shown to occur with dipeptides, but to be relatively inefficient for tripeptides.

- 1 D. A. Phipps, *J. Mol. Catalysis*, 5, 81–107 (1979).
- 2 R. D. Gillard, P. O'Brien, P. R. Norman and D. A. Phipps, *J. Chem. Soc. (Dalton)*, 1988 (1977).
- 3 A. Dempsey and D. A. Phipps, *Inorg. Chim. Acta*, 36, 1425 (1979).

U3

The Correlation between the Activity of Urease Immobilized to Anodized Sheet Aluminium and the Anodizing Conditions

P. GRUNWALD

Institut für Physikalische Chemie, Universität Hamburg, Laufgraben 24, D-2000 Hamburg 13, F.R.G.

The fixation of proteins to water insoluble materials has become an important field in chemistry and related disciplines [1]. Preference is often given to inorganic carriers because of their resistance to bacterial degradation. Moreover, the investigation of the interaction between proteins and nonbiological surfaces is of great practical interest with respect to the increasing use of prosthetic materials in the body [2].

Recently, we could show with the enzyme urease as an example that anodized sheet aluminium is a suitable carrier material for enzyme immobilisation [3, 4]. Further, the anodizing process leads to an adsorbent with a homogeneous surface, which is required for the interpretation of adsorption experiments. In this paper the influence of the different anodizing parameters on the activity of urease adsorbed to anodized sheet aluminium is described. The results are discussed with regard to the surface structure that has been studied by scanning electron micrographs.

The anodizing of aluminium was carried out in sulfuric acid as electrolyte. The rolled Al-sheets were placed between two cathodes of lustrous carbon. A dc power supply allowed anodizing at constant voltage and constant current. The investigated parameters were: the concentration of sulfuric acid (c_a), the applied voltage U and current density (i), the anodizing temperature (T_a), the anodizing time (t_a), and the contact time (t_{ad}) between anodized sheet

aluminium and the enzyme solution. We found that optimum activities are obtained if the anodizing conditions are: $c_a = 26$ wt%, $t_a = 50$ min, $T_a = 308$ K, and $i = 85$ mA/cm². The connection between the single anodizing parameters and the activity of the adsorbed enzyme is as follows: in the starting experiment with $U = 18$ Volt, $t_a = 30$ min, $t_{ad} = 40$ min, $T_a = 298$ K and c_a between 2 and 40 wt% the best results were achieved with $c_a = 26$ wt%; the corresponding i -value was 50 mA/cm². Higher acid concentrations lower the current density at which optimum activities are obtained. The activity itself differs only negligibly. At t_a -values less than 20 min the activity of the immobilized enzyme is low. For $30 \text{ min} < t_a < 50 \text{ min}$ the activity is raised by a factor of 1.5 to 2, dependent on c_a . If T_a is 308 K instead of 298 K the activity is three times as high as in the starting experiment. A nearly tenfold activity results if i is raised up to 80 mA/cm², and the activity increases again by a factor of 2 if t_{ad} is extended by 100 min. The importance of the contact time t_{ad} for the activity of the adsorbed enzyme distinctly increases for $i > 50$ mA/cm² as a consequence of the increasing porosity of the anodized sheet aluminium. The complex influence of the anodizing parameters on the surface structure and its correlation to the activity of the adsorbed enzyme is conclusively interpretable by means of scanning electron micrographs.

- 1 K. Mosbach, *FEBS Lett.*, 62 (suppl.), E80 (1976).
- 2 M. Schaldach, *Klin. Wochenschrift*, 53, 1029 (1975).
- 3 P. Grunwald, W. Gunsser, K. P. Pfaff, R. Krause and K. Lutz, *Z. Naturforsch.*, 35c, 819 (1980).
- 4 P. Grunwald, H. Kleine and W. Gunsser, 28th IUPAC Congress, BR 132, Vancouver, Canada (1981).

U4

Stereochemistry of Pyridoxal–Amino Acid Model Systems

LUIGI CASELLA and MICHELE GULLOTTI

Dipartimento di Chimica Inorganica e Metallorganica, Centro CNR, Via Venezian 21, 20133 Milan, Italy

Pyridoxal phosphate is the essential cofactor for most of the enzymic reactions undergone by amino acids during metabolism [1]. All these reactions proceed through the formation of a Schiff base, such as I, followed by cleavage of one of the bonds to the amino acid α -carbon atom. The nature of the substrate and the conformation of the apoenzyme dictate which of the bonds to C_α is broken and control the further course of the reaction. The stereospecificity of the reaction depends on the ability of the dissymmetric apoenzyme to bind the