

Synthesis and Reactions of $[M(CO)_4(Ph_2PSiMe_3)X]$ Complexes ($M = Mn, Re$; $X = Halogen$)

PETER J. MANNING, LOUIS K. PETERSON*, FUMIO WADA and RANDI S. DHAMI

Department of Chemistry, Simon Fraser University, Burnaby, B. C., V5A 1S6, Canada

Received May 23, 1985

Abstract

The silylphosphine ligand Ph_2PSiMe_3 reacts readily with a slurry of $[Re(CO)_5X]$ ($X = Cl, Br$) in polar and in non-polar solvents to yield soluble *cis*- $[Re(CO)_4(Ph_2PSiMe_3)X]$ (**Ia**, $X = Cl$; **Ib**, $X = Br$) via CO substitution. Compound **I** is readily hydrolyzed by water or silica gel to *cis*- $[Re(CO)_4(Ph_2PH)X]$. Compound **Ib** reacts with $[Re(CO)_5Br]$ to yield $[Re_2(CO)_8(\mu-PPh_2)(\mu-Br)]$ (**II**), and with $[Mn(CO)_5Br]$ to yield $[MnRe(CO)_8(\mu-PPh_2)(\mu-Br)]$ (**III**).

The reaction of Ph_2PSiMe_3 with $[Mn(CO)_5X]$ ($X = Cl, Br, I$) is highly dependent upon reaction conditions. In polar and in non-polar solvents, an excess of ligand gives mainly *cis*- $[Mn(CO)_4(Ph_2PSiMe_3)X]$ (**IVa**, $X = Cl$; **IVb**, $X = Br$; **IVc**, $X = I$). With ligand: $[Mn(CO)_5X]$ reacting ratios in the range 0.5–1.0:1, the products from the three respective halomanganese complexes in THF were: (a) mainly $[Mn_2(CO)_8(\mu-PPh_2)(\mu-Cl)]$ (**Va**); (b) both $[Mn(CO)_4(Ph_2PSiMe_3)Br]$ and $[Mn_2(CO)_8(\mu-PPh_2)(\mu-Br)]$ (**Vb**); and (c) exclusively $[Mn(CO)_4(Ph_2PSiMe_3)I]$. The compounds **IVa–c** are stable in solution at ambient temperatures and are readily hydrolyzed by water or methanol to $[Mn(CO)_4(Ph_2PH)X]$. Compound **IVb** reacts at room temperature with $[Mn(CO)_5Cl]$ to yield only $[Mn_2(CO)_8(\mu-PPh_2)(\mu-Br)]$ (**Vb**); compound **IVc** reacts in hot toluene with $[Mn(CO)_5Cl]$ to yield mainly $[Mn_2(CO)_8(\mu-PPh_2)(\mu-I)]$ (**Vc**), together with a small amount of the chloro-bridged analog.

The dinuclear species **II**, **III** and **Va–c** appear to be formed mainly via an intermolecular elimination of Me_3SiX from the appropriate $[M(CO)_4(Ph_2PSiMe_3)X]$ and metalpentacarbonylhalide (chloride or bromide) complexes.

Introduction

The potential utility of phosphido-bridged polynuclear species as possible multi-site catalysts is an area of substantial current interest [1]. On the one hand, bridging ligands may inhibit cluster fragmenta-

tion [1c, 1e–h], or, alternatively, ligand lability may provide routes to active site formation [1a, g, i, j] and to the formation of novel dynamic polynuclear systems [1b].

Several procedures for the synthesis of phosphido-bridged complexes have been described [1j, 2]. In this paper, we report our studies of the reactions of Ph_2PSiMe_3 with $[M(CO)_5X]$ ($M = Mn, Re$; $X = halogen$), and the general properties of the new complexes $[M(CO)_4(Ph_2PSiMe_3)X]$, $[M_2(CO)_8(\mu-PPh_2)(\mu-X)]$ and $[MnRe(CO)_8(\mu-PPh_2)(\mu-Br)]$.

Discussion

The reaction of Ph_2PSiMe_3 with $[M(CO)_5X]$ ($M = Mn, Re$; $X = halogen$) in refluxing dimethoxyethane (DME) was reported to yield $[M(CO)_4PPh_2]_2$ ($M = Mn, Re$) and $[Mn(CO)_3PPh_2]_3$ [3a]. None of the possible intermediates – $[Mn(CO)_4(Ph_2PSiMe_3)X]$, $[M(CO)_4,5(PPh_2)]$, or $[(X)(CO)_4M(\mu-PPh_2)M(CO)_4(Ph_2PSiMe_3)]$ – were detected by the earlier workers. Thus, the details of the reaction pathway(s) leading to the claimed phosphido-bridged products have remained a mystery. Reactions between $P(EMe_3)_3$ ($E = Ge, Sn$) and $[Mn(CO)_5X]$ have also been reported to yield dinuclear $[Mn(CO)_4P(EMe_3)_2]_2$ [3b].

In our studies of the reaction of Ph_2PSiMe_3 with slurries of $[Re(CO)_5X]$ ($X = Cl, Br$) in hexane, benzene, toluene, THF and DME at ambient temperatures, we observed the formation of *cis*- $[Re(CO)_4(Ph_2PSiMe_3)X]$ (**Ia**, $X = Cl$; **Ib**, $X = Br$). The free ligand and compounds **Ia, b** are extremely sensitive to moisture, so that the pale yellow solid products obtained upon removal of solvent were invariably contaminated with $[Re(CO)_4(Ph_2PH)X]$ [4]. When **Ia** and **Ib** were chromatographed on silica gel, with benzene as the eluant, the latter secondary phosphine complexes were obtained in high yield. Compounds **Ia, b** were characterized by IR, 1H and ^{31}P NMR spectroscopy (see 'Experimental'). Compounds **Ia, b** are stable in solution at ambient temperatures but, at elevated temperatures, highly insoluble products were formed. These products showed infrared bands in the

*Author to whom correspondence should be addressed.

$\nu(\text{CO})$ region that have been ascribed in the literature to the dimer $[\text{Re}(\text{CO})_4\text{PPh}_2]_2$ [3a]; on the basis of later evidence, it appears that this product may consist of an equilibrium mixture of polynuclear species [5].

When combined with $[\text{Re}(\text{CO})_5\text{Br}]$ and with $[\text{Mn}(\text{CO})_5\text{Br}]$, $[\text{Re}(\text{CO})_4(\text{Ph}_2\text{PSiMe}_3)\text{Br}]$ gave the dinuclear species $[\text{Re}_2(\text{CO})_8(\mu\text{-PPh}_2)(\mu\text{-Br})]$ (**II**) and $[\text{MnRe}(\text{CO})_8(\mu\text{-PPh}_2)(\mu\text{-Br})]$ (**III**), respectively. Compounds **II** and **III** were characterized by IR, ^{31}P and mass spectroscopy. The IR spectra of compound **II** and the related $[\text{Re}_2(\text{CO})_8(\mu\text{-P}(\text{CF}_3)_2)(\mu\text{-X})]$ [6] compounds show five major strong $\nu(\text{CO})$ bands, in a pattern indicative of their structural similarities, viz., equivalent C_1 'M(CO) $_4$ ' units linked by halogen and phosphido bridges. The mixed metal complex $[\text{MnRe}(\text{CO})_8(\mu\text{-PPh}_2)(\mu\text{-Br})]$ (**III**), containing inequivalent 'M(CO) $_4$ ' units, showed only six resolvable $\nu(\text{CO})$ bands. The IR spectra of $[\text{Re}(\text{CO})_5\text{X}_2]$ (X = Cl, Br, I) [7], and $[\text{Re}_2(\text{CO})_8(\text{PPh}_2)_2]$ [3a] show four major bands. The ^{31}P chemical shifts of **II** and **III** are typical for phosphido-bridged structures that do not contain a metal-metal bond [8].

The reactions of $\text{Ph}_2\text{PSiMe}_3$ with $[\text{Mn}(\text{CO})_5\text{X}]$ (X = Cl, Br, I) were monitored by IR ($\nu(\text{CO})$ region) and NMR (^1H and ^{31}P) spectroscopy, using various ligand-to-complex molar ratios in the solvents hexane, d_6 -benzene, toluene, THF and DME. In polar solvents, rapid evolution of gas bubbles was observed and deep red solutions were formed. The toluene solutions reacted more slowly and showed variable induction periods before CO evolution was apparent. The hexane solutions reacted even more slowly at room temperature, being substantially incomplete even after two days. Under comparable conditions in hexane solution, it was found that the reaction of $\text{Ph}_2\text{PSiMe}_3$ with $[\text{Mn}(\text{CO})_5\text{Cl}]$ was much slower than the reaction of Ph_3P . The more rapid reaction of $\text{Ph}_2\text{PSiMe}_3$ with $[\text{Mn}(\text{CO})_5\text{X}]$ in polar solvents is noteworthy and is in contrast to reports that tertiary phosphines in general react more rapidly in non-polar media [5, 9]. In addition, the rates of substitution of CO by $\text{Ph}_2\text{PSiMe}_3$ vary in the order $\text{Cl} > \text{Br} > \text{I}$ in polar solvents, similar to relative rates of substitution by tertiary phosphines [9].

As noted above, the complete removal of Ph_2PH from $\text{Ph}_2\text{PSiMe}_3$ was difficult to achieve (but usually less than 5%, based on ^{31}P NMR signal integrals). It was found, however, that the secondary phosphine reacted more rapidly in hexane with $[\text{Mn}(\text{CO})_5\text{X}]$ than $\text{Ph}_2\text{PSiMe}_3$, yielding $[\text{Mn}(\text{CO})_4(\text{Ph}_2\text{PH})\text{X}]$ [4, 10] and, occasionally, $[\text{Mn}(\text{CO})_3(\text{Ph}_2\text{PH})_2\text{X}]$ [10].

The products in solution resulting from the silylphosphine- $[\text{Mn}(\text{CO})_5\text{X}]$ reactions were identified as *cis*- $[\text{Mn}(\text{CO})_4(\text{Ph}_2\text{PSiMe}_3)\text{X}]$ (**IVa**, X = Cl; **IVb**, X = Br; **IVc**, X = I), $[\text{Mn}_2(\text{CO})_8(\mu\text{-PPh}_2)(\mu\text{-X})]$ (**Va**, X = Cl; **Vb**, X = Br; the iodo analog **Vc** was not observed) and traces of $[\text{Mn}_2(\text{CO})_8(\mu\text{-H})(\mu\text{-PPh}_2)]$

[2p, 11]. Compounds **IVa-c** were identified in solution by IR and NMR (^1H and ^{31}P) spectroscopy and by their essentially quantitative hydrolysis to $[\text{Mn}(\text{CO})_4(\text{Ph}_2\text{PH})\text{X}]$ [4, 10]. Compound **Vc**, $[\text{Mn}_2(\text{CO})_8(\mu\text{-PPh}_2)(\mu\text{-I})]$, was prepared in a separate experiment, via the reaction of **IVc** with $[\text{Mn}(\text{CO})_5\text{-Cl}]$. The complexes **Va-c** were characterized by IR, NMR and mass spectroscopy. Their IR spectra ($\nu(\text{CO})$ region) showed a weak high energy band and four other strong bands, in a pattern very similar to those of $[\text{Mn}_2(\text{CO})_8(\mu\text{-P}(\text{CF}_3)_2)(\mu\text{-X})]$ [12] and the ($\mu\text{-X}$)-(μ -phosphido)-dirhenium analogs. The relative proportions of **IV** and **V** were found to vary, depending on nature of solvent, relative amount of ligand present, nature of halogen 'X' and time of reaction. In all cases, dark brown components which could not be eluted from silica gel chromatographic columns were also obtained.


The reaction of $[\text{Mn}(\text{CO})_5\text{Cl}]$ with excess ligand gave mainly compound **IVa** in polar and in non-polar solvents, whereas a deficiency of ligand gave rise to appreciable quantities of **Va** as well as **IVa**; the latter apparently reacts with $[\text{Mn}(\text{CO})_5\text{Cl}]$ to yield **Va**. The formation of **Va** was favored in polar solvents.

The reaction of $[\text{Mn}(\text{CO})_5\text{Br}]$ with equivalent and with excess amounts of $\text{Ph}_2\text{PSiMe}_3$ in hexane gave mainly compound **IVb**. In polar solvents, however, both **IVb** and **Vb** were observed in the early stages of the reaction, regardless of the molar ratios of reactants employed. In a separate study, it was shown that **IVb** reacts very slowly with $[\text{Mn}(\text{CO})_5\text{Br}]$, indicating that there is another pathway to compound **Vb** in the early stages of the reaction involving $\text{Ph}_2\text{PSiMe}_3$. Compound **Vb** was also prepared by the reaction of LiPPh_2 with $[\text{Mn}(\text{CO})_5\text{Br}]$ (1:2 molar ratio). The iodo system was different yet again, in that $[\text{Mn}(\text{CO})_5\text{I}]$ gave only **IVc**, regardless of solvent type or reactant ratios.


In one experiment, the photolytic preparation of $[\text{Mn}(\text{CO})_4(\text{Ph}_2\text{PSiMe}_3)\text{Cl}]$ from $\text{Mn}(\text{CO})_5\text{Cl}$ and $\text{Ph}_2\text{PSiMe}_3$ was attempted. The reaction was complex, however, giving a variety of products, including short-lived paramagnetic species. The photolytic studies were therefore discontinued.

The possible role of the $[\text{Mn}(\text{CO})_{4,3}(\text{Ph}_2\text{PH})_{1,2}\text{X}]$ impurities as intermediates in reaction sequences leading to phosphido-bridged species was examined. In separate studies, it was found that the above complexes do not thermally eliminate HX in solution, nor do they react significantly with $[\text{Mn}(\text{CO})_5\text{X}]$ or $[\text{Mn}(\text{CO})_4(\text{Ph}_2\text{PSiMe}_3)\text{X}]$. For comparison, it is reported that complexes of the type $[\text{M}(\text{CO})_n(\text{Ph}_2\text{PH})]$ (M = Fe, Ni) undergo dehydrohalogenation with various metal halide complexes, in the presence of diethylamine, to yield mono- μ -phosphido derivatives [13], while several $[\text{M}(\text{CO})_n(\text{Ph}_2\text{PH})_m]$ complexes polymerize via dehydrogenation [14, 15].

On the basis of the above studies, it is evident that Ph₂PSiMe₃ reacts with [Re(CO)₅X] to produce [Re(CO)₄(Ph₂PSiMe₃)X] in high yields, via displacement of carbon monoxide (eqn. (1)).



The reactions of Ph₂PSiMe₃ with [Mn(CO)₅X] are more complex than those of the rhenium analogues. The dominant primary process again appears to be displacement of CO, leading to [Mn(CO)₄(Ph₂PSiMe₃)X] (IVa–c) via eqn. (1). Because of the greater reactivity of the manganese derivatives, condensation products are readily formed. It appears that one route to such dinuclear species involves an initial nucleophilic attack by halide (chloride and bromide, but not iodide) at the silicon center of a coordinated trimethylsilyldiphenylphosphine ligand (eqn. (2)). Evidence supporting this conclusion



includes: (i) The reactions of [Mn(CO)₅Cl] with [Mn(CO)₄(Ph₂PSiMe₃)X] (X = Br, IVb; X = I, IVc) to yield Mn₂(CO)₈(μ-PPh₂)(μ-X) (X = Br, Vb; X = I, Vc), via the elimination of Me₃SiCl. (ii) The reactions of [Re(CO)₄(Ph₂PSiMe₃)Br] (Ib) with [Re(CO)₅Br] and with [Mn(CO)₅Br], at ambient temperature, to yield Re₂(CO)₈(μ-PPh₂)(μ-Br) (II) and MnRe(CO)₈(μ-PPh₂)(μ-Br) (III), respectively. (iii) Attempts to trap the possible 'Re(CO)₄PPh₂' intermediate [3a] as a saturated complex [Re(CO)₄(L)(PPh₂)], by carrying out reactions in the presence of an excess of various ligands L, were unsuccessful. (iv) Hydrolysis of the coordinated Ph₂PSiMe₃ ligand occurs readily, giving complexes of Ph₂PH. In addition, the compounds [M(CO)₄(Ph₂PSiMe₃)X] are relatively stable in solution at ambient temperature, indicating that the intramolecular elimination of Me₃SiX is not a favourable reaction pathway.

Contrasting with the mechanism proposed above, it is suggested in the literature [6] that the reaction of [M(CO)₅X] (M = Mn, Re; X = halogen) with P₂(CF₃)₄ or As₂(CF₃)₄ proceeds first via metal–halogen bond cleavage rather than CO substitution, giving a terminal phosphido or arsenido complex [M(CO)₅E(CF₃)₂] (E = P, As). The latter then reacts further with [M(CO)₅X] to generate [M₂(CO)₈(μ-E(CF₃)₂(μ-X))] species. The participation of [Mn(CO)₅PPh₂] in the formation of dinuclear species, in competition with halide attack at silicon (eqn. (2)), cannot be ruled out completely in our work. Such terminal phosphido species are known to dimerize or polymerize readily, unless the basic center is modified by the presence of highly electro-negative substituents [16].

Experimental

Reactions were carried out under a purified nitrogen atmosphere, using conventional Schlenk and vacuum equipment and a Vacuum Atmospheres Model HE493 drybox, as appropriate. Solvents were dried and distilled under nitrogen before use. The halocomplexes [M(CO)₅X] (M = Mn, Re; X = Cl, Br, I) were prepared from [Mn₂(CO)₁₀] or [Re₂(CO)₁₀] by reaction with halogen, according to published procedures [17, 18]. The ligand Ph₂PSiMe₃ was prepared as described in the literature [19] but, despite careful distillation, some Ph₂PH impurity was invariably present. The purity was checked by ³¹P NMR spectroscopy.

Spectra were obtained using the Perkin-Elmer 983 infrared spectrometer, the Hewlett Packard 5985 mass spectrometer and the Bruker WM400 and Varian EM360 NMR spectrometers. The phosphorus coordination chemical shift Δ(δ³¹P) was evaluated as δ³¹P (free ligand)–δ³¹P (complex). In a few cases, evolved carbon monoxide was measured by means of a Sprengle pump and its purity checked by gas chromatography.

Unless otherwise noted, reactions were performed on 0.3–0.6 mmol scales. The following selected descriptions are typical of the procedures used.

(a) [Re(CO)₅X] + Ph₂PSiMe₃ in Toluene and Benzene

(i) A mixture (1:1 molar ratio) of [Re(CO)₅X] (X = Cl, Br) and Ph₂PSiMe₃ was heated until CO evolution was complete, giving a pale yellow solution. An IR spectrum of the solution showed ν(CO) bands attributable to *cis*-[Re(CO)₄(Ph₂PSiMe₃)X] (Ia, X = Cl; Ib, X = Br). The solvent was removed, giving compounds Ia, b as a pale yellow solid in high yield, contaminated with *cis*-[Re(CO)₄(Ph₂PH)X].

(ii) Reactions were also carried out in C₆D₆, in order to obtain ¹H and ³¹P NMR spectra.

(iii) A mixture (2:1 molar ratio) of [Re(CO)₅Br] and Ph₂PSiMe₃ in toluene was heated to reflux for 3 h. The IR spectrum showed that all of the [Re(CO)₅-Br] had been consumed. The solvent was removed under vacuum and the residue was dissolved in CDCl₃. The ³¹P NMR spectrum showed only two phosphorus-containing compounds, [Re(CO)₄(Ph₂PH)Br] and [Re₂(CO)₈(μ-PPh₂)(μ-Br)] (II) (50–60%). These products were separated and purified by TLC.

(b) [Re(CO)₅Br] + Ph₂PSiMe₃ in 1,2-dimethoxyethane (DME)

In this experiment, a reaction vessel consisting of two compartments separated by 'Rotaflo' valves was used so that evolved carbon monoxide could be measured. A solution of Ph₂PSiMe₃ in toluene was placed in one compartment and solid [Re(CO)₅Br]

(1 molar equivalent) was placed in the other compartment. The apparatus was evacuated and degassed. The reagents were mixed and allowed to react for several hours. The vessel was cooled in liquid N₂, and the evolved carbon monoxide was collected by means of a Sprengle pump. 96% of the expected amount was obtained. Its purity was checked by gas chromatography. The initial IR spectrum of the DME solution was essentially similar to that of [Re(CO)₄(Ph₂PSiMe₃)Br] (**Ib**) in benzene. After several hours, changes in the IR spectrum were noted and a white precipitate, identified by IR and mass spectroscopy as [Re(CO)₄PPh₂]₂ [**3a**], was obtained.

(c) *Conversion of cis-[Re(CO)₄(Ph₂PSiMe₃)Br] (**Ib**) into cis-[Re(CO)₄(Ph₂PH)Br]*

A benzene solution of compound **Ib** was placed on a silica gel column and eluted with benzene. The solvent was removed under vacuum and the residue, *cis*-[Re(CO)₄(Ph₂PH)Br], was recrystallized from hexane. Yield, 85%, based on [Re(CO)₅Br] taken.

Characterization of Rhenium Complexes

[Re(CO)₄(Ph₂PH)Br], $\nu(\text{CO})$ in hexane, 2108 m, 2023 s, 2012 vs, 1944 m; δ ³¹P (in CDCl₃), -11.7d, $J(\text{PH}) = 383$ Hz; $\Delta(\delta$ ³¹P) = -28 ppm; lit. [4], δ ¹H (PH), 7.1 ppm, $J(\text{PH}) = 382$ Hz. The mass spectrum showed the parent ion multiplet at *m/e* 562, 564 and 566 and stepwise loss of four CO groups. Melting point 127.5–129 °C, lit. [4] 128 °C. *Anal. calc.* for C₁₆H₁₁BrO₄PRE: C, 34.04; H, 1.95. Found: C, 34.2; H, 1.95%. [Re(CO)₄(Ph₂PSiMe₃)Br], **Ib**, $\nu(\text{CO})$ in hexane, 2107 s, 2010 s, 2002 m, 1946 s; δ ¹H (PSiMe₃), 0.24 d, $J(\text{PH}) = 5.7$ Hz; δ ³¹P, -56.3; $\Delta(\delta$ ³¹P) = 0.4 ppm [Re₂(CO)₈(μ -PPh₂)(μ -Br)] (**II**), $\nu(\text{CO})$ in hexane, 2111 w, 2095 m, 2018 vs, 2010 m, 2003 s, 1951 s; δ ³¹P (in CDCl₃), -93.3.

(d) *[Mn(CO)₅Cl] + Ph₂PSiMe₃ in Benzene*

In an NMR tube, a slurry of [Mn(CO)₅Cl] in C₆D₆ and Ph₂PSiMe₃ was photolyzed for 7 min, giving an orange solution. The evolved gas pressure was released. The first ¹H NMR spectra showed no fine structure, due to the presence of paramagnetic species. After several minutes, the ¹H NMR spectrum showed a strong methyl singlet (due to Me₃SiCl) and two weaker doublets (due to free Ph₂PSiMe₃ and [Mn(CO)₄(Ph₂PSiMe₃)Cl] (**IVa**). The solution was allowed to stand for several days. The products were separated by silica gel TLC, giving many bands from which [Mn₂(CO)₈(μ -PPh₂)(μ -Cl)] (**Va**) and [Mn₂(CO)₈(μ -H)(μ -PPh₂)] (trace) were isolated.

(e) *[Mn(CO)₅Cl] + Ph₂PSiMe₃ in THF*

(i) The ligand Ph₂PSiMe₃ (0.68 mmol) was syringed through a rubber septum into a solution of [Mn(CO)₅Cl] (0.68 mmol) in THF in a 10 mm NMR tube. Almost immediately, carbon monoxide was

evolved and the solution turned red. Excess pressure was released via the septum. After \approx 45 min, the ³¹P {H} NMR spectrum was observed. Many signals were present, the main ones being due to [Mn(CO)₄(Ph₂PSiMe₃)Cl] (**IVa**) (relative intensity = 1); [Mn₂(CO)₈(μ -PPh₂)(μ -Cl)] (**Va**) (intensity \approx 1.7); [Mn(CO)₄(Ph₂PH)Cl] (intensity \approx 2); and [Mn(CO)₃(Ph₂PH)₂Cl] (intensity \approx 1.7). Upon exposure to air, the signal due to **IVa** was lost. Chromatographic separation (using 4:1 hexane/chloroform eluant) yielded **Va** (0.05 mmol) and [Mn(CO)₃(Ph₂PH)₂Cl].

(ii) The above reaction was repeated, using a 2:1 molar ratio of ligand to manganese complex. The ³¹P NMR spectrum showed the presence of [Mn(CO)₄(Ph₂PSiMe₃)Cl] (**IVa**) and *fac*-[Mn(CO)₃(Ph₂PH)₂Cl], but only trace amounts of [Mn₂(CO)₈(μ -PPh₂)(μ -Cl)] (**Va**) and [Mn(CO)₄(Ph₂PH)Cl]. The secondary phosphine Ph₂PH was also present. The solution was stored under anhydrous conditions for 4 weeks. The only major change in the ³¹P NMR spectrum was a decrease in the intensity of the signal due to *fac*-[Mn(CO)₃(Ph₂PH)₂Cl], coupled with the appearance of signals due to other complexes of Ph₂PH. The *fac*-to-*mer-trans*-isomerization of [Mn(CO)₃(Ph₂PH)X] is a known process [10]. The components were separated by TLC. [Mn₂(CO)₈(μ -PPh₂)(μ -H)] (trace) and *fac*-[Mn(CO)₃(Ph₂PH)₂Cl] were isolated. The IR $\nu(\text{CO})$ spectrum of the latter was in agreement with literature data [10].

(iii) A 2:1 reaction mixture of [Mn(CO)₅Cl] and Ph₂PSiMe₃ in THF was monitored by IR spectroscopy over a 48 h period. After 75 min, $\nu(\text{CO})$ bands due to [Mn(CO)₅Cl], [Mn(CO)₄(Ph₂PSiMe₃)Cl] (**IVa**) and [Mn₂(CO)₈(μ -PPh₂)(μ -Cl)] (**Va**) were observed. The bands due to the first two components slowly disappeared over the 48 h period. The solvent was removed to yield compound **Va**, which was recrystallized from toluene, as an orange solid.

(f) *[Mn(CO)₅Br] + Ph₂PSiMe₃ in Hexane*

The ligand Ph₂PSiMe₃ was injected through a rubber septum into a refluxing solution of [Mn(CO)₅Br] in hexane (1:1 reacting ratio). An immediate evolution of gas bubbles occurred. The IR spectrum showed mainly unreacted [Mn(CO)₅Br] and some [Mn(CO)₄(Ph₂PH)Br]. Further heating for 30 min produced [Mn(CO)₄(Ph₂PSiMe₃)Br] (**IVb**), and a trace of [Mn₂(CO)₈(μ -PPh₂)(μ -Br)] (**Vb**). The mixture was hydrolyzed and its ³¹P NMR spectrum taken, showing the presence of [Mn(CO)₄(Ph₂PH)Br] and a trace of **Vb** as the only phosphorus-containing compounds.

(g) *[Mn(CO)₅Br] + Ph₂PSiMe₃ in DME*

The silylphosphine (2.43 mmol) was added to a warm solution of [Mn(CO)₅Br], (3.36 mmol) in DME (60 cm³), to give an orange solution. After 10 min, the solution was red. It was cooled and methanol was

added to hydrolyze the silylphosphine. The solvent was removed under vacuum, giving an orange residue which was separated by column chromatography (eluant, 1:1 toluene/cyclohexane) into orange [Mn₂(CO)₈(μ-PPh₂)(μ-Br)] (**Vb**) (0.2 mmol), [Mn(CO)₄(Ph₂PH)Br] (1.26 mmol) and a minor amount of yellow [Mn(CO)₃(Ph₂PH)₂Br].

(h) Reaction of [Mn(CO)₄(Ph₂PSiMe₃)Br] (**IVb**) with [Mn(CO)₅Cl]

Compound **IVb** was prepared *in situ* in THF, using a 1:1 molar ratio of Ph₂PSiMe₃ and [Mn(CO)₅Br]. A ³¹P NMR spectrum of the solution showed the presence of **IVb**, [Mn(CO)₃(Ph₂PH)₂Br] and excess Ph₂PSiMe₃. The solution was transferred under N₂ to a flask containing [Mn(CO)₅Cl] in THF. IR spectroscopy showed the formation of [Mn₂(CO)₈(μ-PPh₂)(μ-Br)] (**Vb**), over a period of two days. This product was separated by TLC and characterized by IR, ³¹P NMR and mass spectroscopy. Yield, 15%, based on [Mn(CO)₅Br] taken.

(i) Reaction of [Mn(CO)₄(Ph₂PSiMe₃)I] (**IVc**) with [Mn(CO)₅Cl]

A toluene solution of [Mn(CO)₅I] (0.34 mmol) was allowed to react with Ph₂PSiMe₃ (1:1 molar ratio) for 1.5 h, forming [Mn(CO)₄(Ph₂PSiMe₃)I] (**IVc**) in solution. [Mn(CO)₅Cl] (0.26 mmol) was added, the solution was allowed to stand for 3 h at ambient temperatures, then heated to 80 °C for 10 min to complete the reaction. The solvent was removed under vacuum and the residue was taken up in CH₂Cl₂ (0.5 cm³) and separated by TLC (eluant 1:1, petrol/chloroform) into [Mn₂(CO)₁₀] (trace), [Mn₂(CO)₈(μ-PPh₂)(μ-I)] (0.09 mmol), [Mn(CO)₅Cl] (trace) and [Mn(CO)₄(Ph₂PH)I] (0.06 mmol). A small amount of CH₂Cl₂-insoluble material was found to be [Mn₂(CO)₈(μ-PPh₂)(μ-Cl)].

Characterization of Manganese Complexes

[Mn(CO)₄(Ph₂PSiMe₃)Cl], **IVa**, δ ³¹P in THF, -13.8, Δ (δ ³¹P) = -42 ppm. [Mn₂(CO)₈(μ-PPh₂)(μ-Cl)], **Va**, ν(CO) in cyclohexane, 2097 w, 2078 m, 2021 vs, 2008 s, 1956 s; δ ³¹P in CDCl₃, 1.6. The mass spectrum showed the parent ion at m/e = 524/526, together with fragments corresponding to the sequential loss of eight CO groups and the μ-Ph₂P group. [Mn(CO)₄(Ph₂PSiMe₃)Br], **IVb**, δ ³¹P in THF, -14.3, Δ (δ ³¹P) = -41.6 ppm. [Mn₂(CO)₈(μ-PPh₂)(μ-Br)], **Vb**, ν(CO) in cyclohexane, 2095 w, 2075 m, 2018 vs, 2006 s, 1959 s; δ ³¹P in CDCl₃, -10.8. The mass spectrum showed the parent ion doublet at m/e = 598, 600, corresponding to the presence of ^{79,81}Br isotopes, and the stepwise loss of eight CO groups and the μ-PPh₂ group. [Mn(CO)₄(Ph₂PSiMe₃)I], **IVc**, δ ³¹P in THF, -14.3, Δ (δ ³¹P) = -41.6 ppm. [Mn₂(CO)₈(μ-PPh₂)(μ-I)], **Vc**, ν(CO) in cyclohexane, 2089 w, 2069 m, 2014 vs, 2002 s, 1962 s; δ ³¹P in

CDCl₃, -28.2. [Mn(CO)₄(Ph₂PH)X], δ ³¹P data, X = Cl, 34.5 ppm, J(PH) = 367 Hz, Δ (δ ³¹P) = -74.4 ppm; X = Br, δ ³¹P, 33.5, J(PH) = 364 Hz, Δ (δ ³¹P) = -73.4 ppm; X = I, δ ³¹P, 34.2, J(PH) = 373 Hz, Δ (δ ³¹P) = -74.1 ppm; ν(CO), X = Br, 2091 m, 2026 s, 2015 s, 1959 m. [Mn₂(CO)₈(μ-PPh₂)(μ-H)], ν(CO) in hexane, 2093 m, 2063 m, 2010 vs, 1999 m, 1965 s; δ ³¹P in CDCl₃, +156 ppm. *fac*-[Mn(CO)₃(Ph₂PH)₂X]; X = Cl, ν(CO) in CHCl₃, 2035 s, 1970 m, 1914 m; δ ³¹P (in CDCl₃), 42.2, J(PH) = 318 Hz, Δ (δ ³¹P) = -82.1 ppm; X = Br, ν(CO), 2036 s, 1972 m, 1915 m; δ ³¹P, 49.0, Δ (δ ³¹P) = -88.9 ppm; X = I, ν(CO), 2030 s, 2010 m, 1910 m. [MnRe(CO)₈(μ-PPh₂)(μ-Br)] (**III**). ν(CO) in cyclohexane, 2105 w, 2082 m, 2076 w, 2019 vs, 2005 m, 1955 m; δ ³¹P in CDCl₃, -57.8; the mass spectrum showed species corresponding to the loss of Cl and eight consecutive CO groups, as well as fragments attributed to the formation and breakdown of the recombination species [M₂(CO)₈(μ-PPh₂)(μ-Br)].

Acknowledgement

The financial support of this research work by the Natural Sciences and Engineering Research Council of Canada is acknowledged.

References

- (a) D. A. Roberts, G. R. Steinmetz, M. J. Breen, P. M. Shulman, E. D. Morrison, M. R. Duttera, C. W. DeBrasse, R. R. Whittle and G. L. Geoffroy, *Organometallics*, **2**, 846 (1983); (b) A. D. Harley, G. J. Guskey and G. L. Geoffroy, *Organometallics*, **2**, 53 (1982); (c) M. J. Breen, M. R. Duttera, G. L. Geoffroy, G. C. Novotnak, D. A. Roberts, P. M. Shulman and G. R. Steinmetz, *Organometallics*, **1**, 1008 (1982); (d) G. F. Pregaglia, A. Andreatta, G. F. Ferrari, G. Montrasi and R. Ugo, *J. Organomet. Chem.*, **33**, 73 (1971); (e) A. J. Carty, *Adv. Chem. Ser.*, **196**, 163 (1982); (f) R. B. Finke, G. Gaughan, C. Pierpont and M. E. Cass, *J. Am. Chem. Soc.*, **103**, 1394 (1981); (g) G. Huttner, J. Schneider, H. D. Muller, G. Mohr, J. von Seyerl and L. Wohlfahrt, *Angew. Chem., Int. Ed. Engl.*, **18**, 76 (1979); (h) H. Beurich, T. Madach, F. Richter and H. Vahrenkamp, *Angew. Chem., Int. Ed. Engl.*, **18**, 690 (1979); (i) R. A. Jones, T. C. Wright, J. L. Atwood and W. E. Hunter, *Organometallics*, **2**, 470 (1983); (j) A. J. Carty, S. A. MacLaughlin and N. J. Taylor, *J. Organomet. Chem.*, **204**, C27 (1981).
- (a) R. P. Rosen, G. L. Geoffroy, C. L. Bueno, M. R. Churchill and R. G. Ortega, *J. Organomet. Chem.*, **254**, 89 (1983); (b) H. Schafer, *Angew. Chem., Int. Ed. Engl.*, **20**, 608 (1981); (c) P. Braunstein, D. Matt, O. Bars, M. Louer, D. Grandjean, J. Fischer and A. Mitschler, *J. Organomet. Chem.*, **213**, 79 (1981); (d) M. R. Churchill, C. Bueno and D. A. Young, *J. Organomet. Chem.*, **213**, 139 (1981); (e) R. J. Haines, N. D. C. T. Steen and R. B. English, *Chem. Commun.*, 587 (1981); *J. Organomet. Chem.*, **209**, C34 (1981); (f) F. Iwasaki, M. J. Mays, P. R. Raithby, P. L. Taylor and P. J. Wheatley, *J. Organomet. Chem.*, **213**, 185 (1981); (g) G. Huttner, J. Schneider,

- G. Mohr and J. von Seyerl, *J. Organomet. Chem.*, **191**, 161 (1980); (h) R. Lal, D. J. von Seyerl and G. Huttner, *J. Organomet. Chem.*, **178**, 319 (1979); (i) E. Keller and H. Vahrenkamp, *Chem. Ber.*, **112**, 2347 (1979); (j) R. Muller and H. Vahrenkamp, *J. Organomet. Chem.*, **170**, C25 (1979); (k) W. Malish and P. Panster, *Angew. Chem., Int. Ed. Engl.*, **16**, 408 (1977); (l) L. Staudacher and H. Vahrenkamp, *Chem. Ber.*, **109**, 218 (1979); (m) K. Yasufuku and H. Yamazaki, *Bull. Chem. Soc., Jpn.*, **46**, 1502 (1973); (n) B. C. Benson, R. Jackson, K. K. Joshi and D. T. Thompson, *Chem. Commun.*, 1506 (1968); (o) J. Chatt and D. T. Thompson, *J. Chem. Soc.*, 2713 (1964); (p) R. G. Hayter, *J. Am. Chem. Soc.*, **86**, 823 (1964); (q) J. Grobe and N. Sheppard, *Z. Naturforsch., Teil B*, **23**, 901 (1968); (r) H. Schafer, *Z. Naturforsch., Teil B*, **33**, 351 (1978); (s) J. Grobe and F. Kober, *Z. Naturforsch., Teil B*, **24**, 1660 (1969).
- 3 (a) E. W. Abel and I. H. Sabherwal, *J. Organomet. Chem.*, **10**, 491 (1967); (b) H. Schumann and H.-J. Kroth, *J. Organomet. Chem.*, **32**, C47 (1971).
- 4 E. Lindner and B. Schilling, *Chem. Ber.*, **110**, 3725 (1977).
- 5 P. M. Treichel, in G. Wilkinson, F. G. A. Stone and E. J. Abel (eds.), 'Comprehensive Organometallic Chemistry', Vol. 4, Pergamon, 1982.
- 6 J. Grobe and W. Mohr, *J. Fluorine Chem.*, **8**, 145, 341 (1976).
- 7 M. A. El-Sayed and H. D. Kaesz, *Inorg. Chem.*, **2**, 158 (1963).
- 8 R. L. Keiter and M. J. Madigan, *Organometallics*, **1**, 409 (1982).
- 9 R. J. Angelici and F. Basolo, *J. Am. Chem. Soc.*, **84**, 2495 (1962).
- 10 P. M. Treichel, W. K. Dean and W. M. Douglas, *J. Organomet. Chem.*, **42**, 145 (1972).
- 11 J. A. Iggo, M. J. Mays and P. R. Raithby, *J. Chem. Soc. Dalton*, 205 (1983).
- 12 (a) H. J. Emelius and J. J. Grobe, *Angew. Chem.*, **74**, 400 (1962); (b) R. C. Dobbie, *J. Chem. Soc., A*, 230 (1971); (c) J. Grobe, J. E. Helgerud and H. Stierand, *Z. Anorg. Allg. Chem.*, **371**, 123 (1969).
- 13 (a) K. Yasufuku and H. Yamazaki, *J. Organomet. Chem.*, **28**, 415 (1971); (b) K. Yasufuku and H. Yamazaki, *Bull. Chem. Soc. Jpn.*, **46**, 1502 (1973).
- 14 (a) W. Hieber and E. Winter, *Chem. Ber.*, **97**, 1037 (1964); (b) P. M. Treichel, W. K. Dean and W. M. Douglas, *Inorg. Chem.*, **11**, 1609 (1972); (c) R. C. Dobbie and D. Whittaker, *J. Chem. Soc. D*, 796 (1970).
- 15 R. C. Dobbie and D. Whittaker, *J. Chem. Soc., Dalton Trans.*, 2427 (1973).
- 16 (a) G. Beysel, J. Grobe and W. Mohr, *J. Organomet. Chem.*, **170**, 319 (1979); (b) J. Grobe and R. Rau, *J. Organomet. Chem.*, **157**, 281 (1978); (c) J. Grobe and R. Rau, *J. Fluorine Chem.*, **11**, 265 (1978).
- 17 (a) K. J. Reimer and A. Shaver, *Inorg. Synth.*, **19**, 159 (1979); (b) M. H. Quick and R. J. Angelici, *Inorg. Synth.*, **19**, 160 (1979).
- 18 (a) G. Dolcetti and J. R. Norton, *Inorg. Synth.*, **16**, 35 (1976); (b) H. D. Kaesz, R. Bau, D. Hendrickson and J. M. Smith, *J. Am. Chem. Soc.*, **89**, 2844 (1967).
- 19 G. W. Luther and G. Beyerle, *Inorg. Synth.*, **17**, 186 (1977).
- 20 P. M. Treichel, in G. Wilkinson, F. G. A. Stone and E. W. Abel (eds.), 'Comprehensive Organometallic Chemistry', Vol. 4, Pergamon, 1984.