

ERRATUM

Int J Legal Med (1998) 111 (2):67–77

Location and frequency of polymorphic positions in the mtDNA control region of individuals from Germany

S. Lutz · H.-J. Weisser · J. Heizmann · S. Pollak
Institut für Rechtsmedizin, Klinikum der Universität Freiburg,
Albertstr. 9, D-79104 Freiburg, Germany

In transferring the sequencing results from the manuscript to the printed tabular form shifts have occurred in some positions. The correct version of the table is presented here.

Table with columns for individual ID and nucleotide sequence (A, T, C, G). The table lists 181 individuals and their corresponding mtDNA control region sequences. The sequence is presented in a grid format where each row represents an individual and each column represents a position in the sequence. The sequence is: A T C A A C T T T T C A C C T T C G G T G C A A C C T C C C C A C D T C - C T G C A A C C T A T C C C T G C C C T A A C C C T C C C A.

individual		
158-200		
197		
170		
188		
191		
174		
152-196		
117,171,182,183,185		
148		
140		
146		
139		
121		
179		
114		
180		
142		
153		
75		
108		
118		
162		
158		
110		
93		
126		
132		
152		
176		
165		
164		
147		
137		
151		
156		
173		
71		
189		
175		
128		
99		
145		
155		
115		
80		
190		
154		
82		
136		
100		
107		
105		
59		
172		
133		
130		
5		
104		
61		
73		
68		
53		
54		
27		
169		
83		
41		
161		
34		
177,186		
77		
181		
98		
128		
149		
116		
184		
98		
96		
65		
20		
60		
79		
120		
166		
97		
168		
163		
36		
112		
89		
127,16		
85		
167		
25		
96		
55		
74		
159		
9		
122		
123		
90		
40		
17		
150		
113		
129		
125		
106		
157		
131		
134		
72		
50		

