

Radiographic analysis of epiphyseal fusion at knee joint to assess likelihood of having attained 18 years of age

R. Cameriere · M. Cingolani · A. Giuliadori ·
S. De Luca · L. Ferrante

Received: 11 June 2012 / Accepted: 31 July 2012 / Published online: 12 August 2012
© Springer-Verlag 2012

Abstract Radiological analysis of the epiphyses of the knee joint provides new valuable information, which may be used in combination with these well-established techniques in order to maximise the accuracy in the assessment of age of 18 years. A total of 215 antero-posterior radiographs of the knee was reviewed retrospectively in patients aged between 14 and 24 years old (99 boys, 116 girls). Fusion was scored as stage 1, epiphysis not fused; stage 2, epiphysis is fully ossified and epiphyseal scar is visible; and stage 3, epiphysis is fully ossified and epiphyseal scar is not visible. Scores of 0, 1 and 2 were assigned to stages 1, 2 and 3, respectively. Lastly, the score related to epiphyseal fusion at the knee joint was obtained by adding the three scores of the distal femur, proximal tibia and proximal fibula. Age distribution gradually increased with each score, for both genders. The mean age (\pm standard error) in each score category varied between genders, but the differences were not significant ($p > 0.11$). Five tests were performed to discriminate between individuals who were or were not at age 18 years or more,

according to the receiver operating curve. For boys, the highest value of accuracy was obtained with score 3, with high sensitivity (Se=93.33 %) and specificity (Sp=89.29 %). For girls, it was obtained with score 4, with high accuracy (Acc=85.86 %). These results indicate that radiographic analysis of the knee is a valuable alternative as a non-invasive method of estimation of 18 years of age.

Keywords Age estimation · Knee · Forensic science · ROC curve · Logistic regression · Reliability

Introduction

During the last few years, a growing section of forensic anthropology concerns the study of living individuals and, in particular, the problem of identifying and estimating age in living young adults for legal purposes [1–3]. Verified documentation of the date of birth is the only way of determining the exact age of an individual. However, in subjects who do not possess proper identification documents, it is of the utmost importance to verify whether these persons should be accepted as juveniles or adults [1–6]. This is so not only in criminal prosecutions but also in civil hearings, including determination of refugee status. Although age thresholds change according to country of interest, in most countries, the age of relevance to criminal liability ranges between 14 and 18 years [7–9].

To estimate the age of an individual for whom proper documentation is not available, forensic practitioners must rely on population data concerning growth and development [10–13]. These data enable the creation of a biological profile of the developmental status of an individual of unknown age, thus leading to an estimate of age. This profile is based on assessment of skeletal and dental anatomical growth markers [9, 14–18]. The search for the best

R. Cameriere (✉) · A. Giuliadori · S. De Luca · L. Ferrante
AgEstimation Project, Institute of Legal Medicine,
University of Macerata,
Via Don Minzoni,
62100 Macerata, Italy
e-mail: r.cameriere@unimc.it

M. Cingolani
Institute of Legal Medicine, University of Macerata,
Macerata, Italy

S. De Luca
Department of Legal Medicine,
Toxicology and Physical Anthropology, Faculty of Medicine,
University of Granada,
Granada, Spain

L. Ferrante
Department of Biomedical Sciences and Public Health,
Polytechnic University of Marche,
Ancona, Italy

anatomical growth markers which provide the most reliable age assessments has been ongoing for decades [1].

It must be stressed that biological variations in human development mean that any assessment of age based on analysis of anatomical growth markers is only an estimate and will, thus, contain a degree of error. The size of that error depends to a great extent on the anatomical markers used, the statistical methods applied to construct an age estimation table or formula based on those markers, and also on the developmental characteristics of the population assessed [19–21].

Forensic practitioners have a variety of age estimation methods available to them [1]. These include physical examination by a suitably qualified medical practitioner; assessment of the degree of dental development, which should include inspection of the oral cavity and analysis of panoramic radiographies; and assessment of skeletal maturity by radiological analysis of the left hand [22]. In addition, if it is uncertain whether the subject has reached the age of 21, conventional radiological, CT (computed tomography) or ultrasound examination of the clavicles should also be performed to expand the analytical spectrum. Depending on the question to be answered, these methods may be more or less reliable [1–3].

Nowadays, a new research focuses on the development of multi-factorial methods for age estimation, whereby a combination of methods (or anatomical growth markers) can provide the most accurate age estimate with the smallest possible error [1]. The difficult task for forensic researchers assessing age for the purposes of the legal system is still to convey accurately the degree of error inherent in whatever method is used and to clarify the likelihood that an individual below 18 years of age will be wrongly classified as an adult [1, 2, 8]. New techniques have been developed for older teenage individuals, which can provide age estimates with substantially narrower age ranges at the 95 % confidence level [23–26].

From the mid-teens to the early twenties, there are only a few reliable predictors of age of individuals, and dentition is one of them [27, 28]. In childhood, observation of dentition status results in highly accurate age assessment. However, this accuracy decreases as a person's dental development is completed. As a person grows beyond these years, developmental variability increases as the effects of the environment begin to outweigh those of genetics, thus making age estimations far more precise in early childhood and relatively imprecise during adolescence [29]. At around the age of 14 years, most teeth are fully developed, and age estimation becomes increasingly difficult. Generally, at this stage, after formation of the premolars and canines, the third molars are the only teeth still developing. Hence, although development of the third molar may be unpredictable and the tooth itself has a relatively high incidence of agenesis, the third

molar has been the subject of great interest as a predictor of 18 years of age and status of majority [30–32].

Although the hand–wrist region and third molar development have received the greatest attention in the assessment of age [33–35], radiological analysis of the epiphyses of long bones can overcome the deficiencies of non-adult skeletal collections and allow researchers to obtain contemporary information on epiphyseal union. The knee represents an ideal anatomical area which can also provide new information and which can be used in combination with the above well-established techniques in order to maximise age estimation accuracy. In this study, the knee joint was selected for several reasons. It is an anatomical area which can easily be radiographed at low radiation doses, with no interposed anatomical structures; the knee can also be easily positioned for antero-posterior radiographs; and, last but not the least, it yields information for three epiphyses at the same time, i.e. the distal femur, proximal tibia and proximal fibula.

There are many anthropological studies of skeletal maturation of the knee, based on dry bone and X-ray examinations, but they differ with regard to numerous variables: study population, gender, number of individuals, age range and number of bone fusion stages [36–41].

The most exhaustive anthropological study of the knee was that of McKern and Stewart, who examined the bodies of American soldiers who had died during the Korean War [36].

On the other hand, radiographic data on the development of the knee had been collected during a number of longitudinal studies. This has resulted in a number of reference standards which can be used in age estimation, and all are based on the assessment of radiographs [37, 42, 43].

The most exhaustive in terms of skeletal maturation was that of Pyle and Hoerr [37]. This atlas was developed from data collected during the Cleveland Study which ran in North America in 1926. The longitudinal study into child growth involved the collection of anthropometric data and radiographs from birth through to 21 years of age. In total, the data of 4,483 children formed the dataset which was also enhanced by radiographs of children collected in Boston by Dr. Harold C. Stuart [37]. The results were collected in an atlas published in 1955, consisting of X-rays of male and female subjects and including antero-posterior and lateral incidences of the right knee. The atlas method differs from that presented in the previous two approaches since it is not based upon a scoring system but is based on finding the best match between the radiographic image of the subject and the series of radiographs displayed in the atlas.

Although a wealth of knowledge is apparently provided, it is extremely difficult to find a consensus view on the definable stages of change between no mineralized bridge formation, non-union and complete ossification. Unfortunately, the

authors do not agree on the definition of complete fusion; some take into account the persistence of an epiphyseal scar, some do not [37–41].

The aim of the present study was to develop a radiographic scoring system for the assessment of epiphyseal union at the knee joint in order to estimate the likelihood of having attained 18 years of age in an Italian sample.

Materials and methods

Sample

This is a retrospective cross-sectional study of the relationship between epiphyseal fusion at the knee joint and chronological age. A sample of knee radiographs from Macerata Hospital (Italy) was studied. All X-ray projections were obtained with the same X-ray device (technical data of X-rays equipment and settings were as follows: FFD 110 cm, no grid, 110 mA, 7 mAs, 55 kV, CR Kodak® software, multifrequency processing). They belong to 215 healthy subjects (99 females and 116 males) aged between 14 and 24 years. Table 1 lists the age and gender distribution of each age category. About 55 % of the individuals had attained 18 years. Upper and lower age limits for the sample were derived from an extensive review of the literature and were refined during the data gathering process. All individuals had an antero-posterior radiograph of the knee available for assessment.

Radiographs were excluded from the study if they showed fractures or dislocations involving the growth plate, medical fixatives (i.e. surgical implants or casts) near the diaphyseal–epiphyseal junction or if they had been obtained from subjects with a medical history of chronic disease which may significantly alter skeletal development. Radiographs of subjects with known endocrine, metabolic or nutritional disorders were also excluded from the study. Protocols to collect radiographs for human subjects were approved by the Ethics Committee for Research Involving Human Subjects of the University of Macerata (Italy), and the study was conducted in accordance with the ethical standards laid down by the Declaration of Helsinki. The World Medical Association developed the Declaration of Helsinki as a statement of ethical principles for medical research involving human subjects, including research on identifiable human material and data. The chronological age

of the individual was calculated by calculating the difference between date of the image and the date of birth.

When necessary, digital images were submitted to contrast enhancement for easier interpretation. The following information was recorded for each subject: (1) date of birth and date of radiograph; (2) gender, as reported on the radiograph; (3) ancestry, reported by either self/family or visual evaluation by hospital staff; and (4) stage of fusion for each epiphysis. However, the observer was blinded to information on the subject's gender and age. Based on the combined use of these criteria, the observer was asked to assign a stage of union for each of the epiphyses of the distal femur, proximal tibia and proximal fibula.

Radiographic staging

To assess the degree of ossification of the distal femur, proximal tibia and proximal fibula, classification into three stages for each bone was applied as follows: stage 1, epiphysis is not fused (Fig. 1); stage 2, epiphysis is fully ossified and epiphyseal scar is visible (Fig. 2); stage 3, epiphysis is fully ossified and epiphyseal scar is not visible (Fig. 3). Distal femoral, proximal tibial and fibular epiphyses were separately evaluated according to these stages.

Scores of 0, 1 and 2 were assigned to stages 1, 2 and 3, respectively. Lastly, the score related to the epiphyseal fusion at the knee joint (SKJ) was obtained by adding the three scores of the distal femur, proximal tibia and proximal fibula. In cases of uncertainty during an observation, the lower stage was retained, according to *in dubio pro reo*, i.e. the principle of benefit of the doubt used in criminal proceedings.

Several subjects in this sample had radiographs of both knees available for assessment. However, in this study, knee laterality was not taken into account and only the right knee joint was used for the analysis. Previous studies indicate that epiphyseal union usually occurs at the same time on both sides [36, 44–46]. Auerbach and Ruff [47], in their study of limb bone bilateral asymmetry in modern humans, showed that all asymmetries are more pronounced in the upper limb, favouring the right side. Conversely, most lower limb dimensions, especially those around the knee, show much less asymmetry [47]. This information is extremely useful if only the epiphyses of one side are available, as may occur in some forensic investigations.

Table 1 Frequency distribution by gender and age cohort

Age (years)	14	15	16	17	18	19	20	21	22	23	24	Total
Girls	12	14	9	6	15	11	7	8	6	5	6	99
Boys	8	19	17	12	10	9	9	11	6	5	10	116
Total	20	33	26	18	25	20	16	19	12	10	16	215



Fig. 1 Stage 1: epiphysis is not fused

Statistical analysis

Subjects' radiographs were analysed by the same author (RC). In order to evaluate intra-observer reliability, the observer made repeated measures of 30 images at an interval of 2 weeks. Inter-observer error was assessed based on a random sub-sample of radiographs (30 images), which were assessed by three investigators, one odontologist and two physicians with many years of experience in the field of radiology, anatomy and forensic research. Cohen's kappa non-parametric test was used to evaluate intra- and inter-observer agreement.

With individual age as a dichotomous response variable ($E=1$ if an individual is at least 18 years of age, $E=0$ otherwise) and with gender and score of epiphyseal union at the knee joint (SKJ) as predictor variables, a generalised linear model was developed to predict whether an individual is older ($E=1$) or younger ($E=0$) than 18 years of age, with a logistic model as link function.

The significant variables between gender and SKJ were used to test the medico-legal question as to whether an individual is older or younger than 18 years of age. The predictive accuracy of the model was assessed by determining the receiver operating characteristic (ROC).

ROC curves provide three important ways of examining the efficacy of a test [48, 49]. First, they can evaluate the

overall ability of an indicator variable to make correct classifications, as in this study, of whether an individual is older than 18 (adult age) or not. Their shape shows the discriminatory ability of the indicator variable examined. The area under the curve (AUC) is widely recognised as the measure of the discriminatory power of a diagnostic test. The maximum value for the AUC is 1.0, thus indicating a (theoretically) perfect test (i.e. 100 % sensitive and 100 % specific). An AUC value of 0.5 indicates no discriminative value (i.e. 50 % sensitive and 50 % specific) and is represented by a straight diagonal line extending from the lower left corner to the upper right of the graph. That is, the closer the apex of the curve to the upper left corner, the greater the discriminatory ability of the test (i.e. the true-positive rate is high and the false-positive [1–specificity] rate is low). This is measured quantitatively by the AUC, so that a value of >0.96 indicates excellent discriminatory ability. Ninety-five percent confidence intervals for accuracy were calculated with a non-parametric method [48, 49].

The second function of ROC curves is to allow direct comparison of the abilities of different indicator variables to make correct classifications through a common metric, the AUC. Lastly, ROC curves facilitate the selection of a



Fig. 2 Stage 2: epiphysis is fully ossified and epiphyseal scar is visible



Fig. 3 Stage 3: epiphysis is fully ossified and epiphyseal scar is not visible

maximum threshold value of the indicator variable which best balances sensitivity and specificity. For each indicator variable, the curve shows the trade-off between sensitivity and specificity for any potential threshold [48, 49].

The test was performed to identify a threshold (cutoff) among the seven scores of epiphyseal union at the knee joint which could be used to assign an individual to the population of those younger ($T=0$) or older ($T=1$) than 18. Sensitivity p_1 (i.e. the proportion of individuals who had attained 18 years of age which verifies event $T=1$) was evaluated, as were its specificity p_2 (i.e. the proportion of individuals younger than 18 who verify event $T=0$) and correct classification (test accuracy, i.e. the proportion of all correct results). Lastly, the SKJ can help to discriminate between individuals who are or are not at age 18 years or more by the positive predictive value, PPV or post-test probability of being 18 years of age or more (i.e. the proportion of individuals aged 18 or over in whom event $E=1$ is verified). According to Bayes' theorem, PPV may be written as follows:

$$PPV = \frac{p_1 p_0}{p_1 p_0 + (1 - p_2)(1 - p_0)}$$

where PPV is the positive predictive value and p_0 is the probability that an individual is 18 or older, being between 14 and 24 years of age (our target population). This probability, p_0 , was evaluated with data from Italian Institute of Statistics [50] and was $p_0=0.66$. Statistical analyses were performed with the R statistical program [51]. The significance threshold was set at 5 %.

Results


Age distribution gradually increases with each SKJ category for both girls and boys (Fig. 4). The mean age (\pm standard error) for girls and boys in each score category varies between genders (Table 2), but the differences were not significant ($p>0.11$). The mean age provides an indication of the typical age at which each stage of union occurs.

Regarding to the repeatability and reproducibility of the method, κ statistics did not reveal significant intra-observer effects ($\kappa=0.974$ with 95 % bootstrapped confidence interval, 0.934–0.988), indicating substantial homogeneity of evaluation between two observations of the same knee radiographs. Cohen's kappa test demonstrated no statistically significant inter-observer differences in the age estimation performed by the three examiners: good inter-observer agreement for the distal femoral epiphysis ($\kappa=0.89$), for the proximal tibial epiphysis ($\kappa=0.81$) and for the proximal fibula ($\kappa=0.80$) was observed.

In order to study the extent to which the age of 18 years or more of an individual is related to SKJ and the dichotomous variable gender (1 for boys, 0 for girls), probability p , that an individual is at least 18 years of age, was modelled by a linear logistic model. To examine the effect of including or excluding one of the two factors in the model, the difference in deviance between two nested models was examined (Table 3). The change in deviance on adding the variable gender to a model which includes a constant term alone (null model) was not significant ($p=0.313$). Instead, when the scores were added to the null model, the deviance was reduced by a significant amount ($p<0.001$). In summary, the probability that an individual is aged 18 years or more depends on the score of epiphyseal union at the knee joint but does not significantly depend on gender.

Both estimated probability with the logistic model and frequency distribution by the scores of individuals who had attained 18 years are listed in Table 4, which shows that SKJ values 0 and 1 cannot be used to test adult age. In fact, we found that a very small percentage of individuals in the first two scores were adults. Consequently, taking into account only the SKJ values, five tests were performed to discriminate between individuals who were or were not aged

Fig. 4 Age distribution of boys (left) and girls (right) by scores



18 years or more, considering the scores from 2 to 6 as cutoffs. Results are listed in Table 5.

The predictive accuracy of these tests and their discrimination capacity were also assessed by determining the ROC curves by classification matrices for different levels of predicted probability that an individual is of age. The resulting ROC curves for boys and girls are shown in Fig. 5. The values for the area under the ROC (\pm standard error) were 0.961 (\pm 0.018) and 0.915 (\pm 0.023) for boys and girls, respectively.

For boys, the highest value of accuracy, i.e. the proportion of all tests which give a correct result, was obtained with score 3 (Acc=91.38 %) with high sensitivity (Se=93.33 %) and specificity (Sp=89.29 %). The estimated positive predictive value PPV was 94.42 %, with a 95 % confidence interval, CI (%)=(88.78; 97.31).

For girls, the highest value of accuracy was obtained with both scores 3 and 4 (Acc=85.86 %). However, when score 4 was chosen as cutoff, the ROC curve for girls appeared closest to the point (0, 1) of the graph which represents the ideal point at which the true-positive fraction is 1.0 (sensitivity=1.0) and the false-positive fraction is 0 (specificity=1.0).

Table 2 Age in relation to scores of epiphyseal union at knee joint for girls and boys

Score	Males <i>n</i>	M \pm SD	Females <i>n</i>	M \pm SD
0	14	14.5 \pm 0.2	19	14.9 \pm 0.2
1	10	15.8 \pm 0.5	18	15.7 \pm 0.2
2	15	16.8 \pm 0.7	17	16.9 \pm 0.5
3	8	18.1 \pm 0.8	11	18.2 \pm 0.3
4	16	18.3 \pm 0.3	14	18.9 \pm 0.4
5	15	20.4 \pm 0.5	15	20.8 \pm 0.5
6	21	21.2 \pm 0.6	22	22.5 \pm 0.3

M mean age, *SD* standard deviation, *n* number of cases

In addition, for forensic purposes, it is important that the test shows a low proportion of individuals younger than 18 whose test is positive (high specificity level). It therefore seemed appropriate to focus on the possibility of a false positive than on that of a false negative. The choice of score 4 to characterise adult age for girls yielded a specificity higher than that obtained when score 3 was chosen (Table 5). The PPV was 94.27 %, with a 95 % CI of 86.56 % (97.68 %).

Discussion

Many papers have been published concerning age estimation in young adults and many different methods have been developed, although some of them have been considered inaccurate [1, 52]. The choice of a particular method depends on the specific conditions of each case and mainly on the accuracy required [1]. In adolescence (14–21 years), third molars are the only teeth undergoing maturation, and they are especially attractive as a study subject because the degree of mineralization can easily be ascertained with non-invasive methods such as radiology [7–10]. However, several biological, ethnic and dietary motivations may substantially influence third molar growth [10]. In addition, as third molars are sometimes missing, for reasons of agenesis or extraction, estimating the age of majority based on these teeth is impossible [52].

Table 3 Deviance on fitting considered linear models to data

	df	Deviance	Residual df	Residual deviance	<i>p</i>
Null	–	–	214	296	–
Gender	1	1.02	213	294.98	0.313
Score	1	166.58	212	128.4	<0.001

Terms were added sequentially (first to last)

Table 4 Conditional probability, *p*, and relative frequency, *fr*, that an individual with a given score has attained 18 years (expressed as percentage)

Score	<i>p</i> (%)	<i>fr</i> (%)
0	2.16	0.00
1	7.95	3.57
2	25.25	28.13
3	56.89	68.42
4	83.76	83.33
5	95.27	96.67
6	98.75	95.35

Previous studies also examined the medial extremity of the clavicle, which has been exhaustively studied by plain X-rays, multi-slice computed tomography and ultrasound [53–58], and the wrist, which has also been closely studied with the Greulich and Pyle radiographic atlas [60]. For this reason, and in order to offer a valuable alternative method for assessing 18 years of age, a different anatomical region was analysed in this study.

Although the skeletal maturation of the knee has already been studied anthropologically on dry bone [36], rare radiographic studies have evaluated its potential forensic interest in the bone maturation process [37, 38]. O'Connor et al. [38] investigated the relationship between stage of epiphyseal union at the knee joint and chronological age in a modern Irish population. They used a novel radiographic method which subdivides the continuum of development into five specific stages of union. Regarding the application of an imaging modality with no radiation risk, one MRI study of the knee concerned only the proximal tibial epiphysis [61]. Dedout et al. [62] developed an original MRI staging system for epiphyseal fusion of growth plate maturation of the knee and evaluated its reliability and validity for age assessment of living individuals.

In this cross-sectional study, the radiographic staging system of growth plate maturation and epiphyseal fusion of the knee is analysed. The present results indicate that, in general, intra-observer variability is good for all bones ($\kappa=0.974$, with 95 % bootstrapped confidence interval, 0.934–0.988). In previous studies [e.g. 38], intra-observer error was higher for the fibula than for the tibia and femur probably because, in some instances, a good view of the fibula may be influenced by the position of the knee, which may partially obscure the fibula. However, in these radiographs, systematic evaluation of image blackening, peripheral blackening, appropriate exposure settings and the visibility of image details and contrast increased the likelihood of good identification of all parameters.

The ROC curves indicate that radiographic study of the bone maturation of the knee joint is highly correlated with the age of majority of living girls and boys. In addition, in these final stages of ossification of the knee, no significant differences between genders were observed. This trend was

Table 5 Percentages of sensitivity, specificity and accuracy (95 % confidence interval) of test of adult age when scores of epiphyseal union at knee joint are used to discriminate between individuals who are or are not at age 18 years or more

Gender	Parameter	Score	Score					
			2	3	4	5	6	
Boys	Se	100.00 % (94.04 %; –)	93.33 % (83.80 %; 98.15 %)	78.33 % (65.80 %; 87.93 %)	60.00 % (46.54 %; 72.44 %)	36.67 % (24.59 %; 50.10 %)		
	Sp	66.07 % (52.19 %; 78.19 %)	89.29 % (78.12 %; 95.97 %)	92.86 % (82.71 %; 98.02 %)	98.21 % (90.45 %; 99.95 %)	100.00 % (93.62 %; –)		
	Acc	83.62 % (75.61 %; 89.84 %)	91.38 % (84.72 %; 95.79 %)	85.34 % (77.58 %; 91.22 %)	78.45 % (69.85 %; 85.54 %)	67.24 % (57.21 %; 75.67 %)		
	PPV	85.12 % (79.88 %; 89.18 %)	94.42 % (88.78 %; 97.31 %)	95.51 % (89.13 %; 98.22 %)	98.49 % (90.24 %; 99.78 %)	100.00 % (–; –)		
Girls	Se	98.28 % (90.76 %; 99.96 %)	89.66 % (78.83 %; 96.11 %)	82.76 % (70.57 %; 91.41 %)	58.62 % (44.93 %; 71.40 %)	32.76 % (21.01 %; 46.34 %)		
	Sp	56.10 % (39.75 %; 71.53 %)	80.49 % (65.13 %; 91.18 %)	90.24 % (76.87 %; 97.28 %)	95.12 % (83.47 %; 99.40 %)	95.12 % (83.47 %; 99.40 %)		
	Acc	80.81 % (71.66 %; 88.03 %)	85.86 % (77.41 %; 92.05 %)	85.86 % (77.41 %; 92.05 %)	73.74 % (63.93 %; 82.07 %)	58.59 % (48.24 %; 68.40 %)		
	PPV	81.29 % (75.43 %; 86.02 %)	89.92 % (82.64 %; 94.35 %)	94.27 % (86.56 %; 97.68 %)	95.89 % (85.58 %; 98.92 %)	92.88 % (76.25 %; 98.15 %)		




Fig. 5 Receiver operating characteristic curves for “18 years of age or older” status for boys (continuous line) and girls (dashed line)

found to be consistent in both girls and boys by O'Connor et al. [38], in which the correlation between age (in years) and stage of union of the femur, tibia and fibula in boys produced correlation coefficients of 0.868, 0.858 and 0.802 ($p < 0.01$), respectively. As regards girls, the correlation test examining the relationship between age (in years) and stage of fusion of the epiphyses of the same bones gave coefficients of 0.880, 0.877 and 0.910 ($p < 0.01$), respectively. However, due the different classifications and staging used to characterise the growth plate, any comparison with previous radiological studies of the knee is unfortunately difficult.

As regards absence of statistical differences between genders, it was noted that the femoral and tibial epiphyses in boys and girls exhibited similar rates of fusion. These results were not consistent with previous radiological studies in which the distal femoral epiphyseal fusion occurs between the ages of 14–18 years in girls and 16–19 years in boys [41, 63]. On the Pyle and Hoerr radiographic standards, for instance, there is definite fusion in the centre of the epiphyseal plate at 14.5 years in girls and 17 years in boys, and the growth plates are replaced by lines of fusion by 15.5 years in girls and 18 years in boys [37]. Concerning proximal tibial fusion times, the standard for completion of distal femoral fusion is 14.5 years in girls and 17 years in boys [37]. Other observations have rather later times for complete fusion, up to 17 years in girls and 19.5 years in boys. However, as with other atlases, the use of the Pyle and Hoerr [37] approach to assess chronological age raises a number of methodological issues. The data which formed the basis of the atlas were collected from children who were

growing to adulthood in the first decades of the twentieth century and were described by the authors as being white children of high socioeconomic class. Additionally, the children were deliberately chosen for their good health and nutritional status [64]. This background, both socioeconomically and ethnically, is very different from that of children being age estimated for forensic purposes today. Maturational rate is known to depend on a large number of factors not least of which are the nutritional intake and the health status of the individual. The applicability of an age estimation method which was developed on a population potentially so far removed from a modern population must be therefore tested robustly if it is to be used and accepted for forensic purposes.

Analysis of the ROC curve showed that, for girls, the choice of score 4 to characterise adult age yielded specificity higher than that obtained when score 3 was chosen. In fact, when score 4 was chosen as cutoff, the ROC curve showed its point closest to the point (0, 1) of the graph, which represents the ideal point at which the true-positive fraction is 1.0 (sensitivity=1.0) and the false-positive fraction is 0 (specificity=1.0). For forensic purposes, it is important that the test shows a low proportion of individuals younger than 18 whose test is positive (high specificity level). From the practical perspective applied to forensic sciences, the above results are of special importance when we consider the fact that, from both a criminal viewpoint and according to undocumented immigrant acceptance policies, the age limit to adopt urgent criminal or acceptance measures is 18 years.

Regarding the responsible use of X-rays in forensic age estimations, it is indispensable not only to use dose-lowering techniques but also to take the presumed age of the person being examined into greater account [59, 65, 66]. Established doses for X-ray examinations in forensic age estimations vary from less than 0.1 mSv (left hand–wrist X-ray) to more than 800 mSv (CT) per examination [65, 67]. However, the use of conventional X-rays is not appropriate for individuals under 18, and, like the collar bone CT, it should be recommended as a method only to answer the question of whether the subject is over 18/21 years [65, 68]. Article 3 of the European Council Directive 97/43/Euratom states that when exposing an individual to ionising radiation, the net benefit to the individual must outweigh the risks [4]. Nowadays, more recent data are also available that compare radiation exposure from medical X-rays with the hazards of everyday living [69, 70]. Based on these studies, the resulting risk from using X-rays in age estimation procedures (with the exception of CT scan on sternoclavicular joints) is very low in comparison to other life risks [65]. However, the need for the moment is to minimise all radiation as far as possible and to evaluate risks from radiation when performing X-ray examinations in age estimation procedures [65].

Conclusions

The radiographic technique offers a valuable alternative as a non-invasive method of estimation of 18 years of age. In this study, very encouraging results were obtained when this technique was applied to analyse epiphyseal fusion at the knee joint. However, there are some problems in using a radiographic data set: one involves the integrity of the X-rays and the operator's experience. Although radiographic positions and exposure variables are standardised, the biological variability of tissue thickness and bone density require a certain level of experience to obtain an optimal film series. Nowadays, as hospitals move toward digital X-rays, these considerations may have less influence. The second problem involves observer experience in interpreting radiographic films. Because of the two-dimensional nature of radiographs, which causes overlaps from surrounding structures, an observer may incorrectly assess the stage of fusion, unless multiple radiographic views are available.

In view of the above difficulties, it is very important to standardise the methodology for assessment of epiphyseal union. Further studies with larger sample groups of both antero-posterior and medio-lateral incidences are therefore recommended in order to verify not only the accuracy of the method but also inter-observer agreement. The scoring technique must be clearly defined so that it can be applied by forensic researchers all over the world. As Cunha et al. [1] noted, sometimes the best methods are not those with the best published standard error but those which have been tested by many researchers on many different populations, which are suitable for a specific forensic scenario, practical, user-friendly, relatively quick and economical.

Acknowledgments The authors would like to thank the staff of the Department of Radiology at Macerata Hospital (Italy) for their assistance in this project and also Ms. Gabriel Walton for editing the English text. The authors are also grateful to anonymous reviewers for their comments and suggestions, which greatly improved the manuscript.

References

- Cunha E, Baccino E, Martrille L, Ramsthaler F, Prieto J, Schuliar Y, Lynnerup N, Cattaneo C (2009) The problem of ageing human remains and living individuals: a review. *Forensic Sci Int* 193:1–13
- Schmelting A, Reisinger W, Geserick G, Olze A (2006) Age estimation of unaccompanied minors Part I. General considerations. *Forensic Sci Int* 159s:s61–s64
- Olze A, Reisinger W, Geserick G, Schmelting A (2006) Age estimation of unaccompanied minors Part II. *Forensic Sci Int* 159s:s65–s67
- Aynsley-Green A (2009) Unethical age assessment. *Br Dent J* 206:337
- Kvittingen AV (2011) Negotiating childhood: age assessment in the UK asylum system. Working paper series Oxford: Refugee Studies Centre, Oxford Department of International Development, University of Oxford
- Department of Homeland Security (2009) Age determination practices for unaccompanied alien children in ICE custody. Department of Homeland Security, Washington, DC
- Lewis ME, Flavel A (2006) Age assessment of child skeletal remains in forensic context. In: Cunha E, Pinhero J (eds) *Forensic Anthropology and Medicine*. Humana, Totowa, p 464
- Schmelting A, Olze A, Reisinger W, Geserick G (2001) Age estimation of living people undergoing criminal proceedings. *Lancet* 358:89–90
- Olze A, Solheim T, Schulz R, Kupfer M, Schmelting A (2010) Evaluation of the radiographic visibility of the root pulp in the lower third molars for the purpose of forensic age estimation in living individuals. *Int J Legal Med* 124:183–186
- Liversidge HM (2008) Timing of human third molar formation. *Ann Hum Biol* 35:294–321
- Knottnerus JA, Muris JW (2009) The evidence base of clinical diagnosis, 2nd edn, Assessment of the accuracy of diagnostic tests: the cross-sectional study. BMJ Publishing, London, pp 42–62
- Habemma JDF, Eijkemans R, Krinjen P, Knottnerus A (2009) Analysis of data on the accuracy of diagnostic tests. In: Knottnerus JA, Buntinx F (ed) *The evidence base of clinical diagnosis*, 2nd edn. BMJ Publishing, London, pp 118–145
- Konigsberg LW, Herrmann NP, Westcott DJ, Kimmerle EH (2008) Estimation and evidence in forensic anthropology: age-at-death. *J Forensic Sci* 53:541–557
- Bhat VJ, Kamath GP (2007) Age estimation from root development of mandibular third molars in comparison with skeletal age of wrist joint. *Am J Forensic Med Pathol* 28:238–241
- Schmidt S, Baumann U, Schulz R, Reisinger W, Schmelting A (2008) Study of age dependence of epiphyseal ossification of the hand skeleton. *Int J Legal Med* 122:51–54
- Schmidt S, Nitz I, Schulz R, Schmelting A (2008) Applicability of the skeletal age determination method of Tanner and Whitehouse for forensic age diagnostics. *Int J Legal Med* 122:309–314
- Cardoso HF (2008) Epiphyseal union at the innominate and lower limb in a modern Portuguese skeletal sample, and age estimation in adolescent and young adult male and female skeletons. *Am J Phys Anthropol* 135:161–170
- Hedge RJ, Sood PB (2002) Dental maturity as an indicator of chronological age: radiographic evaluation of dental age in 6 to 13 years children of Belgaum using Demirjian's methods. *J Indian Soc Pedod Prev Dent* 20:132–138
- Solari AC, Abramovitch AK (2002) The accuracy and precision of third molar development as an indicator of chronological age in Hispanics. *J Forensic Sci* 47:531–535
- Willems G, Van Olmen A, Spiessens B, Carels C (2001) Dental age estimation in Belgian children: Demirjian's technique revisited. *J Forensic Sci* 46:893–895
- Deeks JJ, Altman DG (2004) Likelihood ratios. *Br Med J* 329:168–169
- Schmidt S, Koch B, Schulz R, Reisinger W, Schmelting A (2007) Comparative analysis of the applicability of the skeletal age determination methods of Greulich–Pyle and Thiemann–Nitz for forensic age estimation in living subjects. *Int J Legal Med* 121:293–296
- Cameriere R, De Luca S, Biagi R, Cingolani M, Farronato G, Ferrante L (2012) Accuracy of three age estimation methods in children by measurements of developing teeth and carpals and epiphyses of the ulna and radius. *J Forensic Sci*. doi:10.1111/j.1556-4029.2012.02120

24. Liversidge HM, Smith HB, Maber M (2010) Bias and accuracy of age estimation using developing teeth in 946 children. *Am J Phys Anthropol*. doi:10.1002/ajpa.21349
25. Liversidge HM (2010) Demirjian stage tooth formation results from a large group of children. *Dent Anthropol* 23:16–24
26. Hillewig E, De Tobel J, Cuhe O, Vandemaele P, Piette M, Verstraete K (2010) Magnetic resonance imaging of the medial extremity of the clavicle in forensic bone age determination: a new four-minute approach. *Eur Radiol* 21:757–767
27. Mincer HH, Harris EF, Berryman HE (1993) The ABFO study of third molar development and its use as an estimator of chronological age. *J Forensic Sci* 38:379–390
28. Liliequist B, Lundberg M (1971) Skeletal and tooth development, a methodological investigation. *Acta Radiol* 11:97–112
29. Bassed B, Briggs C, Drummer OH (2011) Age estimation and the developing third molar tooth: an analysis of an Australian population using computed tomography. *J Forensic Sci*. doi:10.1111/J.1556-4029.01769.X
30. Kullman L, Johanson G, Akesson L (1992) Root development of the lower third molar and its relation to chronological age. *Swed Dent J* 16:161–167
31. Kieser JA (1990) Human adult odontometrics: The study of variation in adult tooth size. Cambridge University Press, New York
32. Bolaños MV, Moussa H, Manrique MC, Bolaños MJ (2003) Radiographic evaluation of third molar development in Spanish children and young people. *Forensic Sci Int* 133:212–219
33. Koc A, Karaoglanoglu M, Erdogan M, Kosecik M, Cesur Y (2001) Assessment of bone ages: is the Greulich–Pyle method sufficient for Turkish boys? *Pediatr Int* 43:662–665
34. Cameriere R, Ferrante L (2008) Age estimation in children by measurement of carpals and epiphyses of radius and ulna and open apices in teeth: a pilot study. *Forensic Sci Int* 174:60–63
35. Berst MJ, Dolan L, Bogdanowicz MM, Stevens MA, Chow S, Brandser EA (2001) Effect of knowledge of chronologic age on the variability of pediatric bone age determined using the Greulich and Pyle Standards. *AJR* 176:507–510
36. McKern TW, Stewart TD (1957) Skeletal age changes in young American males. Analysed from the Standpoint of Age Identification. Technical Report EP-45 Natick Mass
37. Pyle SI, Hoerr NL (1955) Radiographic atlas of skeletal development of the knee. A Standard of Reference Thomas Springfield IL
38. O'Connor JE, Bogue C, Spence LD, Last J (2008) A method to establish the relationship between chronological age and stage of union from radiographic assessment of epiphyseal fusion at the knee: an Irish population study. *J Anat* 212:198–209
39. Schaefer MC, Black SM (2005) Comparison of ages of epiphyseal union in North American and Bosnian skeletal material. *J Forensic Sci* 50:777–784
40. Schaefer MC, Black SM (2007) Epiphyseal union sequencing: aiding in the recognition and sorting of commingled remains. *J Forensic Sci* 52:277–285
41. Das Gupta SM, Prasad V, Singh S (1974) A roentgenologic study of epiphyseal union around elbow, wrist and knee joints and the pelvis in boys and girls of Uttar Pradesh. *J Indian Med Assoc* 62:10–12
42. Acheson RM (1952) Radiographs of the hand as an index of skeletal maturity in infants. *Arch Dis Child* 27:382–385
43. Roche AF, Davila GH (1976) The reliability of assessments of the maturity of individual hand-wrist bones. *Hum Biol* 48:585
44. Albert AM, Greene DL (1999) Bilateral asymmetry in skeletal growth and maturation as an indicator of environmental stress. *Am J Phys Anthropol* 110:341–349
45. Schmeling A, Schulz R, Reisinger W, Muhler M, Wernecke KD, Geserick G (2004) Studies on the time frame for ossification of the medial clavicular epiphyseal cartilage in conventional radiography. *Int J Legal Med* 118:5–8
46. Falyys CG, Schutkowski H, Weston DA (2006) Auricular surface aging: worse than expected? A test of the revised method on a documented historic skeletal assemblage. *Am J Phys Anthropol* 130:508–513
47. Auerbach BM, Ruff C (2006) Limb bone bilateral asymmetry: variability and commonality among modern humans. *J Hum Evol* 50:203–218
48. Prieto JL, Barbería E, Ortega R, Magaña C (2005) Evaluation of chronological age based on third molar development in the Spanish population. *Int J Legal Med* 119:349–354
49. Martín-de las Heras S, García-Fortea P, Ortega A, Zococovich S, Valenzuela A (2008) Third molar development according to chronological age in populations from Spanish and Magrebian origin. *Forensic Sci Int* 174:47–53
50. Istituto Nazionale di Statistica (ISTAT) <http://demo.istat.it/pop2004/index.html> (Last access 28/05/2012)
51. R Development Core Team (2008) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org> (Last access 28/05/2012).
52. Schmeling A, Garamendi PM, Prieto JL, Landa MI (2011) Forensic age estimation in unaccompanied minors and young living adults. In: Duarte NV (ed) Forensic medicine—from old problems to new challenges InTech, 77–120.
53. Singh J, Chavali KH (2011) Age estimation from clavicular epiphyseal union sequencing in a Northwest Indian population of the Chandigarh region. *J Forensic Legal Med* 18:82–87
54. Schmeling A, Schulz R, Reisinger W, Mühler M, Wernecke KD, Geserick G (2004) Studies on the time frame for ossification of medial clavicular epiphyseal cartilage in conventional radiography. *Int J Legal Med* 118:5–8
55. Garamendi PM, Landa MI, Botella MC, Alemán I (2011) Forensic age estimation on digital X-ray images: medial epiphyses of the clavicle and first rib ossification in relation to chronological age. *J Forensic Sci* 56:S3–S12
56. Kellinghaus M, Schulz R, Vieth V, Schmidt S, Pfeiffer H, Schmeling A (2010) Enhanced possibilities to make statements on the ossification status of the medial clavicular epiphysis using an amplified staging scheme in evaluating thin-slice CT scans. *Int J Legal Med* 124:321–325
57. Kellinghaus M, Schulz R, Vieth V, Schmidt S, Schmeling A (2010) Forensic age estimation in living subjects based on the ossification status of the medial clavicular epiphysis as revealed by thin-slice multidetector computed tomography. *Int J Legal Med* 124:149–154
58. Hillewig E, De Tobel J, Cuhe O, Vandemaele P, Piette M, Verstraete K (2011) Magnetic resonance imaging of the medial extremity of the clavicle in forensic bone age determination: a new four-minute approach. *Eur Radiol* 21:757–767
59. Schulz R, Zwiesigk P, Schiborr M, Schmidt S, Schmeling A (2008) Ultrasound studies on the time course of clavicular ossification. *Int J Legal Med* 122:163–167
60. Greulich W, Pyle SI (1959) Radiographic atlas of skeletal development of the hand and wrist, 2nd edn. Stanford University Press, Stanford
61. Jopp E, Schröder I, Maas R, Adam G, Püschel K, Hertzog C (2010) A Proximale Tibiaepiphyse im Magnetresonanztomogramm. *Rechtsmedizin* 20:464–468
62. Dedouit F, Auriol J, Rousseau H, Rougé D, Crubézy E, Telmon N (2012) Age assessment by magnetic resonance imaging of the knee: a preliminary study. *Forensic Sci Int* 217:232.e1–232.e7
63. Flecker H (1932) Roentgenographic observations of the times of appearance of epiphyses and their fusion with the diaphyses. *J Anat* 67:118–164

64. Behrents RG, Broadbent BH Jr. A chronological account of the Bolton-Brush Growth Studies, 1984. Available at: <http://www.cwru.edu/dental/bbgsc/phase5.html> (Last access July 3, 2012)
65. Ramsthaler F, Proschek P, Betz W, Verhoff MA (2009) How reliable are the risk estimates for X-ray examinations in forensic age estimations? A safety update. *Int J Legal Med* 123:199–204
66. Willems G, Moulin-Romsee C, Solheim T (2002) Non-destructive dental-age calculation methods in adults: intra- and inter-observer effects. *Forensic Sci Int* 126:221–226
67. Huda W, Atherton JV, Ware DE, Cumming WA (1997) An approach for the estimation of effective radiation dose at CT in pediatric patients. *Radiology* 203:417–422
68. Maher MM, Kalra MK, Toth TL, Wittram C, Saini S, Shepard J (2004) Application of rational practice and technical advances for optimizing radiation dose for chest CT. *J Thorac Imaging* 19:16–23
69. Hall EJ (2009) Radiation biology for pediatric radiologists. *Pediatr Radiol* 39(suppl 1):57–64
70. Hall EJ, Brenner DJ (2008) Cancer risks from diagnostic radiology. *Br J Radiol* 81:362–378