

## Subject index Volume 231

### Ab initio calculations

Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59

Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83

Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197

### Accurate mass

A fragmentation study of kaempferol using electrospray quadrupole-time-of-flight mass spectrometry at high mass resolution, 157

### Acetone

Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69

### *N*-Acetoxy-2-acetylaminofluorene

Analysis of the in vitro digestion of modified DNA to oligonucleotides by LC–MS and LC–MS/MS, 169

### Aluminium fluoride

Mass spectrometric determination of partial pressures of ions in the saturated vapor over the NaF–Na<sub>3</sub>AlF<sub>6</sub> system, 31

### Averaging

Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1

### Binary cluster ion

Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77

### BINOL

Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139

### CAD

Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179

### Carboxylate groups

Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113

### Catalyzed isomerizations

Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83

### Catechol estrogens

Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119

### Central ion effects

Isomeric differentiation and quantification of  $\alpha$ ,  $\beta$ -amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103

### CH bond activation

Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139

### Charge state

Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131

### Chemical ionization mass spectrometer

Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17

### CID

Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59

### Collisionally induced dissociation

Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207

### Collision-induced dissociation

Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69

### Complexes

Isomeric differentiation and quantification of  $\alpha$ ,  $\beta$ -amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103

### Cryolite

Mass spectrometric determination of partial pressures of ions in the saturated vapor over the NaF–Na<sub>3</sub>AlF<sub>6</sub> system, 31

### Density functional theory

Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59

### Diastereoselectivity

Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139

### Dimer

Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207

### 1,4-Dioxane

Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77

### Dissociation

Isomeric differentiation and quantification of  $\alpha$ ,  $\beta$ -amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103

### Distonic ions

Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197

### Distribution

Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131

### Disubstituted benzene

Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69

### Electrospray

Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131

Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189

- Electrospray ionization  
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119  
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
- Electrospray-ionization mass spectrometry  
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
- Enantioselectivity  
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139
- Energy resolved mass spectrometry  
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119
- Enzymatic digestion  
Analysis of the in vitro digestion of modified DNA to oligonucleotides by LC-MS and LC-MS/MS, 169
- Escape velocity  
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1
- ESI-MS  
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113
- Flavonoid  
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
- FT-ICR mass spectrometry  
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83  
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197
- FT-ICR-MS  
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37
- Gas phase derivatization  
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
- Geometry factor  
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37
- Glutathione conjugates  
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119
- Heated capillary  
Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189
- Hydrogen abstraction  
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197
- Initial velocity  
Initial velocity of secondary ions from XY-TOF technique, simultaneous calibration by residual gas ionization, 51
- Internal energy  
Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189
- Ion dynamics  
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1
- Ion pressures  
Mass spectrometric determination of partial pressures of ions in the saturated vapor over the NaF–Na<sub>3</sub>AlF<sub>6</sub> system, 31
- Ion trap mass spectrometry  
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119
- Ionized carbenes  
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197
- Ionized dihydroxycarbenes  
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83
- Ionized formic acid  
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83
- Ion–molecule equilibrium  
Mass spectrometric determination of partial pressures of ions in the saturated vapor over the NaF–Na<sub>3</sub>AlF<sub>6</sub> system, 31
- Ion–molecule reaction  
Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69
- IPM  
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramidate mustard, 147
- Isomeric tripeptides  
Isomeric differentiation and quantification of  $\alpha$ ,  $\beta$ -amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103
- Isophosphoramidate mustard  
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramidate mustard, 147
- Kaempferol  
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
- Kinetic method  
Isomeric differentiation and quantification of  $\alpha$ ,  $\beta$ -amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103
- LC-ES-MS/MS  
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramidate mustard, 147
- Liquid chromatography–tandem mass spectrometry  
Analysis of the in vitro digestion of modified DNA to oligonucleotides by LC-MS and LC-MS/MS, 169
- Low temperature mass spectrometry  
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37
- Manganese complexes  
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139
- Mass spectrometric fragmentation pathway  
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47
- Mass spectrometry  
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131  
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139

- Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramidate mustard, 147
- Metallothionein  
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207
- MM/MD  
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
- Nanospray  
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207
- Non-denaturing conditions  
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
- Nonlinear Mathieu equation  
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1
- Oligonucleotide adducts  
Analysis of the *in vitro* digestion of modified DNA to oligonucleotides by LC–MS and LC–MS/MS, 169
- 1,3,4-Oxadiazoles  
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47
- Paul trap  
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1
- Peptide analysis  
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
- Peptides  
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113
- Perchloric acid  
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramidate mustard, 147
- Phosphopeptides  
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
- Poly(vinyl acetate)  
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
- Position sensitive detector  
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
- Pressure effect  
Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189
- Proteins  
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
- Proteomics  
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113
- Quantification  
Isomeric differentiation and quantification of  $\alpha$ ,  $\beta$ -amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103
- Relative humidity  
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17
- Secondary ion desorption  
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
- Skeletal rearrangement  
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47
- Sodium ion interactions  
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113
- Substituent effect  
Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69
- Sulfur dioxide  
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17
- Sulfuric acid  
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17
- Tandem mass spectrometry  
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
- Temperature-controlled ICR cell  
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37
- 1,3,4-Thiadiazoles  
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47
- Time-of-flight  
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
- Time-of-flight mass spectrometry  
Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77
- Trimethyl borate  
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
- Unimolecular decomposition  
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
- Vanadium complexes  
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139
- Water  
Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77
- Water clusters  
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17
- XY*-TOF  
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
- Zinc chelates  
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207