

Subject index Volume 251

- Accurate mass
Truly “exact” mass: Elemental composition can be determined uniquely from molecular mass measurement at ~0.1 mDa accuracy for molecules up to ~500 Da, 260
- Alpha-hydroxy acid
Collision induced dissociation of alpha hydroxy acids: Evidence of an ion-neutral complex intermediate, 40
- α , β and proton radioactivity
Decay Q -value of ^{105}Sn and of other nuclei near ^{100}Sn , measured at the GSI on-line mass separator, 138
- Antibaryon
Antiproton mass measurements, 273
- Antiproton
Antiproton mass measurements, 273
- Antiquark
Antiproton mass measurements, 273
- Atomic masses
The history of nuclidic masses and of their evaluation, 85
Mass measurements and superallowed β decay, 95
Mass ratio of two ions in a Penning trap by alternating between the trap center and a large cyclotron orbit, 125
On-line commissioning of SHIPTRAP, 146
The Q -value of tritium β -decay and the neutrino mass, 173
Precision experiments on exotic nuclei at IGISOL, 204
The UW-PTMS: Systematic studies, measurement progress, and future improvements, 231
Precision measurements with highly charged ions at rest: The HITRAP project at GSI, 266
Mass measurements of the shortest-lived nuclides *à la MISTRAL*, 286
- Atomic physics
Mass measurements and the bound-electron g factor, 102
- Avogadro constant
Primary mass standard based on atomic masses, 220
- Beta decay
The WITCH experiment: Completion of a set-up to investigate the structure of weak interactions with a Penning trap, 159
The Q -value of tritium β -decay and the neutrino mass, 173
- Binding energies and masses
Decay Q -value of ^{105}Sn and of other nuclei near ^{100}Sn , measured at the GSI on-line mass separator, 138
- Binding energy
On-line commissioning of SHIPTRAP, 146
Precision measurements with highly charged ions at rest: The HITRAP project at GSI, 266
- Biomolecules hydration
Hydration of gas-phase protonated alkylamines, amino acids and dipeptides produced by electrospray, 23
- Bismuth
Primary mass standard based on atomic masses, 220
- Bond enthalpies
Hydration of gas-phase protonated alkylamines, amino acids and dipeptides produced by electrospray, 23
- Bound states
Mass measurements and the bound-electron g factor, 102
- Charge breeding
Mass measurements on highly charged radioactive ions, a new approach to high precision with TITAN, 198
- CO ice
Secondary ion emission from condensed CO bombarded by fission fragments, 1
- Collinear laser spectroscopy
Precision experiments on exotic nuclei at IGISOL, 204
- Condensed gas
Secondary ion emission from condensed CO bombarded by fission fragments, 1
- Cooling of highly charged ions
Mass measurements on highly charged radioactive ions, a new approach to high precision with TITAN, 198
- Cyclotron-frequency systematics
The UW-PTMS: Systematic studies, measurement progress, and future improvements, 231
- Cylindrical ion trap
Fabrication and testing of a miniature cylindrical ion trap mass spectrometer constructed from low temperature co-fired ceramics, 32
- Density functional theory
Large-scale self-consistent nuclear mass calculations, 243
- Density of distribution
The interpretation of mass spectra of inorganic molecules, 59
- Differential cross section
Electron-impact excitation of the $3s3p\ ^1P_1$ state of magnesium: Electron scattering at small angles, 66
- Dissociation mechanism
Collision induced dissociation of alpha hydroxy acids: Evidence of an ion-neutral complex intermediate, 40
- EBIT
Mass measurements on highly charged radioactive ions, a new approach to high precision with TITAN, 198
- Electron affinity
Mass spectrometric studies of trifluoromethylated fullerenes, 16
- Electron excitation
Electron-impact excitation of the $3s3p\ ^1P_1$ state of magnesium: Electron scattering at small angles, 66
- Electron mass
Highly charged ions, quantum-electrodynamics, and the electron mass, 152
Precision measurements with highly charged ions at rest: The HITRAP project at GSI, 266
- Electrospray
Hydration of gas-phase protonated alkylamines, amino acids and dipeptides produced by electrospray, 23
- Electrospray ionization mass spectrometry
Peptide sequencing through N-terminal phosphorylation and multistage electrospray mass spectrometry of sodiated molecules, 82

- Enhanced molecular ion
A new type of GC–MS with advanced capabilities, 47
- Erbium
On-line commissioning of SHIPTRAP, 146
- Evaluation of atomic masses
The history of nuclidic masses and of their evaluation, 85
- Exotic nuclei
Experiments with stored exotic nuclei at relativistic energies, 212
- Fast GC–MS
A new type of GC–MS with advanced capabilities, 47
- Fourier transform mass spectrometry
Truly “exact” mass: Elemental composition can be determined uniquely from molecular mass measurement at ~0.1 mDa accuracy for molecules up to ~500 Da, 260
- Fragment ion kinetic energy
Mean kinetic energy of molecular fragment ions from time-of-flight and momentum analysis, 10
- Fullerene
Mass spectrometric studies of trifluoromethylated fullerenes, 16
- Fundamental constants
QED effects in heavy few-electron ions, 109
- Gas-phase equilibria
Hydration of gas-phase protonated alkylamines, amino acids and dipeptides produced by electrospray, 23
- GC–MS
A new type of GC–MS with advanced capabilities, 47
- g*-Factor
Precision measurements with highly charged ions at rest: The HITRAP project at GSI, 266
Precise measurements of ionic masses for QED tests, 281
- HCI
Stopping highly charged ions in a laser-cooled one component plasma of $^{24}\text{Mg}^+$ ions, 179
- Heavy ions
QED effects in heavy few-electron ions, 109
- High RF fields
The use of high RFQ fields to manipulate ions, 190
- Highly-charged ions
Towards high-accuracy mass spectrometry of highly charged short-lived ions at ISOLTRAP, 131
Experiments with stored exotic nuclei at relativistic energies, 212
- History of atomic masses
The history of nuclidic masses and of their evaluation, 85
- History of mass spectrometry
The history of nuclidic masses and of their evaluation, 85
- Holmium
On-line commissioning of SHIPTRAP, 146
- ICR
Truly “exact” mass: Elemental composition can be determined uniquely from molecular mass measurement at ~0.1 mDa accuracy for molecules up to ~500 Da, 260
- Ion cluster
Secondary ion emission from condensed CO bombarded by fission fragments, 1
- Ion cyclotron resonance
Truly “exact” mass: Elemental composition can be determined uniquely from molecular mass measurement at ~0.1 mDa accuracy for molecules up to ~500 Da, 260
- Ion guide
Precision experiments on exotic nuclei at IGISOL, 204
- Ion mirror
A novel principle for an ion mirror design in time-of-flight mass spectrometry, 73
- Ion temperature
The use of high RFQ fields to manipulate ions, 190
- Ionization energy
Mass spectrometric studies of trifluoromethylated fullerenes, 16
- Ion-neutral complex
Collision induced dissociation of alpha hydroxy acids: Evidence of an ion-neutral complex intermediate, 40
- Isotopes
Precise measurements of ionic masses for QED tests, 281
- Isotopic labelling
Collision induced dissociation of alpha hydroxy acids: Evidence of an ion-neutral complex intermediate, 40
- Krypton isotope
Precision mass measurements with LEBIT at MSU, 300
- Lamor precision frequency
Precise measurements of ionic masses for QED tests, 281
- Large scale calculation
Large-scale self-consistent nuclear mass calculations, 243
- Laser cooling
Stopping highly charged ions in a laser-cooled one component plasma of $^{24}\text{Mg}^+$ ions, 179
- LEAR
Antiproton mass measurements, 273
- LEBIT
Precision mass measurements with LEBIT at MSU, 300
- Low-energy beam
Precision mass measurements with LEBIT at MSU, 300
- LTCC
Fabrication and testing of a miniature cylindrical ion trap mass spectrometer constructed from low temperature co-fired ceramics, 32
- Magnesium
Electron-impact excitation of the $3s3p\ ^1P_1$ state of magnesium: Electron scattering at small angles, 66
- Magnetic moment
Highly charged ions, quantum-electrodynamics, and the electron mass, 152
- Many-body problem
Large-scale self-consistent nuclear mass calculations, 243
- Mass and half-life measurements
Experiments with stored exotic nuclei at relativistic energies, 212
- Mass measurement
Studies of neutron-rich isotopes with the CPT mass spectrometer and the CARIBU project, 252
- Mass spectra of inorganic molecules
The interpretation of mass spectra of inorganic molecules, 59
- Mass spectrometer
The UW-PTMS: Systematic studies, measurement progress, and future improvements, 231
- Mass spectrum coefficient
The interpretation of mass spectra of inorganic molecules, 59
- Metallized ceramic surface
Fabrication and testing of a miniature cylindrical ion trap mass spectrometer constructed from low temperature co-fired ceramics, 32
- Miniature
Fabrication and testing of a miniature cylindrical ion trap mass spectrometer constructed from low temperature co-fired ceramics, 32
- Molecular fragmentation
Mean kinetic energy of molecular fragment ions from time-of-flight and momentum analysis, 10
- Momentum spectrometry
Mean kinetic energy of molecular fragment ions from time-of-flight and momentum analysis, 10
- MS–MS
A new type of GC–MS with advanced capabilities, 47
- Multiple reflection
A novel principle for an ion mirror design in time-of-flight mass spectrometry, 73

- N=Z nuclides
 Mass measurements of the shortest-lived nuclides *à la MISTRAL*, 286
- Neutrino masses
 The Q -value of tritium β -decay and the neutrino mass, 173
- Neutron-rich isotopes
 Studies of neutron-rich isotopes with the CPT mass spectrometer and the CARIBU project, 252
- N-terminal phosphorylation
 Peptide sequencing through N-terminal phosphorylation and multistage electrospray mass spectrometry of sodiated molecules, 82
- Nuclear binding energies
 The history of nuclidic masses and of their evaluation, 85
 Mass measurements of the shortest-lived nuclides *à la MISTRAL*, 286
- Nuclear masses
 Opportunistic mass measurements at the Holifield Radioactive Ion Beam Facility, 119
 Large-scale self-consistent nuclear mass calculations, 243
 The importance of nuclear masses in the astrophysical rp-process, 293
- On-line isotope separation
 Decay Q -value of ^{105}Sn and of other nuclei near ^{100}Sn , measured at the GSI on-line mass separator, 138
- Oxygen
 The UW-PTMS: Systematic studies, measurement progress, and future improvements, 231
- Parity nonconservation effects
 QED effects in heavy few-electron ions, 109
- Penning trap
 Mass ratio of two ions in a Penning trap by alternating between the trap center and a large cyclotron orbit, 125
 Towards high-accuracy mass spectrometry of highly charged short-lived ions at ISOLTRAP, 131
 On-line commissioning of SHIPTRAP, 146
 The WITCH experiment: Completion of a set-up to investigate the structure of weak interactions with a Penning trap, 159
 Stopping highly charged ions in a laser-cooled one component plasma of $^{24}\text{Mg}^+$ ions, 179
 Precision experiments on exotic nuclei at IGISOL, 204
 Studies of neutron-rich isotopes with the CPT mass spectrometer and the CARIBU project, 252
 Precision measurements with highly charged ions at rest: The HITRAP project at GSI, 266
 Precise measurements of ionic masses for QED tests, 281
 Precision mass measurements with LEBIT at MSU, 300
- Penning trap mass spectrometry
 Mass measurements on highly charged radioactive ions, a new approach to high precision with TITAN, 198
- Penning-trap
 The UW-PTMS: Systematic studies, measurement progress, and future improvements, 231
- Peptide sequencing
 Peptide sequencing through N-terminal phosphorylation and multistage electrospray mass spectrometry of sodiated molecules, 82
- Petroleum
 Truly “exact” mass: Elemental composition can be determined uniquely from molecular mass measurement at ~ 0.1 mDa accuracy for molecules up to ~ 500 Da, 260
- Phosphoramidate peptide
 Peptide sequencing through N-terminal phosphorylation and multistage electrospray mass spectrometry of sodiated molecules, 82
- Photolithography
 Fabrication and testing of a miniature cylindrical ion trap mass spectrometer constructed from low temperature co-fired ceramics, 32
- Precision mass measurement
 Precision mass measurements with LEBIT at MSU, 300
- Precision mass spectrometry
 Towards high-accuracy mass spectrometry of highly charged short-lived ions at ISOLTRAP, 131
- Precision measurement
 Mass ratio of two ions in a Penning trap by alternating between the trap center and a large cyclotron orbit, 125
- QED
 Precise measurements of ionic masses for QED tests, 281
- Quantum-electrodynamics
 Mass measurements and the bound-electron g factor, 102
 QED effects in heavy few-electron ions, 109
 Highly charged ions, quantum-electrodynamics, and the electron mass, 152
- Q-value
 Mass measurements and superallowed β decay, 95
- Radio frequency ion cooler
 Precision experiments on exotic nuclei at IGISOL, 204
- Radio frequency quadrupole
 Stopping highly charged ions in a laser-cooled one component plasma of $^{24}\text{Mg}^+$ ions, 179
- Radioactive beams
 Mass measurements on highly charged radioactive ions, a new approach to high precision with TITAN, 198
- Radioactive ion beams
 Opportunistic mass measurements at the Holifield Radioactive Ion Beam Facility, 119
- Reflectron
 A novel principle for an ion mirror design in time-of-flight mass spectrometry, 73
- Relative atomic mass
 Primary mass standard based on atomic masses, 220
- Retardation spectrometer
 The WITCH experiment: Completion of a set-up to investigate the structure of weak interactions with a Penning trap, 159
- RF mass spectrometer
 Mass measurements of the shortest-lived nuclides *à la MISTRAL*, 286
- RFQ confinement
 The use of high RFQ fields to manipulate ions, 190
- rp-Process
 The importance of nuclear masses in the astrophysical rp-process, 293
- Salt systems
 The interpretation of mass spectra of inorganic molecules, 59
- Scalar interaction
 The WITCH experiment: Completion of a set-up to investigate the structure of weak interactions with a Penning trap, 159
- Secondary ions
 Secondary ion emission from condensed CO bombarded by fission fragments, 1
- Silicon
 Primary mass standard based on atomic masses, 220
- Simulation
 Stopping highly charged ions in a laser-cooled one component plasma of $^{24}\text{Mg}^+$ ions, 179
- Sodiated molecule
 Peptide sequencing through N-terminal phosphorylation and multistage electrospray mass spectrometry of sodiated molecules, 82
- Spontaneous symmetry breaking
 Large-scale self-consistent nuclear mass calculations, 243
- Storage rings
 Experiments with stored exotic nuclei at relativistic energies, 212
- Supersonic molecular beams
 A new type of GC-MS with advanced capabilities, 47
- Systematics
 Mass measurements and superallowed β decay, 95
- Temperature dependence
 Secondary ion emission from condensed CO bombarded by fission fragments, 1

- Time-of-flight mass spectrometry
 - A novel principle for an ion mirror design in time-of-flight mass spectrometry, 73
- Time-of-flight spectrometry
 - Mean kinetic energy of molecular fragment ions from time-of-flight and momentum analysis, 10
- Time-of-flight variance
 - Mean kinetic energy of molecular fragment ions from time-of-flight and momentum analysis, 10
- TOF SIMS
 - Secondary ion emission from condensed CO bombarded by fission fragments, 1
- Trifluoromethylated fullerene
 - Mass spectrometric studies of trifluoromethylated fullerenes, 16
- Triple quadrupole
 - Collision induced dissociation of alpha hydroxy acids: Evidence of an ion-neutral complex intermediate, 40
- Weak interaction
 - The WITCH experiment: Completion of a set-up to investigate the structure of weak interactions with a Penning trap, 159
- Wigner energy
 - Mass measurements of the shortest-lived nuclides *à la MISTRAL*, 286
- X-ray bursts
 - The importance of nuclear masses in the astrophysical rp-process, 293