

Subject index Volume 253

Absolute cross sections

Absolute partial cross sections and kinetic energy analysis for the electron impact ionization of propene, 122

Aerosol mass spectrometry

The heterogeneous reaction of particle-phase methyl esters and ozone elucidated by photoelectron resonance capture ionization: Direct products of ozonolysis and secondary reactions leading to the formation of ketones, 38

Aldehydes

On-line monitoring of UV-induced lipid peroxidation products from human skin in vivo using proton-transfer reaction mass spectrometry, 58

Alkali metal cluster

Structures and charge distributions of cationic and neutral Li_nX_m ($X = \text{Na}$ and K), 104

Ammonia ion cluster

Ion cluster desorption from frozen NH_3 induced by impact of fast multi-charged ions, 112

Analyte incorporation

Investigations of theoretical principles for MALDI-MS derived from solvent-free sample preparation. Part I. Preorganization, 13

Asymmetric dissociation

Gas-phase processes and measurements of macromolecular properties in solution: On the possibility of false positive and false negative signals of protein unfolding, 207

Atmospheric pressure reaction

Atmospheric pressure Eberlin transacetalization reactions in the heterogeneous liquid/gas phase, 281

Avizafone

Mass spectrometry study of increased breakdown of an anticonvulsant drug substance: Avizafone, 1

BeC_n^-

Parity alternation in the linear ground-state beryllium-doped carbon clusters BeC_n^- ($n = 1-8$), 30

Beryllium-doped anionic carbon clusters

Parity alternation in the linear ground-state beryllium-doped carbon clusters BeC_n^- ($n = 1-8$), 30

Binary cluster anions

Parity alternation in the linear ground-state beryllium-doped carbon clusters BeC_n^- ($n = 1-8$), 30

Bond dissociation energies

Sequential bond energies of water to sodium proline cation, 288

Calculation method

A calculation method to eliminate gain effect on isotopic measurement using the virtual amplifier multi-collector mass spectrometer, 130

Carbohydrate

From the gas phase to aqueous solution: Vibrational spectroscopy, Raman optical activity and conformational structure of carbohydrates, 193

Cation Mass Spectrometer

SIMS analysis with neutral cesium deposition: Negative secondary ion sensitivity increase and quantification aspects, 71

^{252}Cf -PDMS

Ion cluster desorption from frozen NH_3 induced by impact of fast multi-charged ions, 112

Charge distributions

Structures and charge distributions of cationic and neutral Li_nX_m ($X = \text{Na}$ and K), 104

Charge exchange process

Determination of concentration of excited oxygen atoms on the basis of ion mobility in atmospheric plasma, 22

Charge state distribution

Gas-phase processes and measurements of macromolecular properties in solution: On the possibility of false positive and false negative signals of protein unfolding, 207

Probing solution- and gas-phase structures of Trp-cage cations by chiral substitution and spectroscopic techniques, 263

Chirality

Probing solution- and gas-phase structures of Trp-cage cations by chiral substitution and spectroscopic techniques, 263

Chlorpyrifos

Effect of dissociation energy on ion formation and sensitivity of an analytical method for determination of chlorpyrifos in human blood, using gas chromatography–mass spectrometer (GC–MS in MS/MS), 48

Condensed NH_3

Ion cluster desorption from frozen NH_3 induced by impact of fast multi-charged ions, 112

Conformation

Fluorescence probe of polypeptide conformational dynamics in gas phase and in solution, 172

From the gas phase to aqueous solution: Vibrational spectroscopy, Raman optical activity and conformational structure of carbohydrates, 193

Criegee intermediate

The heterogeneous reaction of particle-phase methyl esters and ozone elucidated by photoelectron resonance capture ionization: Direct products of ozonolysis and secondary reactions leading to the formation of ketones, 38

Decreased desorption threshold

Investigations of theoretical principles for MALDI-MS derived from solvent-free sample preparation. Part I. Preorganization, 13

Degradation products

Mass spectrometry study of increased breakdown of an anticonvulsant drug substance: Avizafone, 1

Density functional study

Parity alternation in the linear ground-state beryllium-doped carbon clusters BeC_n^- ($n = 1-8$), 30

DFT

Gas-phase interaction of protonated lysine with water, 201

Diffusion coefficient

Characterization of a sheet membrane interface for sample introduction into a time-of-flight mass spectrometer, 65

Dissociation

Stability of complementary and mismatched DNA duplexes: Comparison and contrast in gas versus solution phases, 238

- Dissociation energy
Effect of dissociation energy on ion formation and sensitivity of an analytical method for determination of chlorpyrifos in human blood, using gas chromatography–mass spectrometer (GC–MS in MS/MS), 48
- Disulfide bond cleavage
Distance dependence of through-bond electron transfer rates in electron-capture and electron-transfer dissociation, 274
- DNA
ESI-MS fragmentation pathways of *N*-methylpyrrole polyamide/peptide conjugates, 141
G-quadruplexes in telomeric repeats are conserved in a solvent-free environment, 225
Stability of complementary and mismatched DNA duplexes: Comparison and contrast in gas versus solution phases, 238
- Drug–DNA complexes
Positive and negative ion mode ESI-MS and MS/MS for studying drug–DNA complexes, 156
- Duplex
Stability of complementary and mismatched DNA duplexes: Comparison and contrast in gas versus solution phases, 238
- Dynamics
Fluorescence probe of polypeptide conformational dynamics in gas phase and in solution, 172
- Eberlin reaction
Atmospheric pressure Eberlin transacetalization reactions in the heterogeneous liquid/gas phase, 281
- Electron capture dissociation
Probing solution- and gas-phase structures of Trp-cage cations by chiral substitution and spectroscopic techniques, 263
Distance dependence of through-bond electron transfer rates in electron-capture and electron-transfer dissociation, 274
- Electron capture induced dissociation
Atypical behavior in the electron capture induced dissociation of biologically relevant transition metal ion complexes of the peptide hormone oxytocin, 217
- Electron impact ionization
Electron impact ionization mass spectrometry of aliphatic alcohol clusters in helium nanodroplets, 79
Absolute partial cross sections and kinetic energy analysis for the electron impact ionization of propene, 122
- Electron transfer
Distance dependence of through-bond electron transfer rates in electron-capture and electron-transfer dissociation, 274
- Electrospray
Formation of hydrated triply charged metal ions from aqueous solutions using nanodrop mass spectrometry, 256
- Electrospray ionization
Gas-phase processes and measurements of macromolecular properties in solution: On the possibility of false positive and false negative signals of protein unfolding, 207
Segmental charge distributions of Cytochrome *c* on transfer into the gas phase, 249
- Energetics
Stability of complementary and mismatched DNA duplexes: Comparison and contrast in gas versus solution phases, 238
- ESI/MS/MS
ESI-MS fragmentation pathways of *N*-methylpyrrole polyamide/peptide conjugates, 141
- ESI-MS
Positive and negative ion mode ESI-MS and MS/MS for studying drug–DNA complexes, 156
- Evaporation experiments
Structures and charge distributions of cationic and neutral Li_nX_m ($X = \text{Na}$ and K), 104
- Fluorescence
Fluorescence probe of polypeptide conformational dynamics in gas phase and in solution, 172
- Fragmentation
Investigations of theoretical principles for MALDI-MS derived from solvent-free sample preparation. Part I. Preorganization, 13
- FTICR MS
Gas-phase interaction of protonated lysine with water, 201
- Gain factor
A calculation method to eliminate gain effect on isotopic measurement using the virtual amplifier multi-collector mass spectrometer, 130
- Gas-phase basicity
Positive and negative ion mode ESI-MS and MS/MS for studying drug–DNA complexes, 156
- Gas-phase interaction
Gas-phase interaction of protonated lysine with water, 201
- Gas-phase ion/ion proton transfer
Charge-state dependent dissociation of a trypsin/inhibitor complex via ion trap collisional activation, 147
- Gas-phase protein–protein complex
Charge-state dependent dissociation of a trypsin/inhibitor complex via ion trap collisional activation, 147
- Gas-phase structure
Probing solution- and gas-phase structures of Trp-cage cations by chiral substitution and spectroscopic techniques, 263
- GC–MS/MS
Effect of dissociation energy on ion formation and sensitivity of an analytical method for determination of chlorpyrifos in human blood, using gas chromatography–mass spectrometer (GC–MS in MS/MS), 48
- Geometric configuration
Structures and charge distributions of cationic and neutral Li_nX_m ($X = \text{Na}$ and K), 104
- G-quadruplexes
G-quadruplexes in telomeric repeats are conserved in a solvent-free environment, 225
- Guided ion beam mass spectrometry
Sequential bond energies of water to sodium proline cation, 288
- H/D exchange
Gas-phase interaction of protonated lysine with water, 201
- Helium nanodroplet
Electron impact ionization mass spectrometry of aliphatic alcohol clusters in helium nanodroplets, 79
- Heterogeneous reaction
Atmospheric pressure Eberlin transacetalization reactions in the heterogeneous liquid/gas phase, 281
- Human blood
Effect of dissociation energy on ion formation and sensitivity of an analytical method for determination of chlorpyrifos in human blood, using gas chromatography–mass spectrometer (GC–MS in MS/MS), 48
- Human skin
On-line monitoring of UV-induced lipid peroxidation products from human skin in vivo using proton-transfer reaction mass spectrometry, 58
- Hydration
From the gas phase to aqueous solution: Vibrational spectroscopy, Raman optical activity and conformational structure of carbohydrates, 193
Formation of hydrated triply charged metal ions from aqueous solutions using nanodrop mass spectrometry, 256
- In situ U–Pb geochronology
In situ petrographic thin section U–Pb dating of zircon, monazite, and titanite using laser ablation–MC–ICP–MS, 87
- In vivo
On-line monitoring of UV-induced lipid peroxidation products from human skin in vivo using proton-transfer reaction mass spectrometry, 58
- Internal energy
Fragmentation at and above surfaces in SIMS: Effects of biomolecular yield enhancing surface modifications, 181

- Ion fragmentation**
Electron impact ionization mass spectrometry of aliphatic alcohol clusters in helium nanodroplets, 79
- Ion kinetic energy**
Absolute partial cross sections and kinetic energy analysis for the electron impact ionization of propene, 122
- Ion mobility**
Determination of concentration of excited oxygen atoms on the basis of ion mobility in atmospheric plasma, 22
G-quadruplexes in telomeric repeats are conserved in a solvent-free environment, 225
- Ion solvation**
Formation of hydrated triply charged metal ions from aqueous solutions using nanodrop mass spectrometry, 256
- Ion/molecule reaction**
Atmospheric pressure Eberlin transacetalization reactions in the heterogeneous liquid/gas phase, 281
- Ionization mode**
Mass spectrometry study of increased breakdown of an anticonvulsant drug substance: Avizafone, 1
- Isotope analysis**
Resonance enhancement of ion signals in direct isotope analysis of metal samples by adopting 2-color resonant laser ablation mass spectrometry, 136
- Isotope measurement**
A calculation method to eliminate gain effect on isotopic measurement using the virtual amplifier multi-collector mass spectrometer, 130
- Kinetic energy distribution**
Fragmentation at and above surfaces in SIMS: Effects of biomolecular yield enhancing surface modifications, 181
- Labile molecules**
Investigations of theoretical principles for MALDI-MS derived from solvent-free sample preparation. Part I. Preorganization, 13
- Laser ablation**
In situ petrographic thin section U–Pb dating of zircon, monazite, and titanite using laser ablation–MC–ICP–MS, 87
- Laser mass spectrometer**
Laser mass spectrometric studies on rare earth doped UO_2 , 98
- Laser-induced plasma**
Resonance enhancement of ion signals in direct isotope analysis of metal samples by adopting 2-color resonant laser ablation mass spectrometry, 136
- Ligated trications**
Formation of hydrated triply charged metal ions from aqueous solutions using nanodrop mass spectrometry, 256
- Lipid peroxidation**
On-line monitoring of UV-induced lipid peroxidation products from human skin in vivo using proton-transfer reaction mass spectrometry, 58
- MALDI**
Investigations of theoretical principles for MALDI-MS derived from solvent-free sample preparation. Part I. Preorganization, 13
- Mass spectrometer**
A calculation method to eliminate gain effect on isotopic measurement using the virtual amplifier multi-collector mass spectrometer, 130
- Mass spectrometry**
Mass spectrometry study of increased breakdown of an anticonvulsant drug substance: Avizafone, 1
Determination of concentration of excited oxygen atoms on the basis of ion mobility in atmospheric plasma, 22
Absolute partial cross sections and kinetic energy analysis for the electron impact ionization of propene, 122
- Matrix-enhanced SIMS**
Fragmentation at and above surfaces in SIMS: Effects of biomolecular yield enhancing surface modifications, 181
- MC–ICP–MS**
In situ petrographic thin section U–Pb dating of zircon, monazite, and titanite using laser ablation–MC–ICP–MS, 87
- Membrane inlet**
Characterization of a sheet membrane interface for sample introduction into a time-of-flight mass spectrometer, 65
- Metal binding**
Atypical behavior in the electron capture induced dissociation of biologically relevant transition metal ion complexes of the peptide hormone oxytocin, 217
- Metal-assisted SIMS**
Fragmentation at and above surfaces in SIMS: Effects of biomolecular yield enhancing surface modifications, 181
- Mismatches**
Stability of complementary and mismatched DNA duplexes: Comparison and contrast in gas versus solution phases, 238
- Molecular cluster**
Electron impact ionization mass spectrometry of aliphatic alcohol clusters in helium nanodroplets, 79
- MS/MS**
Positive and negative ion mode ESI-MS and MS/MS for studying drug–DNA complexes, 156
- Multi-collector**
A calculation method to eliminate gain effect on isotopic measurement using the virtual amplifier multi-collector mass spectrometer, 130
- Negative ion mode**
Positive and negative ion mode ESI-MS and MS/MS for studying drug–DNA complexes, 156
- Neutral cesium deposition**
SIMS analysis with neutral cesium deposition: Negative secondary ion sensitivity increase and quantification aspects, 71
- Non-covalent complex**
Gas-phase processes and measurements of macromolecular properties in solution: On the possibility of false positive and false negative signals of protein unfolding, 207
- Oxytocin**
Atypical behavior in the electron capture induced dissociation of biologically relevant transition metal ion complexes of the peptide hormone oxytocin, 217
- Ozonolysis**
The heterogeneous reaction of particle-phase methyl esters and ozone elucidated by photoelectron resonance capture ionization: Direct products of ozonolysis and secondary reactions leading to the formation of ketones, 38
- Peptide**
Fluorescence probe of polypeptide conformational dynamics in gas phase and in solution, 172
- Petrographic thin section**
In situ petrographic thin section U–Pb dating of zircon, monazite, and titanite using laser ablation–MC–ICP–MS, 87
- Photoelectron resonance capture ionization**
The heterogeneous reaction of particle-phase methyl esters and ozone elucidated by photoelectron resonance capture ionization: Direct products of ozonolysis and secondary reactions leading to the formation of ketones, 38
- Polyamidic oligomer**
ESI-MS fragmentation pathways of *N*-methylpyrrole polyamide/peptide conjugates, 141
- Polydimethylsiloxane (PDMS)**
Characterization of a sheet membrane interface for sample introduction into a time-of-flight mass spectrometer, 65
- Precursor model**
Fragmentation at and above surfaces in SIMS: Effects of biomolecular yield enhancing surface modifications, 181
- Proline**
Fluorescence probe of polypeptide conformational dynamics in gas phase and in solution, 172
- Propene**
Absolute partial cross sections and kinetic energy analysis for the electron impact ionization of propene, 122

- Proton affinity
Gas-phase processes and measurements of macromolecular properties in solution: On the possibility of false positive and false negative signals of protein unfolding, 207
- Proton transfer
Segmental charge distributions of Cytochrome *c* on transfer into the gas phase, 249
- Proton-transfer reaction mass spectrometry
On-line monitoring of UV-induced lipid peroxidation products from human skin in vivo using proton-transfer reaction mass spectrometry, 58
- Quantification
SIMS analysis with neutral cesium deposition: Negative secondary ion sensitivity increase and quantification aspects, 71
- Raman
From the gas phase to aqueous solution: Vibrational spectroscopy, Raman optical activity and conformational structure of carbohydrates, 193
- Rare earth doped UO₂
Laser mass spectrometric studies on rare earth doped UO₂, 98
- Reaction mechanism
Gas-phase interaction of protonated lysine with water, 201
- Resonant laser ablation
Resonance enhancement of ion signals in direct isotope analysis of metal samples by adopting 2-color resonant laser ablation mass spectrometry, 136
- Secondary ion
Ion cluster desorption from frozen NH₃ induced by impact of fast multi-charged ions, 112
- SIMS
SIMS analysis with neutral cesium deposition: Negative secondary ion sensitivity increase and quantification aspects, 71
- Sodium proline cation
Sequential bond energies of water to sodium proline cation, 288
- Soft ionization
The heterogeneous reaction of particle-phase methyl esters and ozone elucidated by photoelectron resonance capture ionization: Direct products of ozonolysis and secondary reactions leading to the formation of ketones, 38
- Solution structure
Probing solution- and gas-phase structures of Trp-cage cations by chiral substitution and spectroscopic techniques, 263
- Solvated ion
Atmospheric pressure Eberlin transacetalization reactions in the heterogeneous liquid/gas phase, 281
- Solvation energies
Sequential bond energies of water to sodium proline cation, 288
- Solvent-free sample preparation
Investigations of theoretical principles for MALDI-MS derived from solvent-free sample preparation. Part I. Preorganization, 13
- Stability
Structures and charge distributions of cationic and neutral Li_{*n*}X_{*m*} (X = Na and K), 104
- Tandem mass spectrometry
Charge-state dependent dissociation of a trypsin/inhibitor complex via ion trap collisional activation, 147
- Threshold laser power
Investigations of theoretical principles for MALDI-MS derived from solvent-free sample preparation. Part I. Preorganization, 13
- Through-bond
Distance dependence of through-bond electron transfer rates in electron-capture and electron-transfer dissociation, 274
- Time-of-flight mass spectrometer
Characterization of a sheet membrane interface for sample introduction into a time-of-flight mass spectrometer, 65
- TOF
Ion cluster desorption from frozen NH₃ induced by impact of fast multi-charged ions, 112
- Trivalent metal ions
Formation of hydrated triply charged metal ions from aqueous solutions using nanodrop mass spectrometry, 256
- Trypsin/trypsin inhibitor complex
Charge-state dependent dissociation of a trypsin/inhibitor complex via ion trap collisional activation, 147
- Useful yield
SIMS analysis with neutral cesium deposition: Negative secondary ion sensitivity increase and quantification aspects, 71
- Volatile organic compound
On-line monitoring of UV-induced lipid peroxidation products from human skin in vivo using proton-transfer reaction mass spectrometry, 58