



Corrigendum

Corrigendum to ‘Tandem Fourier transform mass spectrometry of block and random copolymers’ [Int. J. Mass Spectrom. 301 (2011) 184–194]

Sasa M. Miladinovic, Cynthia J. Kaeser, Matthias M. Knust, Charles L. Wilkins*

Chemistry and Biochemistry Department, University of Arkansas, Fayetteville, AR 72701, USA

The authors regret that in the above-mentioned article Fig. 3 was published with an incorrect caption, and Figs. 7 and 8 were published in incorrect versions. The corrections can be found below.

The authors would like to apologise for any inconvenience caused.




Fig. 7. ESI-ECD-FT mass spectrum of (A) triply-charged (MMA)₄₃ oligomer, (B) triply-charged (MMA)₄₃ oligomer when SORI pulse was used to eject doubly-charged fragment with proposed fragmentation mechanism.

DOI of original article: [10.1016/j.ijms.2010.08.028](https://doi.org/10.1016/j.ijms.2010.08.028).

* Corresponding author at: 251 Chemistry Bldg., University of Arkansas, Fayetteville, AR 72701, USA. Tel.: +1 479 575 3160; fax: +1 479 575 4049.
 E-mail address: cwilkins@uark.edu (C.L. Wilkins).




Fig. 8. ESI-ECD-FT mass spectrum of (A) doubly-charged random $(\text{MMA})_{13}(\text{BMA})_{11}$ oligomer, (B) doubly-charged block $(\text{MMA})_{18}(\text{BMA})_{11}$ oligomer.

Fig. 3. (A) Full mass range MALDI-FT mass spectrum of rMMABMA, (B) MALDI-QCID-FT mass spectrum of random $(\text{MMA})_7(\text{BMA})_8$ singly-charged oligomer with one Na^+ cation, (C) full mass range MALDI-FT mass spectrum of bMMABMA, (D) MALDI-QCID-FT mass spectrum of block $(\text{MMA})_{14}(\text{BMA})_4$ singly-charged oligomer with one Na^+ cation.