

## AUTHOR INDEX

- Agyilrah, G.A., see Nixon, J.R., 277
- Al-Shora, H., see Said, S., 11
- Ali, H.M. and Farouk, A.M., The effect of Sudanese diet on the bioavailability of ampicillin, 301
- Ali, S.L. and Strittmatter, Th., Separation and determination of four ergot alkaloids, 183
- Andersson, J., see Morén, F., 295
- Armstrong, N.A. and James, K.C., Drug release from lipid-based dosage forms. I, 185
- Armstrong, N.A. and James, K.C., Drug release from lipid-based dosage forms. II, 195
- Badawi, A.A., Fouli, A.M. and El-Sayed, A.A., Drug release from matrices made of polymers with reacting sites, 55
- Barnett, M.I., Nyström, C. and Engvall, H., The importance of optical refraction in connection with the surface area measurement of powders by some photometric apparatus, 131
- Bethke, H., HPLC-separation of codergocrine-mesylate (dihydroergotoxine-methanesulfonate), 87
- Bundgaard, H., Acid-catalyzed hydrolysis of fosfomycin and its implication in oral absorption of the drug, 1
- Bundgaard, H., Falch, E. and Larsen, C., Prodrugs as drug delivery systems XI. Preparation and characterization of a novel water-soluble pro-drug type for barbituric acids, 19
- Bundgaard, H., see Hansen, J., 307
- Burkman, A.M., see Sokoloski, T.D., 63
- Cheng, K.C., see Shefter, E., 179
- Crommelin, D.J.A. and de Blaey, C.J., In vitro release studies on drugs suspended in non-polar media II. The release of paracetamol and chloramphenicol from suspensions in liquid paraffin, 29
- D'Arcy, P.F., see McElnay, J.C., 205
- de Blaey, C.J., see Crommelin, D.J.A., 29
- Dominguez-Gil, A., see Sanchez Camazano, M., 343
- El-Sayed, A.A., see Badawi, A.A., 55
- Engvall, H., see Barnett, M.I., 131
- Falch, E., see Bundgaard, H., 19
- Farouk, A.M., see Ali, H.M., 301
- Fouli, A.M., see Badawi, A.A., 55
- Guy, R.H. and Hadgraft, J., A theoretical description relating skin penetration to the thickness of the applied medicament, 321
- Hadgraft, J., see Guy, R.H., 321
- Hanano, M., see Wateri, N., 155
- Hansen, J. and Bundgaard, H., Studies on the stability of corticosteroids V. The degradation pattern of hydrocortisone in aqueous solution, 307
- Heimlich, K.R., see Kwan, K.C., 225
- Hersey, J.A., see Krycer, I., 119
- Hirai, S., see Iga, K., 43
- Hüttenrauch, R., The influence of molecule-size on the state of order and on the order-dependent properties of ointments, 349
- Iga, K., Hirai, S. and Yashiki, T., Assessment of local and systemic availability of tribenoside after rectal administration in rats, 43
- Inotsume, N. and Nakano, M., Reversible azo-methine bond cleavage of 2'-fluoro derivatives of benzodiazepines in acidic solutions at body temperature, 147
- James, K.C., see Armstrong, N.A., 185
- James, K.C., see Armstrong, N.A., 195
- Kaneniwa, N., see Wateri, N., 155
- Kellaway, I.W. and Najib, N.M., The adsorption of hydrophilic polymers at the polystyrene-water interface, 285
- Kinkel, J.F.M. and Tomlinson, E., Drug liquid-liquid distribution based on the fundamentals of segmented flow, 261
- Krycer, I. and Hersey, J.A., Fine power mixing in a vibratory ball mill, 119
- Kutris, A.M., see Theodorakis, M.C., 333
- Kwan, K.C. and Heimlich, K.R., The role of

- pharmacokinetics in drug product design, 225
- Larsen, C., see Bundgaard, H., 19
- Long, D.F., Repta, A.J. and Sternson, L.A., The reactivity of cisplatin in plasma. Implications for sample handling in pharmacokinetic studies, 167
- Macheras, P. and Rosen, A., Some considerations concerning the effect of an increased rate of oral absorption on drug levels of protein-bound drugs, 175
- Marszall, L., Effective hydrophile-lipophile balance and micelle formation of non-ionic surfactants in the presence of short-chain glycol ethers, 253
- McElroy, J.C. and D'Arcy, P.F., Sites and mechanisms of drug interactions II. Protein binding, renal excretion and pharmacodynamic interactions, 205
- Miyazaki, S., Oshiba, M. and Nadai, T., Unusual solubility and dissolution behavior of pharmaceutical hydrochloride salts in chloride-containing media, 77
- Morén, F. and Anderson, J., Fraction of dose exhaled after administration of pressurized inhalation aerosols, 295
- Nadai, T., see Miyazaki, S., 77
- Najib, N.M., see Kellaway, I.W., 285
- Nakano, M., see Inotsume, N., 147
- Nixon, J.R. and Agyilrah, G.A., The influence of colloidal proportions on the release of phenobarbitone from microcapsules, 277
- Nyström, C., see Barnett, M.I., 131
- Oshiba, M., see Miyazaki, S., 77
- Renitakis, N.K., see Theodorakis, M.C., 333
- Repta, A.J., see Long, D.F., 167
- Rogers, J.A. and Wong, A., The temperature dependence and thermodynamics of partitioning of phenols in the n-octanol-water system, 339
- Rosen, A., see Macheras, P., 175
- Rosoff, M. and Serajuddin, A.T.M., Solubilization of diazepam in bile salts and in sodium cholate-lecithin-water phase, 137
- Said, S. and Al-Shora, H., Sustained release from inert matrices I. Effect of microcrystalline cellulose on aminophylline and theophylline release, 11
- Sanchez, M.J., see Sanchez Camazano, M., 243
- Sanchez Camazano, M., Sanchez, M.J., Vicente, M.T. and Dominguez-Gil, A., Adsorption of chlorpheniramine maleate by montmorillonite, 243
- Sarajuddin, A.T.M., see Rosoff, M., 137
- Shefter, E. and Cheng, K.C., Drug-polyvinylpyrrolidone (PVP) dispersions. A differential scanning calorimetric study, 179
- Sokoloski, T.D., Wu, C.-C. and Burkman, A.M., Rapid absorptive loss of nitroglycerin from aqueous solution to plastic, 63
- Sternson, L.A., see Long, D.F., 167
- Strittmatter, Th., see Ali, S.L., 183
- Theodorakis, M.C., Kutris, A.M. and Renitakis, N.K., A study on particulate formation of silicone-coated glass surfaces, 333
- Tomlinson, E., see Kinkel, J.F.M., 261
- Vicente, M.T., see Sanchez Camazano, M., 243
- Wateri, N., Kaneniwa, N. and Hanano, M., Comparison of in vivo dissolution of sulfa drugs calculated by deconvolution to absorption rates calculated by multicompartment model method, 155
- Wong, A., see Rogers, J.A., 339
- Wu, C.-C., see Sokoloski, T.D., 63
- Yashiki, T., see Iga, K., 43
- York, P., Powder failure testing - pharmaceutical applications, 89